

# MAN'S PLACE IN NATURE

T. H. HUXLEY

*The University Library  
Leeds*



*Medical and Dental  
Library*

Stack  
311 312  
1004



30106

004071113

University of Leeds Medical and Dental Library  
DATE DUE FOR RETURN

27 MAR 1996 <del>CANCELLED</del>		

UPS/4442/5/82



# MAN'S PLACE IN NATURE

AND A SUPPLEMENTARY ESSAY

BY

THOMAS HENRY HUXLEY

[ISSUED FOR THE RATIONALIST PRESS ASSOCIATION, LIMITED]

LONDON:

WATTS & CO.,

17, JOHNSON'S COURT, FLEET STREET, E.C.

1908



820066



## PREFATORY NOTE

---

DARWIN closed his *Origin of Species* with the brief but pregnant hint that the theory of natural selection "would throw light on the origin of man and his history." In the *Descent of Man*, published twelve years later, he explained that his reticence in 1859 was due to a desire "not to add to the prejudice against his views." He found a doughty champion in Huxley, who, heedless of prejudices based on ignorance or intellectual indolence, pushed the theory of organic evolution to its logical issue in including man, physically and psychically, in the processes of development.

In the spring of 1861 he gave a course of weekly lectures to working men "On the Relation of Man to the Rest of the Animal Kingdom." This was followed by lectures on the same subject before the Philosophical Institution of Edinburgh. The applause of his Scotch audience was not echoed in the Press. It resented the presentment of "an anti-scriptural and most debasing theory .....standing in blasphemous contradiction to Biblical narrative and doctrine." And this group of adjectives—ever convenient weapons where cogent arguments are lacking—was representative of the well-nigh universal feeling of the time at the audacious demolition of "the barrier between man and the rest of the animal world."

The two sets of lectures formed the basis of the book now re-published. Heedless of the advice of "a very shrewd friend [Sir William Lawrence<sup>1</sup>] not to ruin all his prospects by so rash a venture," Huxley issued the book, under the title of *Evidences as to Man's Place in Nature*, in 1863. He explains that,

inasmuch as Development and Vertebrate Anatomy were not among Darwin's many specialities, it appeared to me that I should not be intruding on the ground he had made his own if I discussed this part of the general question. In fact, I thought that I might probably serve the cause of Evolution by doing so.

<sup>1</sup> Died 1867. Author of *The Physiology, Zoology, and Natural History of Man* (1819), in which "contemporary theologians discerned an attempt to undermine the foundations of religion. Lord Eldon refused an injunction to protect the rights of the author on the ground that the book contradicted the Scriptures" (*Dict. Nat. Biog.*, xxxii., p. 286).

Moreover, he was well qualified sociologically. As he told an audience in 1882, it had been his fate

to be familiar with almost every form of society, from the uncivilised savage of Papua and Australia and the civilised savages of the slums and dens of the poverty-stricken parts of great cities, to those who, perhaps, are occasionally the somewhat over-civilised members of our upper ten thousand.

And when Huxley spoke, with full warrant, of his equipment in Vertebrate Anatomy, he included therein knowledge of the apparatus of the mind as well as of the mechanism of the body. Consequently, his demonstration of the essentially indistinguishable structural likeness between man and his nearest congeners, the four anthropoid apes—the Orang-utan, the Chimpanzee, the Gorilla, and the Gibbon—is followed by the conclusion that

the attempt to draw a psychical distinction is futile, and that even the highest faculties of feeling and of intellect begin to germinate in lower forms of life.

Forty-five years have passed since *Evidences as to Man's Place in Nature* appeared. Not a statement therein has needed any material qualification; biology and psychology bring their "great cloud of witnesses" to its unchallengeable accuracy and abiding value.

EDWARD CLODD.

*February, 1908.*



## CONTENTS

---

	PAGE
I.—ON THE NATURAL HISTORY OF THE MAN-LIKE APES - - - -	7
II.—ON THE RELATIONS OF MAN TO THE LOWER ANIMALS - - - -	39
III.—ON SOME FOSSIL REMAINS OF MAN -	80
IV.—ON THE METHODS AND RESULTS OF ETHNOLOGY [1865] - - - -	104

\* \* The first three essays were published in January, 1863, under the title of "Man's Place in Nature"; the fourth essay originally appeared in the *Fortnightly Review* (1865).



# MAN'S PLACE IN NATURE

## I.

### ON THE NATURAL HISTORY OF THE MAN-LIKE APES

ANCIENT traditions, when tested by the severe processes of modern investigation, commonly enough fade away into mere dreams; but it is singular how often the dream turns out to have been a half-waking one, presaging a reality. Ovid foreshadowed the discoveries of the geologist: the Atlantis was an imagination, but Columbus found a western world; and, though the quaint forms of Centaurs and Satyrs have an existence only in the realms of art, creatures approaching man more nearly than they in essential structure, and yet as thoroughly brutal as the goat's or horse's half of the mythical compound, are now not only known, but notorious.

I have not met with any notice of one of these MAN-LIKE APES of earlier date than that contained in Pigafetta's *Description of the Kingdom of Congo*,<sup>1</sup> drawn up from the notes of a Portuguese sailor, Eduardo Lopez, and published in 1598. The tenth chapter

<sup>1</sup> *Regnum Congo: hoc est, Vera Descriptio Regni Africani quod tam ab incolis quam Lusitanis Congus appellatur*, per Philippum Pigafettam, olim ex Edoardo Lopez acroamatis lingua Italica excerpta, num Latio sermone donata ab August. Cassiod. Reinio. Iconibus et imaginibus rerum memorabilium quasi vivis, opera et industria Joan. Theodori et Joan. Israelis de Bry, fratrum exornata. Francofurti, MDXCVIII.

of this work is entitled "De Animalibus quæ in hac provincia reperiuntur," and contains a brief passage to the effect that "in the Songan country, on the banks of the Zaire, there are multitudes of apes, which afford great delight to the nobles by imitating human gestures." As this might apply to almost any kind of apes, I should have thought little of it, had not the brothers De Bry, whose engravings illustrate the work, thought fit, in their eleventh "Argumentum," to figure two of these "Simiæ magnatum deliciæ." So much of the plate as contains these apes is faithfully copied in the woodcut (Fig. 1), and it will be observed that they are tail-less, long-armed, and large-eared; and about the size of Chimpanzees. It may be that these apes are as much figments of the imagination of the ingenious brothers as the winged, two-legged, crocodile-headed dragon which adorns the same plate; or, on the other hand, it may be that the artists have constructed their drawings from some essentially faithful description of a Gorilla or a Chimpanzee. And, in either case, though these figures are worth a passing notice, the oldest trustworthy and definite accounts of any animal of this kind

date from the seventeenth century, and are due to an Englishman.

The first edition of that most amusing old book, *Purchas his Pilgrimage*, was published in 1613, and therein are to be found many references to the statements of one whom Purchas terms "Andrew Battell (my neere neighbour, dwelling at Leigh in Essex) who served under Manuel Silvera Perera, Governor under the King of Spaine, at his city of Saint Paul, and with him went farre into the countrey of Angola"; and again, "my friend,



FIG. 1.—*Simiæ magnatum deliciæ*.—De Bry, 1598.

Andrew Battle, who lived in the kingdom of Congo many yeares," and who, "upon some quarell betwixt the Portugals (among whom he was a sergeant of a band) and him, lived eight or nine moneths in the woodes." From this weather-beaten old soldier, Purchas was amazed to hear "of a kinde of Great Apes, if they might so bee termed, of the height of a man, but twice as bigge in feature of their limmes, with strength proportionable, hairie all over, otherwise altogether like men and women in their whole

bodily shape.<sup>1</sup> They lived on such wilde fruits as the trees and woods yielded, and in the night time lodged on the trees."

This extract is, however, less detailed and clear in its statements than a passage in the third chapter of the second part of another work—*Purchas his Pilgrimes*, published in 1625, by the same author—which has been often, though hardly ever rightly, cited. The chapter is entitled, "The strange adventures of Andrew Battell, of Leigh in Essex, sent by the Portugals prisoner to Angola, who lived there and in the adioining regions neere eightene yeeres." And the sixth section of this chapter is headed—"Of the Provinces of Bongo, Calongo, Mayombe, Manikesocke, Motimbas: of the Ape Monster Pongo, their hunting: Idolatries; and divers other observations."

This province (Calongo) toward the east bordereth upon Bongo, and toward the north upon Mayombe, which is nineteen leagues from Longo along the coast.

This province of Mayombe is all woods and groves, so overgrowne that a man may travaile twentie days in the shadow without any sunne or heat. Here is no kind of corne nor graine, so that the people liveth onely upon plantanes and roots of sundrie sorts, very good; and nuts; nor any kinde of tame cattell, nor hens.

But they have great store of elephants' flesh, which they greatly esteeme, and many kinds of wild beasts; and great store of fish. Here is a great sandy bay, two leagues to the northward of Cape Negro,<sup>2</sup> which is the port of Mayombe. Sometimes the Portugals lade log-wood in this bay. Here is a great river, called Banna: in the winter it hath no barre,

<sup>1</sup> "Except this that their legges had no calves" (ed. 1626). And in a marginal note, "These great apes are called Pongo's."

<sup>2</sup> *Purchas's note*.—Cape Negro is in sixteen degrees south of the Line.



because the generall winds cause a great sea. But when the sunne hath his south declination, then a boat may goe in ; for then it is smooth because of the raine. This river is very great, and hath many islands and people dwelling in them. The woods are so covered with baboones, monkees, apes and parrots, that it will feare any man to travaile in them alone. Here are also two kinds of monsters, which are common in these woods, and very dangerous.

The greatest of these two monsters is called Pongo in their language, and the lesser is called Engeco. This Pongo is in all proportion like a man ; but that he is more like a giant in stature than a man ; for he is very tall, and hath a man's face, hollow-eyed, with long haire upon his browes. His face and eares are without haire, and his hands also. His bodie is full of haire, but not very thicke ; and it is of a dunnish colour.

He differeth not from a man but in his legs ; for they have no calfe. Hee goeth alwaies upon his legs, and carrieth his hands clasped in the nape of his necke when he goeth upon the ground. They sleepe in the trees, and build shelters for the raine. They feed upon fruit that they find in the woods, and upon nuts, for they eate no kind of flesh. They cannot speake, and have no understanding more than a beast. The people of the countrie, when they travaile in the woods, make fires where they sleepe in the night ; and in the morning when they are gone, the Pongoes will come and sit about the fire till it goeth out ; for they have no understanding to lay the wood together. They goe many together and kill many negroes that travaile in the woods. Many times they fall upon the elephants which come to feed where they be, and so beate them with their clubbed fists, and pieces of wood, that they will runne roaring away from them. Those Pongoes are never taken alive because they are so strong that ten men cannot hold one of them ; but yet they take many of their young ones with poisoned arrowes.

The young Pongo hangeth on his mother's belly with his hands fast clasped about her ; so that when the countrie people kill any of the females they take the young one, which hangeth fast upon his mother.

When they die among themselves, they cover the dead with great heaps of

boughs and wood, which is commonly found in the forest.<sup>1</sup>

It does not appear difficult to identify the exact region of which Battell speaks. Longo is doubtless the name of the place usually spelled "Loango" on our maps. Mayombe still lies some nineteen leagues northward from Loango, along the coast ; and Cilongo or Kilonga, Manikesocke, and Motimbas are yet registered by géographers. The Cape Negro of Battell, however, cannot be the modern Cape Negro in 16° S., since Loango itself is in 4° S. latitude. On the other hand, the "great river called Banna" corresponds very well with the "Camma" and "Fernand Vas" of modern geographers, which form a great delta on this part of the African coast.

Now, this "Camma" country is situated about a degree and a half south of the Equator ; while a few miles to the north of the Line lies the Gaboon, and a degree or so north of that the Money River—both well known to modern naturalists as localities where the largest of man-like Apes has been obtained. Moreover, at the present day, the word "Engeco," or N'schego, is applied by the natives of these regions to the smaller of the two great Apes which inhabit

<sup>1</sup> *Purchas's marginal note*, p. 982 :—"The Pongo is a giant ape. He told me in conference with him that one of these Pongoes tooke a negro boy of his which lived a moneth with them. For they hurt not those which they surprise at unawares, except they look on them, which he avoyded. He said their highth was like a man's, but their bignes twice as great. I saw the negro boy. What the other monster should be he hath forgotten to relate ; and these papers came to my hand since his death, which, otherwise, in my often conferences, I might have learned. Perhaps he meaneth the pigmy Pongo killers mentioned."



them; so that there can be no rational doubt that Andrew Battell spoke of that which he knew of his own knowledge, or, at any rate, by immediate report from the natives of Western Africa. The "Engeco," however, is that "other monster," whose nature Battell "forgot to relate"; while the name "Pongo" — applied to the animal whose characters and habits are so fully and carefully described — seems to have died out, at least

*Homo Sylvestris.*  
*Orang Outang.*



FIG. 2.—The Orang of Tulpus, 1641.

in its primitive form and significance. Indeed, there is evidence that not only in Battell's time, but up to a very recent date, it was used in a totally different sense from that in which he employs it.

For example, the second chapter of Purchas's work, which I have just quoted, contains "A Description and Historicall Declaration of the Golden Kingdom of Guinea, &c., &c. Translated from the Dutch, and compared also with the Latin," wherein it is stated (p. 986) that

the River Gaboon lyeth about fifteen miles northward from Rio de Angra, and eight miles northward from Cape de Lope Gonsalvez (Cape Lopez), and is right under the Equinoctial line, about fiftene miles from St. Thomas, and is a great land, well and easily to be knowne. At the mouth of the river there lieth a sand, three or foure fathoms deepe, whereon it beateth mightily with the streame which runneth out of the river into the sea. This river, in the mouth thereof, is at least four miles broad; but, when you are about the Iland called *Pongo*, it is not above two miles broad.....On both sides the river there standeth many trees..... The Iland called *Pongo*, which hath a monstrous high hill.

The French naval officers, whose letters are appended to the late M. Isidore Geoff. Saint Hilaire's excellent essay on the Gorilla,<sup>1</sup> note in similar terms the width of the Gaboon, the trees that line its banks down to the water's edge, and the strong current that sets out of it. They describe two islands in its estuary—one low, called Perroquet; the other high, presenting three conical hills, called Coniquet; and one of them, M. Franquet, expressly states that formerly the Chief of Coniquet was called *Meni-Pongo*, meaning thereby Lord of *Pongo*; and that the *N'Pongues* (as, in agreement with Dr. Savage, he affirms the natives call themselves) term the estuary of the Gaboon itself *N'Pongo*.

It is so easy, in dealing with savages, to misunderstand their applications of words to things that one is at first inclined to suspect Battell of having confounded the name of this region, where his "greater monster" still abounds, with the name of the animal itself. But he is so right about other matters (including the name of the "lesser monster") that one is loth to suspect the old traveller

<sup>1</sup> *Archives du Museum*, tome x.

of error; and, on the other hand, we shall find that a voyager of a hundred years' later date speaks of the name "Boggoe" as applied to a great Ape by the inhabitants of quite another part of Africa—Sierra Leone.

But I must leave this question to be settled by philologists and travelers; and I should hardly have dwelt so long upon it except for the curious part played by this word "*Pongo*" in the later history of the man-like Apes.

The generation which succeeded Battell saw the first of the man-like Apes which was ever brought to Europe, or, at any rate, whose visit found a historian. In the third book of Tulpus's *Observationes Medicæ*, published in 1641, the 56th chapter or section is devoted to what he calls *Satyrus indicus*, "called by the Indians Orang-outang, or Man-of-the-Woods, and by the Africans Quoias Morrou." He gives a very good figure, evidently from the life, of the specimen of this animal, "*nostra memoria ex Angolâ delatum*," presented to Frederick Henry Prince of Orange. Tulpus says it was as big as a child of three years old, and as stout as one of six years; and that its back was covered with black hair. It is plainly a young Chimpanzee.

In the meanwhile, the existence of other, Asiatic, man-like Apes became known, but at first in a very mythical fashion. Thus Bontius (1658) gives an altogether fabulous and ridiculous account and figure of an animal which he calls "Orang-outang"; and, though he says "*vidi Ego cujus effigiem hic exhibeo*," the said effigies (see Fig. 6 for Hoppius's copy of it) is nothing but a very hairy woman of rather comely aspect, and with proportions

and feet wholly human. The judicious English anatomist, Tyson, was justified in saying of this description by Bontius: "I confess I do mistrust the whole representation."

It is to the last-mentioned writer and his coadjutor Cowper that we owe the first account of a man-like ape which has any pretensions to scientific accuracy and completeness. The treatise *Orang-outang, sive Homo Sylvestris; or, the Anatomy of a Pygmie Compared with that of a Monkey, an Ape, and a Man*, published by the Royal Society in 1699, is, indeed, a work of remarkable merit, and has, in some respects, served as a model to subsequent inquirers. This "Pygmie," Tyson tells us, "was brought from Angola, in Africa, but was first taken a great deal higher up the country." Its hair "was of a coal-black colour, and strait"; and, "when it went as a quadruped on all four, 'twas awkwardly, not placing the palm of the hand flat to the ground; but it walk'd upon its knuckles, as I observed it to do when weak and had not strength enough to support its body." "From the top of the head to the heel of the foot, in a strait line, it measured twenty-six inches."

These characters, even without Tyson's good figures (Figs. 3 and 4), would have been sufficient to prove his "Pygmie" to be a young Chimpanzee. But the opportunity of examining the skeleton of the very animal Tyson anatomised having most unexpectedly presented itself to me, I am able to bear independent testimony to its being a veritable *Trogodytes niger*,<sup>1</sup> though still very young.

<sup>1</sup> I am indebted to Dr. Wright, of Cheltenham.



Although fully appreciating the resemblances between his Pygmie and Man, Tyson by no means overlooked the differences between the two; and he concludes his memoir by summing up, first, the points in which "the Ourang-outang or Pygmie more resembled a Man than Apes and Monkeys do," under forty-seven distinct heads; and then giving, in thirty-

resembled more the Ape and Monkey kind."

After a careful survey of the literature of the subject extant in his time, our author arrives at the conclusion that his "Pygmie" is identical neither with the Orangs of Tulpius and Bon-tius, nor with the Quoias Morrou of Dapper (or rather of Tulpius), the Barris of d'Arcos, nor with the Pongo



FIG. 3.—The "Pygmie," reduced from Tyson's figure 1, 1699.



FIG. 4.—The "Pygmie," reduced from Tyson's figure 2, 1699.

four similar brief paragraphs, the respects in which "the Ourang-outang or Pygmie differ'd from a man, and

ham whose paleontological labours are so well known, for bringing this interesting relic to my knowledge. Tyson's granddaughter, it appears, married Dr. Allardyce, a physician of repute in Cheltenham, and brought, as part of her dowry, the skeleton of the "Pygmie." Dr. Allardyce presented it to the Cheltenham Museum; and, through the good offices of my friend Dr. Wright, the authorities of the museum have permitted me to borrow what is, perhaps, its most remarkable ornament.

of Battell; but that it is a species of ape probably identical with the Pygmies of the ancients; and, says Tyson, though it "does so much resemble a *Man* in many of its parts, more than any of the ape kind, or any other *animal* in the world that I know of, yet by no means do I look upon it as the product of a *mixt* generation—'tis a *Brute-Animal sui generis*, and a particular *species of Ape*."

The name of "Chimpanzee," by

which one of the African Apes is now so well known, appears to have come into use in the first half of the eighteenth century; but the only important addition made, in that period, to our acquaintance with the man-like apes of Africa is contained in *A New Voyage to Guinea*, by William Smith, which bears the date 1744.

In describing the animals of Sierra Leone, p. 51, this writer says:—

I shall next describe a strange sort of animal, called by the white men in this

before, neither can those who call them so tell, except it be for their near resemblance of a human creature, though nothing at all like an Ape. Their bodies, when full grown, are as big in circumference as a middle-sized man's—their legs much shorter, and their feet larger; their arms and hands in proportion. The head is monstrously big, and the face broad and flat, without any other hair but the eyebrows; the nose very small, the mouth wide, and the lips thin. The face, which is covered by a white skin, is monstrously ugly, being all over wrinkled as with old age; the teeth broad and yellow; the hands have no more hair than the face, but the same white skin, though all the rest of



FIG. 5.—Facsimile of William Smith's figure of the "Mandrill," 1744.

country Mandrill;<sup>1</sup> but why it is so called I know not, nor did I ever hear the name

<sup>1</sup> "Mandrill" seems to signify a "man-like ape," the word "drill" or "dril" having been anciently employed in England to denote an ape or baboon. Thus, in the fifth edition of Blount's *Glossographia; or a Dictionary interpreting the hard words of whatsoever language now used in our refined English tongue.....very useful for all such as desire to understand what they read*, published in 1681, I find "Dril—a stone-cutter's tool, wherewith he bores little holes in marble, etc. Also a large overgrown Ape and Baboon, so called." "Drill" is used in the same sense in Charleton's *Onomasticon Zoicon*, 1668. The singular etymology of the word given by Buffon seems hardly a probable one.

the body is covered with long black hair, like a bear. They never go upon all fours, like apes; but cry, when vexed or teased, just like children.....

When I was at Sherbro, one Mr. Cummerbus, whom I shall have occasion hereafter to mention, made me a present of one of these strange animals, which are called by the natives 'Boggoe': it was a she-cub, of six months' age, but even then larger than a Baboon. I gave it in charge to one of the slaves, who knew how to feed and nurse it, being a very tender sort of animal; but whenever I went off the deck the sailors began to tease it—some loved to see its tears and hear it cry; others hated its snotty-nose; one who hurt it, being checked by the negro that took care of it, told the slave



he was very fond of his country-woman, and asked him if he should not like her for a wife? To which the slave very readily replied, "No, this no my wife; this a white woman—this fit wife for you." This unlucky wit of the negro's, I fancy, hastened its death, for next morning it was found dead under the windlass.

William Smith's "Mandrill" or "Boggoe," as his description and figure testify, was, without doubt, a Chimpanzee.

Linnæus knew nothing, of his own observation, of the man-like Apes of either Africa or Asia, but a dissertation by his pupil Hoppius in the *Amæ-*

believed; for in the standard edition of the *Systema Naturæ* it is enumerated as a second species of Homo, "H. nocturnus." *Lucifer Aldrovandi* is a copy of a figure in Aldrovandus, *De Quadrupedibus digitatis viviparis*, Lib. 2, p. 249 (1645), entitled "Cercopithecus formæ raræ *Barbilius* vocatus et originem a China ducebat." Hoppius is of opinion that this may be one of that cat-tailed people, of whom Nicolaus Köping affirms that they eat a boat's crew, "gubernator navis" and all! In the *Systema Naturæ* Linnæus calls it, in a note, *Homo cau-*




FIG. 6.—The Anthropomorpha of Linnæus.

*nitates Academicæ* (VI. "Anthropomorpha") may be regarded as embodying his views respecting these animals.

The dissertation is illustrated by a plate, of which the accompanying woodcut, Fig. 6, is a reduced copy. The figures are entitled (from left to right) 1. *Troglodyta Bontii*; 2. *Lucifer Aldrovandi*; 3. *Satyrus Tulpæ*; 4. *Pygmæus Edwardi*. The first is a bad copy of Bontius' fictitious "Ourang-outang," in whose existence, however, Linnæus appears to have fully

*dated*, and seems inclined to regard it as a third species of man. According to Temminck, *Satyrus Tulpæ* is a copy of the figure of a Chimpanzee published by Scotin in 1738, which I have not seen. It is the *Satyrus indicus* of the *Systema Naturæ*, and is regarded by Linnæus as possibly a distinct species from *Satyrus sylvestris*. The last, named *Pygmæus Edwardi*, is copied from the figure of a young "Man of the Woods," or true Orang-Utan, given in Edwards's *Gleanings of Natural History* (1758).



Buffon was more fortunate than his great rival. Not only had he the rare opportunity of examining a young Chimpanzee in the living state, but he became possessed of an adult Asiatic man-like Ape—the first and the last adult specimen of any of these animals brought to Europe for many years. With the valuable assistance of Daubenton, Buffon gave an excellent description of this creature, which, from its singular proportions, he termed the long-armed Ape, or Gibbon. It is the modern *Hylobates lar*.

Thus when, in 1766, Buffon wrote the fourteenth volume of his great work, he was personally familiar with the young of one kind of African man-like Ape, and with the adult of an Asiatic species; while the Orang-Utan and the Mandrill of Smith were known to him by report. Furthermore, the Abbé Prevost had translated a good deal of Purchas's *Pilgrimes* into French, in his *Histoire générale des Voyages* (1748), and there Buffon found a version of Andrew Battell's account of the Pongo and the Engeco. All these data Buffon attempts to weld together into harmony in his chapter entitled "Les Orang-outangs ou le Pongo et le Jocko." To this title the following note is appended:—

Orang-outang, nom de cet animal aux Indes orientales: Pongo, nom de cet animal à Lowando, Province de Congo.

Jocko, Enjocko, nom de cet animal à Congo que nous avons adopté. *En* est l'article, que nous avons retranché.

Thus it was that Andrew Battell's "Engeco" became metamorphosed into "Jocko," and, in the latter shape, was spread all over the world, in consequence of the extensive popularity of Buffon's works. The Abbé Prevost

and Buffon between them, however, did a good deal more disfigurement to Battell's sober account than "cutting off an article." Thus Battell's statement that the Pongos "cannot speake, and have no understanding more than a beast," is rendered by Buffon "*qu'il ne peut parler quoiqu'il ait plus d'entendement que les autres animaux*"; and again, Purchas's affirmation, "He told me, in conference with him, that one of these Pongos tooke a negro boy of his which lived a moneth with them," stands in the French version, "*un pongo lui enleva un petit nègre qui passa un an entier dans la société de ces animaux.*"

After quoting the account of the great Pongo, Buffon justly remarks that all the "Jockos" and "Orangs" hitherto brought to Europe were young; and he suggests that, in their adult condition, they might be as big as the Pongo or "great Orang"; so that, provisionally, he regarded the Jockos, Orangs, and Pongos as all of one species. And perhaps this was as much as the state of knowledge at the time warranted. But how it came about that Buffon failed to perceive the similarity of Smith's "Mandrill" to his own "Jocko," and confounded the former with so totally different a creature as the blue-faced Baboon, is not so easily intelligible.

Twenty years later Buffon changed his opinion,<sup>1</sup> and expressed his belief that the Orangs constituted a genus with two species—a large one, the Pongo of Battell, and a small one, the Jocko; that the small one (Jocko) is the East Indian Orang; and that

<sup>1</sup> *Histoire Naturelle*, Suppl. tome 7ème, 1789.

the young animals from Africa, observed by himself and Tulpius, are simply young Pongos.

In the meanwhile, the Dutch naturalist Vosmaer gave, in 1778, a very good account and figure of a young Orang, brought alive to Holland; and his countryman, the famous anatomist Peter Camper, published (1779) an essay on the Orang-Utan of similar value to that of Tyson on the Chimpanzee. He dissected several females and a male, all of which, from the state of their skeleton and their dentition, he justly supposes to have been young. However, judging by the analogy of man, he concludes that they could not have exceeded four feet in height in the adult condition. Furthermore, he is very clear as to the specific distinctness of the true East Indian Orang.

"The Orang," says he, "differs not only from the Pigmy of Tyson and from the Orang of Tulpius by its peculiar colour and its long toes, but also by its whole external form. Its arms, its hands, and its feet are longer, while the thumbs, on the contrary, are much shorter, and the great toes much smaller in proportion."<sup>1</sup> And again: "The true Orang—that is to say, that of Asia, that of Borneo—is consequently not the Pithecus, or tailless Ape, which the Greeks, and especially Galen, have described. It is neither the Pongo nor the Jocko, nor the Orang of Tulpius, nor the Pigmy of Tyson—it is an animal of a peculiar species, as I shall prove in the clearest manner by the organs of voice and the skeleton in the following chapters" (l. c. p. 64).

A few years later, M. Radermacher, who held a high office in the Government of the Dutch dominions in India, and was an active member of the Batavian Society of Arts and Sciences, published in the second part of the Transactions of that Society<sup>1</sup> a Description of the Island of Borneo, which was written between the years 1779 and 1781, and, among much other interesting matter, contains some notes upon the Orang. The small sort of Orang-Utan—viz., that of Vosmaer and of Edwards—he says, is found only in Borneo, and chiefly about Banjermassing, Mampauwa, and Landak. Of these he had seen some fifty during his residence in the Indies; but none exceeded two and a half feet in length. The larger sort, often regarded as a chimæra, continues Radermacher, would perhaps long have remained so, had it not been for the exertions of the Resident at Rembang, M. Palm, who, on returning from Landak towards Pontiana, shot one, and forwarded it to Batavia in spirit, for transmission to Europe.

Palm's letter describing the capture runs thus:—"Herewith I send your Excellency, contrary to all expectation (since long ago I offered more than a hundred ducats to the natives for an Orang-Utan of four or five feet high) an Orang which I heard of this morning about eight o'clock. For a long time we did our best to take the frightful beast alive in the dense forest about half way to Landak. We forgot even to eat, so anxious were we not to let him escape; but it was necessary to take care that he did not revenge

<sup>1</sup> *Verhandelingen van het Bataviaasch Genootschap. Tweede Deel. Derde Druk. 1826.*

<sup>1</sup> Camper, *Œuvres*, i., p. 56.



himself, as he kept continually breaking off heavy pieces of wood and green branches, and dashing them at us. This game lasted till four o'clock in the afternoon, when we determined to shoot him; in which I succeeded very well, and indeed better than I ever shot from a boat before; for the bullet went just into the side of his chest, so that he was not much damaged. We got him into the prow still living, and bound him fast, and next morning he died of his wounds. All Pontiana came on board to see him when we arrived." Palm gives

states, in a letter dated Batavia, February 18th, 1781,<sup>1</sup> that the specimen was sent to Europe in brandy to be placed in the collection of the Prince of Orange; "unfortunately," he continues, "we hear that the ship has been wrecked." Von Wurm died in the course of the year 1781, the letter in which this passage occurs being the last he wrote; but in his posthumous papers, published in the fourth part of the Transactions of the Batavian Society, there is a brief description, with measurements, of a female Pongo four feet high.



FIG. 7.—The Pongo Skull, sent by Radermacher to Camper, after Camper's original sketches, as reproduced by Lucæ.

his height from the head to the heel as forty-nine inches.

A very intelligent German officer, Baron Von Wurm, who at this time held a post in the Dutch East India service, and was Secretary of the Batavian Society, studied this animal, and his careful description of it, entitled "*Beschrijving van der Groote Borneosche Orang-outang, of de Oost-Indische Pongo*," is contained in the same volume of the Batavian Society's Transactions. After Von Wurm had drawn up his description he

Did either of these original specimens, on which Von Wurm's descriptions are based, ever reach Europe? It is commonly supposed that they did; but I doubt the fact. For, appended to the memoir "*De l'Ourang-outang*," in the collected edition of Camper's works, tome i., pp. 64-66, is a note by Camper himself, referring to Von Wurm's papers, and continuing thus:—"Heretofore this

<sup>1</sup> *Briefe des Herrn v. Wurm und des H. Baron von Wollzogen.* Gotha, 1794.

kind of ape had never been known in Europe. Radermacher has had the kindness to send me the skull of one of these animals, which measured fifty-three inches, or four feet five inches, in height. I have sent some sketches of it to M. Soemmering at Mayence, which are better calculated, however, to give an idea of the form than of the real size of the parts."

These sketches have been reproduced by Fischer and by Lucæ, and bear date 1783, Soemmering having received them in 1784. Had either of Von Wurmb's specimens reached Holland, they would hardly have been unknown at this time to Camper, who, however, goes on to say:—"It appears that since this some more of these monsters have been captured, for an entire skeleton, very badly set up, which had been sent to the Museum of the Prince of Orange, and which I saw only on the 27th of June, 1784, was more than four feet high. I examined this skeleton again on the 19th December, 1785, after it had been excellently put to rights by the ingenious Onymus."

It appears evident, then, that this skeleton, which is doubtless that which has always gone by the name of Wurmb's Pongo, is not that of the animal described by him, though unquestionably similar in all essential points.

Camper proceeds to note some of the most important features of this skeleton; promises to describe it in detail by-and-by; and is evidently in doubt as to the relation of this great "Pongo" to his "petit Orang."

The promised further investigations were never carried out; and so it happened that the Pongo of Von

Wurmb took its place by the side of the Chimpanzee, Gibbon, and Orang as a fourth and colossal species of man-like Ape. And indeed nothing could look much less like the Chimpanzees or the Orangs then known than the Pongo; for all the specimens of Chimpanzee and Orang which had been observed were small of stature, singularly human in aspect, gentle and docile; while Wurmb's Pongo was a monster almost twice their size, of vast strength and fierceness, and very brutal in expression; its great projecting muzzle, armed with strong teeth, being further disfigured by the outgrowth of the cheeks into fleshy lobes.

Eventually, in accordance with the usual marauding habits of the Revolutionary armies, the "Pongo" skeleton was carried away from Holland into France, and notices of it, expressly intended to demonstrate its entire distinctness from the Orang and its affinity with the Baboons, were given, in 1798, by Geoffroy St. Hilaire and Cuvier.

Even in Cuvier's *Tableau Elementaire*, and in the first edition of his great work, the *Règne Animal*, the "Pongo" is classed as a species of Baboon. However, so early as 1818 it appears that Cuvier saw reason to alter this opinion, and to adopt the view suggested several years before by Blumenbach,<sup>1</sup> and after him by Tilesius, that the Bornean Pongo is simply an adult Orang. In 1824 Rudolphi demonstrated, by the con-

<sup>1</sup> See Blumenbach, *Abbildungen Naturhistorischen Gegenstände*, No. 12, 1810; and Tilesius, *Naturhistorische Früchte der ersten Kaiserlich-Russischen Erdumseglung*, p. 115, 1813.



dition of the dentition, more fully and completely than had been done by his predecessors, that the Orangs described up to that time were all young animals, and that the skull and teeth of the adult would probably be such as those seen in the Pongo of Wurmb. In the second edition of the *Régne Animal* (1829) Cuvier infers, from the "proportions of all the parts" and "the arrangements of the foramina and sutures of the head," that the Pongo is the adult of the Orang-Utan, "at least of a very closely allied species," and this conclusion was eventually placed beyond all doubt by Professor Owen's Memoir published in the *Zoological Transactions* for 1835, and by Temminck in his *Monographies de Mammalogie*. Temminck's memoir is remarkable for the completeness of the evidence which it affords as to the modification which the form of the Orang undergoes according to age and sex. Tiedemann first published an account of the brain of the young Orang, while Sandifort, Müller, and Schlegel described the muscles and the viscera of the adult, and gave the earliest detailed and trustworthy history of the habits of the great Indian Ape in a state of nature; and as important additions have been made by later observers, we are at this moment better acquainted with the adult of the Orang-Utan than with that of any of the other greater man-like Apes.

It is certainly the Pongo of Wurmb;<sup>1</sup> and it is as certainly not the Pongo of Battell, seeing that the Orang-Utan is

entirely confined to the great Asiatic islands of Borneo and Sumatra.

And, while the progress of discovery thus cleared up the history of the Orang, it also became established that the only other man-like Apes in the eastern world were the various species of Gibbon—Apes of smaller stature, and therefore attracting less attention than the Orangs, though they are spread over a much wider range of country, and are hence more accessible to observation.

Although the geographical area inhabited by the "Pongo" and "Engeco" of Battell is so much nearer to Europe than that in which the Orang and Gibbon are found, our acquaintance with the African Apes has been of slower growth; indeed, it is only within the last few years that the truthful story of the old English adventurer has been rendered fully intelligible. It was not until 1835 that the skeleton of the adult Chimpanzee became known, by the publication of Professor Owen's above-mentioned very excellent memoir, "On the Osteology of the Chimpanzee and Orang," in the *Zoological Transactions*—a memoir which, by the accuracy of its descriptions, the carefulness of its comparisons, and the excellence of its figures, made an epoch in the history of our knowledge of the bony framework, not only of the Chimpanzee, but of all the anthropoid Apes.

By the investigations herein detailed, it became evident that the old Chimpanzee acquired a size and aspect as different from those of the young known to Tyson, to Buffon, and to Traill, as those of the old Orang from

<sup>1</sup> Speaking broadly, and without prejudice to the question whether there be more than one species of Orang.



the young Orang; and the subsequent very important researches of Messrs. Savage and Wyman, the American missionary and anatomist, have not only confirmed this conclusion, but have added many new details.<sup>1</sup>

One of the most interesting among the many valuable discoveries made by Dr. Thomas Savage is the fact that the natives in the Gaboon country at the present day apply to the Chimpanzee a name—"Enché-eko"—which is obviously identical with the "Engeko" of Battell; a discovery which has been confirmed by all later inquirers. Battell's "lesser monster" being thus proved to be a veritable existence, of course a strong presumption arose that his "greater monster," the "Pongo," would sooner or later be discovered. And, indeed, a modern traveller, Bowdich, had, in 1819, found strong evidence, among the natives, of the existence of a second great Ape, called the "Ingena," "five feet high, and four across the shoulders," the builder of a rude house, on the outside of which it slept.

In 1847 Dr. Savage had the good fortune to make another and most important addition to our knowledge of the man-like Apes; for, being unexpectedly detained at the Gaboon river, he saw in the house of the Rev. Mr. Wilson, a missionary resident there, "a skull represented by the natives to be a monkey-like animal,

remarkable for its size, ferocity, and habits." From the contour of the skull, and the information derived from several intelligent natives, "I was induced," says Dr. Savage (using the term Orang in its old general sense), "to believe that it belonged to a new species of Orang. I expressed this opinion to Mr. Wilson, with a desire for further investigation; and, if possible, to decide the point by the inspection of a specimen, alive or dead." The result of the combined exertions of Messrs. Savage and Wilson was not only the obtaining of a very full account of the habits of this new creature, but a still more important service to science, the enabling the excellent American anatomist already mentioned, Professor Wyman, to describe, from ample materials, the distinctive osteological characters of the new form. This animal was called by the natives of the Gaboon "Engé-ena," a name obviously identical with the "Ingena" of Bowdich; and Dr. Savage arrived at the conviction that this last discovered of all the great Apes was the long-sought "Pongo" of Battell.

The justice of this conclusion, indeed, is beyond doubt; for not only does the "Engé-ena" agree with Battell's "greater monster" in its hollow eyes, its great stature, and its dun or iron-grey colour, but the only other man-like Ape which inhabits these latitudes—the Chimpanzee—is at once identified, by its smaller size, as the "lesser monster," and is excluded from any possibility of being the "Pongo" by the fact that it is black and not dun, to say nothing of the important circumstance already mentioned, that it still retains the

<sup>1</sup> See "Observations on the external characters and habits of the *Troglodytes niger*, by Thomas N. Savage, M.D., and on its organisation, by Jeffries Wyman, M.D.," *Boston Journal of Natural History*, vol. iv., 1843-4; and "External characters, habits, and osteology of *Troglodytes Gorilla*," by the same authors, *ibid*, vol. v., 1847.

name of "Engeko," or "Enché-eko," by which Battell knew it.

In seeking for a specific name for the "Engé-ena," however, Dr. Savage wisely avoided the much-misused "Pongo"; but, finding in the ancient *Periplus* of Hanno the word "Gorilla" applied to certain hairy savage people, discovered by the Carthaginian voyager in an island on the African coast, he attached the specific name *Gorilla* to his new ape, whence arises its present well-known appellation. But Dr. Savage, more cautious than some of his successors, by no means identifies his ape with Hanno's "wild men." He merely says that the latter were "probably one of the species of the Orang"; and I quite agree with M. Brullé that there is no ground for identifying the modern "Gorilla" with that of the Carthaginian admiral.

Since the memoir of Savage and Wyman was published the skeleton of the Gorilla has been investigated by Professor Owen and by the late Professor Duvernoy, of the Jardin des Plantes, the latter having further supplied a valuable account of the muscular system and of many of the other soft parts; while African missionaries and travellers have confirmed and expanded the account originally given of the habits of this great man-like Ape, which has had the singular fortune of being the first to be made known to the general world and the last to be scientifically investigated.

Two centuries and a half have passed away since Battell told his stories about the "greater" and the "lesser monsters" to Purchas, and it has taken nearly that time to arrive at the clear result that there are four

distinct kinds of Anthropoids—in Eastern Asia, the Gibbons and the Orangs; in Western Africa, the Chimpanzees and the Gorilla.

The man-like Apes, the history of whose discovery has just been detailed, have certain characters of structure and of distribution in common. Thus they all have the same number of teeth as man—possessing four incisors, two canines, four false molars, and six true molars in each jaw, or thirty-two teeth in all, in the adult condition; while the milk dentition consists of twenty teeth—or four incisors, two canines, and four molars in each jaw. They are what are called catarrhine Apes—that is, their nostrils have a narrow partition and look downwards; and, furthermore, their arms are always longer than their legs, the difference being sometimes greater and sometimes less; so that, if the four were arranged in the order of the length of their arms in proportion to that of their legs, we should have this series—Orang ( $1\frac{1}{3}$ —1), Gibbon ( $1\frac{1}{4}$ —1), Gorilla ( $1\frac{1}{6}$ —1), Chimpanzee ( $1\frac{1}{10}$ —1). In all, the fore limbs are terminated by hands, provided with longer or shorter thumbs; while the great toe of the foot, always smaller than in Man, is far more movable than in him, and can be opposed, like a thumb, to the rest of the foot. None of these apes have tails, and none of them possess the cheek-pouches common among monkeys. Finally, they are all inhabitants of the Old World.

The Gibbons are the smallest, slenderest, and longest-limbed of the man-like Apes: their arms are longer in proportion to their bodies than

those of any of the other man-like Apes, so that they can touch the ground when erect; their hands are longer than their feet, and they are the only Anthropoids which possess callosities like the lower monkeys. They are variously coloured. The Orangs have arms which reach to the ankles in the erect position of the animal; their thumbs and great toes are very short, and their feet are longer than their hands. They are covered with reddish-brown hair, and the sides of the face, in adult males, are commonly produced into two crescentic, flexible excrescences, like fatty tumours. The Chimpanzees have arms which reach below the knees; they have large thumbs and great toes, their hands are longer than their feet, and their hair is black, while the skin of the face is pale. The Gorilla, lastly, has arms which reach to the middle of the leg, large thumbs and great toes, feet longer than the hands, a black face, and dark grey or dun hair.

For the purpose which I have at present in view, it is unnecessary that I should enter into any further minutiae respecting the distinctive characters of the genera and species into which these man-like Apes are divided by naturalists. Suffice it to say, that the Orangs and the Gibbons constitute the distinct genera, *Simia* and *Hylobates*; while the Chimpanzees and Gorillas are by some regarded simply as distinct species of one genus, *Troglodytes*; by others as distinct genera — *Troglodytes* being reserved for the Chimpanzees, and *Gorilla* for the Engé-ena or Pongo.

Sound knowledge respecting the

habits and mode of life of the man-like Apes has been even more difficult of attainment than correct information regarding their structure.

Once in a generation, a Wallace may be found physically, mentally, and morally qualified to wander unscathed through the tropical wilds of America and of Asia; to form magnificent collections as he wanders; and withal to think out sagaciously the conclusions suggested by his collections; but, to the ordinary explorer or collector, the dense forests of equatorial Asia and Africa, which constitute the favourite habitation of the Orang, the Chimpanzee, and the Gorilla, present difficulties of no ordinary magnitude; and the man who risks his life by even a short visit to the malarious shores of those regions may well be excused if he shrinks from facing the dangers of the interior; if he contents himself with stimulating the industry of the better seasoned natives, and collecting and collating the more or less mythical reports and traditions with which they are too ready to supply him.

In such a manner most of the earlier accounts of the habits of the man-like Apes originated; and even now a good deal of what passes current must be admitted to have no very safe foundation. The best information we possess is that, based almost wholly on direct European testimony, respecting the Gibbons; the next best evidence relates to the Orangs; while our knowledge of the habits of the Chimpanzee and the Gorilla stands much in need of support and enlargement by additional testimony from instructed European eye-witnesses.



It will therefore be convenient, in endeavouring to form a notion of what we are justified in believing about these animals, to commence with the best known man-like Apes, the Gibbons and Orangs; and to make use of the perfectly reliable information respecting them as a sort of criterion of the probable truth or falsehood of assertions respecting the others.

Of the GIBBONS, half a dozen species are found scattered over the Asiatic islands, Java, Sumatra, Borneo, and through Malacca, Siam, Arracan, and an uncertain extent of Hindostan, on the mainland of Asia. The largest attain a few inches above three feet in height, from the crown to the heel, so that they are shorter than the other man-like Apes; while the slenderness of their bodies renders their mass far smaller in proportion even to this diminished height.

Dr. Salomon Müller, an accomplished Dutch naturalist, who lived for many years in the Eastern Archipelago, and to the results of whose personal experience I shall frequently have occasion to refer, states that the Gibbons are true mountaineers, loving the slopes and edges of the hills, though they rarely ascend beyond the limit of the fig-trees. All day long they haunt the tops of the tall trees; and though, towards even-

ing, they descend in small troupes to the open ground, no sooner do they spy a man than they dart up the hill-sides, and disappear in the darker valleys.



FIG. 8.—A Gibbon (*H. Pileatus*), after Wolff.

All observers testify to the prodigious volume of voice possessed by these animals. According to the writer whom I have just cited, in one of them, the Siamang, "the voice is

grave and penetrating, resembling the sounds gōek, gōek, gōek, gōek, goek ha ha ha haaāā, and may easily be heard at a distance of half a league." While the cry is being uttered, the great membranous bag under the throat which communicates with the organ of voice, the so-called "laryngeal sac," becomes greatly distended, diminishing again when the creature relapses into silence.

M. Duvaucel likewise affirms that the cry of the Siamang may be heard for miles—making the woods ring again. So Mr. Martin<sup>1</sup> describes the cry of the agile Gibbon as "overpowering and deafening" in a room, and, "from its strength, well calculated for resounding through the vast forests." Mr. Waterhouse, an accomplished musician as well as zoologist, says: "The Gibbon's voice is certainly much more powerful than that of any singer I ever heard." And yet it is to be recollected that this animal is not half the height of, and far less bulky in proportion than, a man.

There is good testimony that various species of Gibbon readily take to the erect posture. Mr. George Bennett,<sup>2</sup> a very excellent observer, in describing the habits of a male *Hylobates syndactylus* which remained for some time in his possession, says: "He invariably walks in the erect posture when on a level surface; and then the arms either hang down, enabling him to assist himself with his knuckles; or, what is more usual, he keeps his arms uplifted in nearly an erect position, with the hands pendent ready to seize a

rope, and climb up on the approach of danger or on the obtrusion of strangers. He walks rather quick in the erect posture, but with a waddling gait, and is soon run down if, whilst pursued, he has no opportunity of escaping by climbing. . . . When he walks in the erect posture he turns the leg and foot outwards, which occasions him to have a waddling gait and to seem bow-legged."

Dr. Burrough states of another Gibbon, the Horlack or Hooluk :

They walk erect; and when placed on the floor, or in an open field, balance themselves very prettily, by raising their hands over their head and slightly bending the arm at the wrist and elbow, and then run tolerably fast, rocking from side to side; and, if urged to greater speed, they let fall their hands to the ground, and assist themselves forward, rather jumping than running, still keeping the body, however, nearly erect.

Somewhat different evidence, however, is given by Dr. Winslow Lewis<sup>3</sup> :—

"Their only manner of walking was on their posterior or inferior extremities, the others being raised upwards to preserve their equilibrium, as rope-dancers are assisted by long poles at fairs. Their progression was not by placing one foot before the other, but by simultaneously using both, as in jumping." Dr. Salomon Müller also states that the Gibbons progress upon the ground by a short series of tottering jumps, effected only by the hind limbs, the body being held altogether upright.

But Mr. Martin (l. c., p. 418), who also speaks from direct observation, says of the Gibbons generally :—

Pre-eminently qualified for arboreal

<sup>1</sup> *Man and Monkeys*, p. 423.

<sup>2</sup> *Wanderings in New South Wales*, vol. ii., chap. viii., 1834.

<sup>3</sup> *Boston Journal of Natural History*, vol. i., 1834.



habits, and displaying among the branches amazing activity, the Gibbons are not so awkward or embarrassed on a level surface as might be imagined. They walk erect, with a waddling or unsteady gait, but at a quick pace; the equilibrium of the body requiring to be kept up, either by touching the ground with the knuckles, first on one side, then on the other, or by uplifting the arms so as to poise it. As with the Chimpanzee, the whole of the narrow, long sole of the foot is placed upon the ground at once and raised at once, without any elasticity of step.

After this mass of concurrent and independent testimony, it cannot reasonably be doubted that the Gibbons commonly and habitually assume the erect attitude.

But level ground is not the place where these animals can display their very remarkable and peculiar locomotive powers, and that prodigious activity which almost tempts one to rank them among flying rather than among ordinary climbing mammals.

Mr. Martin (l. c., p. 430) has given so excellent and graphic an account of the movements of a *Hylobates agilis*, living in the Zoological Gardens, in 1840, that I will quote it in full:—

It is almost impossible to convey in words an idea of the quickness and graceful address of her movements: they may indeed be termed aerial, as she seems merely to touch in her progress the branches among which she exhibits her evolutions. In these feats her hands and arms are the sole organs of locomotion; her body hanging as if suspended by a rope, sustained by one hand (the right, for example), she launches herself, by an energetic movement, to a distant branch, which she catches with the left hand; but her hold is less than momentary: the impulse for the next launch is acquired: the branch then aimed at is attained by the right hand again, and quitted instantaneously, and so on, in alternate succession. In this manner spaces of twelve and eighteen feet are cleared, with the greatest ease and uninterruptedly, for hours together, without the slightest

appearance of fatigue being manifested; and it is evident that, if more space could be allowed, distances very greatly exceeding eighteen feet would be as easily cleared; so that Duvaucel's assertion that he has seen these animals launch themselves from one branch to another, forty feet asunder, startling as it is, may be well credited. Sometimes, on seizing a branch in her progress, she will throw herself, by the power of one arm only, completely round it, making a revolution with such rapidity as almost to deceive the eye, and continue her progress with undiminished velocity. It is singular to observe how suddenly this Gibbon can stop, when the impetus given by the rapidity and distance of her swinging leaps would seem to require a gradual abatement of her movements. In the very midst of her flight a branch is seized, the body raised, and she is seen, as if by magic, quietly seated on it, grasping it with her feet. As suddenly she again throws herself into action.

The following facts will convey some notion of her dexterity and quickness. A live bird was let loose in her apartment; she marked its flight, made a long swing to a distant branch, caught the bird with one hand in her passage, and attained the branch with her other hand; her aim, both at the bird and at the branch, being as successful as if one object only had engaged her attention. It may be added that she instantly bit off the head of the bird, picked its feathers, and then threw it down without attempting to eat it.

On another occasion this animal swung herself from her perch, across a passage at least twelve feet wide, against a window which it was thought would be immediately broken: but not so; to the surprise of all, she caught the narrow framework between the panes with her hand, in an instant attained the proper impetus, and sprang back again to the cage she had left—a feat requiring not only great strength, but the nicest precision.

The Gibbons appear to be naturally very gentle, but there is very good evidence that they will bite severely when irritated—a female *Hylobates agilis* having so severely lacerated one man with her long canines that he died; while she had injured others

so much that, by way of precaution, these formidable teeth had been filed down; but, if threatened, she would still turn on her keeper. The Gibbons eat insects, but appear generally to avoid animal food. A Siamang, however, was seen by Mr. Bennett to seize and devour greedily a live lizard. They commonly drink by dipping their fingers in the liquid and then licking them. It is asserted that they sleep in a sitting posture.

Duvaucel affirms that he has seen the females carry their young to the waterside and there wash their faces, in spite of resistance and cries. They are gentle and affectionate in captivity—full of tricks and pettishness, like spoiled children, and yet not devoid of a certain conscience, as an anecdote, told by Mr. Bennett (l. c., p. 156), will show. It would appear that his Gibbon had a peculiar inclination for disarranging things in the cabin. Among these articles, a piece of soap would especially attract his notice, and for the removal of this he had been once or twice scolded. "One morning," says Mr. Bennett, "I was writing, the ape being present in the cabin, when, casting my eyes towards him, I saw the little fellow taking the soap. I watched him without his perceiving that I did so: and he occasionally would cast a furtive glance towards the place where I sat. I pretended to write; he, seeing me busily occupied, took the soap, and moved away with it in his paw. When he had walked half the length of the cabin, I spoke quietly, without frightening him. The instant he found I saw him he walked back again, and deposited the soap nearly in the same place from whence he had taken it.

There was certainly something more than instinct in that action: he evidently betrayed a consciousness of having done wrong both by his first and last actions—and what is reason if that is not an exercise of it?"

The most elaborate account of the natural history of the ORANG-UTAN extant is that given in the *Verhandelingen over de Natuurlijke Geschiedenis der Nederlandsche overzeesche Bezittingen* (1839-45), by Dr. Salomon Müller and Dr. Schlegel, and I shall base what I have to say upon this subject almost entirely on their statements, adding, here and there, particulars of interest from the writings of Brooke, Wallace, and others.

The Orang-Utan would rarely seem to exceed four feet in height, but the body is very bulky, measuring two-thirds of the height in circumference.<sup>1</sup>

The Orang-Utan is found only in Sumatra and Borneo, and is common in neither of these islands—in both of which it occurs always in low, flat

<sup>1</sup> The largest Orang-Utan, cited by Temminck, measured, when standing upright, 4 ft. ; but he mentions having just received news of the capture of an Orang 5 ft. 3 in. high. Schlegel and Müller say that their largest old male measured, upright, 1.25 Netherlands "el"; and from the crown to the end of the toes, 1.5 el; the circumference of the body being about 1 el. The largest old female was 1.09 el high, when standing. The adult skeleton in the College of Surgeons' Museum, if set upright, would stand 3 ft. 6-8 in. from crown to sole. Dr. Humphry gives 3 ft. 8 in. as the mean height of two Orangs. Of seventeen Orangs examined by Mr. Wallace, the largest was 4 ft. 2 in. high, from the heel to the crown of the head. Mr. Spencer St. John, however, in his *Life in the Forests of the Far East*, tells us of an Orang of "5 ft. 2 in., measuring fairly from the head to the heel," 15 in. across the face, and 12 in. round the wrist. It does not appear, however, that Mr. St. John measured this Orang himself.



plains, never in the mountains. It loves the densest and most sombre of the forests, which extend from the sea-shore inland, and thus is found only in the eastern half of Sumatra,

lating is dense. In favourable places the hunter may, by good fortune, see three or four in a day.

Except in the pairing time, the old males usually live by themselves.



FIG. 9.—An adult male Orang-Utan, after Müller and Schlegel.

where alone such forests occur, though, occasionally, it strays over to the western side.

On the other hand, it is generally distributed through Borneo, except in the mountains, or where the popu-

The old females and the immature males, on the other hand, are often met with in twos and threes; and the former occasionally have young with them, though the pregnant females usually separate themselves, and



sometimes remain apart after they have given birth to their offspring. The young Orangs seem to remain unusually long under their mother's protection, probably in consequence of their slow growth. While climbing, the mother always carries her young against her bosom, the young holding on by his mother's hair.<sup>1</sup> At what time of life the Orang-Utan becomes capable of propagation, and how long the females go with young, is unknown; but it is probable that they are not adult until they arrive at ten or fifteen years of age. A female which lived for five years at Batavia had not attained one-third the height of the wild females. It is probable that, after reaching adult years, they go on growing, though slowly, and that they live to forty or fifty years. The Dyaks tell of old Orangs, which have not only lost all their teeth, but which find it so troublesome to climb that they maintain themselves on windfalls and juicy herbage.

The Orang is sluggish, exhibiting none of that marvellous activity characteristic of the Gibbons. Hunger alone seems to stir him to exertion, and when it is stilled he relapses into repose. When the animal sits, it curves its back and bows its head, so as to look straight down on the ground; sometimes it holds on with its hands by a higher branch, sometimes lets them hang phlegmatically down by his side—and in these posi-

tions the Orang will remain, for hours together, in the same spot, almost without stirring, and only now and then giving utterance to its deep, growling voice. By day he usually climbs from one tree-top to another, and only at night descends to the ground, and, if then threatened with danger, he seeks refuge among the underwood. When not hunted, he remains a long time in the same locality, and sometimes stops for many days on the same tree—a firm place among its branches serving him for a bed. It is rare for the Orang to pass the night in the summit of a large tree, probably because it is too windy and cold there for him; but, as soon as night draws on, he descends from the height and seeks out a fit bed in the lower and darker part, or in the leafy top of a small tree, among which he prefers Nibong Palms, Pandani, or one of those parasitic Orchids which give the primæval forests of Borneo so characteristic and striking an appearance. But whenever he determines to sleep, there he prepares himself a sort of nest: little boughs and leaves are drawn together round the selected spot, and bent crosswise over one another; while, to make the bed soft, great leaves of Ferns, of Orchids, of *Pandanus fascicularis*, *Nipa fruticans*, &c., are laid over them. Those which Müller saw, many of them being very fresh, were situated at a height of ten to twenty-five feet above the ground, and had a circumference, on the average, of two or three feet. Some were packed many inches thick with *Pandanus* leaves; others were remarkable only for the cracked twigs, which, united in a common centre, formed a regular

<sup>1</sup> See Mr. Wallace's account of an infant Orang-Utan, in the *Annals of Natural History* for 1856. Mr. Wallace provided his interesting charge with an artificial mother of buffalo-skin, but the cheat was too successful. The infant's entire experience led it to associate teats with hair, and, feeling the latter, it spent its existence in vain endeavours to discover the former.

platform. "The rude *hut*," says Sir James Brooke, "which they are stated to build in the trees, would be more properly called a seat or nest, for it has no roof or cover of any sort. The facility with which they form this nest is curious, and I had an opportunity of seeing a wounded female weave the branches together and seat herself, within a minute."

According to the Dyaks, the Orang rarely leaves his bed before the sun is well above the horizon and has dissipated the mists. He gets up about nine, and goes to bed again about five; but sometimes not till late in the twilight. He lies sometimes on his back; or, by way of change, turns on one side or the other, drawing his limbs up to his body, and resting his head on his hand. When the night is cold, windy, or rainy, he usually covers his body with a heap of *Pandanus*, *Nipa*, or Fern leaves, like those of which his bed is made, and he is especially careful to wrap up his head in them. It is this habit of covering himself up which has probably led to the fable that the Orang builds huts in the trees.

Although the Orang resides mostly amid the boughs of great trees during the daytime, he is very rarely seen squatting on a thick branch, as other apes, and particularly the Gibbons, do. The Orang, on the contrary, confines himself to the slender leafy branches, so that he is seen right at the top of the trees—a mode of life which is closely related to the constitution of his hinder limbs, and especially to that of his seat. For this is provided with no callosities, such as are possessed by many of the lower apes, and even by the Gibbons; and

those bones of the pelvis which are termed the ischia, and which form the solid framework of the surface on which the body rests in the sitting posture, are not expanded like those of the apes which possess callosities, but are more like those of man.

An Orang climbs so slowly and cautiously<sup>1</sup> as, in this act, to resemble a man more than an ape, taking great care of his feet, so that injury of them seems to affect him far more than it does other apes. Unlike the Gibbons, whose forearms do the greater part of the work as they swing from branch to branch, the Orang never makes even the smallest jump. In climbing, he moves alternately one hand and one foot, or, after having laid fast hold with the hands, he draws up both feet together. In passing from one tree to another he always seeks out a place where the twigs of both come close together, or interlace. Even when closely pursued, his circumspection is amazing; he shakes the branches to see if they will bear him, and then, bending an overhanging bough down by throwing his weight gradually along it, he makes a bridge from the tree he wishes to quit to the next.<sup>2</sup>

On the ground the Orang always goes laboriously and shakily, on all fours. At starting he will run faster than a man, though he may soon be overtaken. The very long arms which, when he runs, are but little

<sup>1</sup> "They are the slowest and least active of all the monkey tribe, and their motions are surprisingly awkward and uncouth."—Sir James Brooke, in the *Proceedings of the Zoological Society*, 1841.

<sup>2</sup> Mr. Wallace's account of the progression of the Orang almost exactly corresponds with this.



bent, raise the body of the Orang remarkably, so that he assumes much the posture of a very old man bent down by age, and making his way along by the help of a stick. In walking the body is usually directed straight forward, unlike the other apes, which run more or less obliquely; except the Gibbons, who in these, as in so many other respects, depart remarkably from their fellows.

The Orang cannot put its feet flat on the ground, but is supported upon their outer edges, the heel resting more on the ground, while the curved toes partly rest upon the ground by the upper side of their first joint, the two outermost toes of each foot completely resting on this surface. The hands are held in the opposite manner, their inner edges serving as the chief support. The fingers are then bent out in such a manner that their foremost joints, especially those of the two innermost fingers, rest upon the ground by their upper sides, while the point of the free and straight thumb serves as an additional fulcrum.

The Orang never stands on its hind legs, and all the pictures representing it as so doing are as false as the assertion that it defends itself with sticks and the like.

The long arms are of especial use, not only in climbing, but in the gathering of food from boughs to which the animal could not trust his weight. Figs, blossoms, and young leaves of various kinds constitute the chief nutriment of the Orang; but strips of bamboo two or three feet long were found in the stomach of a male. They are not known to eat living animals.

Although, when taken young, the

Orang-Utan soon becomes domesticated, and indeed seems to court human society, it is naturally a very wild and shy animal, though apparently sluggish and melancholy. The Dyaks affirm that, when the old males are wounded with arrows only, they will occasionally leave the trees and rush raging upon their enemies, whose sole safety lies in instant flight, as they are sure to be killed if caught.<sup>\*</sup>

But, though possessed of immense

<sup>\*</sup> Sir James Brooke, in a letter to Mr. Waterhouse, published in the *Proceedings of the Zoological Society* for 1841, says:—"On the habits of the Orangs, as far as I have been able to observe them, I may remark that they are as dull and slothful as can well be conceived, and on no occasion, when pursuing them, did they move so fast as to preclude my keeping pace with them easily through a moderately clear forest; and even when obstructions below (such as wading up to the neck) allowed them to get away some distance, they were sure to stop and allow me to come up. I never observed the slightest attempt at defence, and the wood which sometimes rattled about our ears was broken by their weight, and not thrown, as some persons represent. If pushed to extremity, however, the *Pappan* could not be otherwise than formidable, and one unfortunate man, who, with a party, was trying to catch a large one alive, lost two of his fingers, besides being severely bitten on the face, whilst the animal finally beat off his pursuers and escaped." Mr. Wallace, on the other hand, affirms that he has several times observed them throwing down branches when pursued. "It is true he does not throw them *at* a person, but casts them down vertically; for it is evident that a bough cannot be thrown to any distance from the top of a lofty tree. In one case a female *Mias*, on a durian tree kept up for at least ten minutes a continuous shower of branches and of the heavy, spined fruits, as large as 32-pounders, which most effectually kept us clear of the tree she was on. She could be seen breaking them off and throwing them down with every appearance of rage, uttering at intervals a loud pumping grunt, and evidently meaning mischief."—"On the Habits of the Orang-Utan," *Annals of Nat. History*, 1856. This statement, it will be observed, is quite in accordance with that contained in the letter of the Resident Palm quoted above (p. 17).



strength, it is rare for the Orang to attempt to defend itself, especially when attacked with fire-arms. On such occasions he endeavours to hide himself, or to escape along the top-most branches of the trees, breaking off and throwing down the boughs as he goes. When wounded he betakes himself to the highest attainable point of the tree, and emits a singular cry, consisting at first of high notes, which at length deepen into a low roar, not unlike that of a panther. While giving out the high notes the Orang thrusts out his lips into a funnel shape; but in uttering the low notes he holds his mouth wide open, and at the same time the great throat bag, or laryngeal sac, becomes distended.

According to the Dyaks, the only animal the Orang measures his strength with is the crocodile, who occasionally seizes him on his visits to the water side. But they say that the Orang is more than a match for his enemy, and beats him to death, or rips up his throat by pulling the jaws asunder!

Much of what has been here stated was probably derived by Dr. Müller from the reports of his Dyak hunters; but a large male, four feet high, lived in captivity, under his observation, for a month, and receives a very bad character.

"He was a very wild beast," says Müller, "of prodigious strength, and false and wicked to the last degree. If anyone approached, he rose up slowly with a low growl, fixed his eyes in the direction in which he meant to make his attack, slowly passed his hand between the bars of his cage, and then, extending his long arm, gave a sudden grip—usually at

the face." He never tried to bite (though Orangs will bite one another), his great weapons of offence and defence being his hands.

His intelligence was very great; and Müller remarks that, though the faculties of the Orang have been estimated too highly, yet Cuvier, had he seen this specimen, would not have considered its intelligence to be only a little higher than that of the dog.

His hearing was very acute, but the sense of vision seemed to be less perfect. The under lip was the great organ of touch, and played a very important part in drinking, being thrust out like a trough, so as either to catch the falling rain, or to receive the contents of the half cocoa-nut shell full of water with which the Orang was supplied, and which, in drinking, he poured into the trough thus formed.

In Borneo the Orang-Utan of the Malays goes by the name of *Mias* among the Dyaks, who distinguish several kinds as *Mias Pappan*, or *Zimo*, *Mias Kassu*, and *Mias Rambi*. Whether these are distinct species, however, or whether they are mere races, and how far any of them are identical with the Sumatran Orang, as Mr. Wallace thinks the *Mias Pappan* to be, are problems which are at present undecided; and the variability of these great Apes is so extensive that the settlement of the question is a matter of great difficulty. Of the form called "*Mias Pappan*" Mr. Wallace<sup>1</sup> observes:—

It is known by its large size, and by the lateral expansion of the face into fatty protuberances, or ridges, over the tem-

<sup>1</sup> "On the Orang-Utan, or *Mias* of Borneo," *Annals of Natural History*, 1856.

poral muscles, which have been mis-termed *callosities*, as they are perfectly soft, smooth, and flexible. Five of this form, measured by me, varied only from 4 feet 1 inch to 4 feet 2 inches in height, from the heel to the crown of the head, the girth of the body from 3 feet to 3 feet  $7\frac{1}{2}$  inches, and the extent of the outstretched arms from 7 feet 2 inches to 7 feet 6 inches; the width of the face from 10 to  $13\frac{1}{4}$  inches. The colour and length of the hair varied in different individuals, and in different parts of the same individual; some possessed a rudimentary nail on the great toe, others none at all; but they otherwise present no external differences on which to establish even varieties of a species.

Yet, when we examine the crania of these individuals, we find remarkable differences of form, proportion, and dimension, no two being exactly alike. The slope of the profile, and the projection of the muzzle, together with the size of the cranium, offer differences as decided as those existing between the most strongly marked forms of the Caucasian and African crania in the human species. The orbits vary in width and height, the cranial ridge is either single or double, either much or little developed, and the zygomatic aperture varies considerably in size. This variation in the proportions of the crania enables us satisfactorily to explain the marked difference presented by the single-crested and double-crested skulls, which have been thought to prove the existence of two large species of Orang. The external surface of the skull varies considerably in size, as do also the zygomatic aperture and the temporal muscle; but they bear no necessary relation to each other, a small muscle often existing with a large cranial surface, and *vice versa*. Now, those skulls which have the largest and strongest jaws and the widest zygomatic aperture have the muscles so large that they meet on the crown of the skull, and deposit the bony ridge which separates them, and which is the highest in that which has the smallest cranial surface. In those which combine a large surface with comparatively weak jaws, and small zygomatic aperture, the muscles, on each side, do not extend to the crown, a space of from 1 to 2 inches remaining between them, and along their margins small ridges are formed. Intermediate forms are found, in which the ridges meet only in the hinder part of the skull. The

form and size of the ridges are therefore independent of age, being sometimes more strongly developed in the less aged animal. Professor Temminck states that the series of skulls in the Leyden Museum shows the same result.

Mr. Wallace observed two male adult Orangs (Mias Kassu of the Dyaks), however, so very different from any of these that he concludes them to be specifically distinct; they were respectively 3 feet  $8\frac{1}{2}$  inches and 3 feet  $9\frac{1}{2}$  inches high, and possessed no sign of the cheek excrescences, but otherwise resembled the larger kinds. The skull has no crest, but two bony ridges,  $1\frac{3}{4}$  inches to 2 inches apart, as in the *Simia morio* of Professor Owen. The teeth, however, are immense, equalling or surpassing those of the other species. The females of both these kinds, according to Mr. Wallace, are devoid of excrescences, and resemble the smaller males, but are shorter by  $1\frac{1}{2}$  to 3 inches, and their canine teeth are comparatively small, subtruncated and dilated at the base, as in the so-called *Simia morio*, which is, in all probability, the skull of a female of the same species as the smaller males. Both males and females of this smaller species are distinguishable, according to Mr. Wallace, by the comparatively large size of the middle incisors of the upper jaw.

So far as I am aware, no one has attempted to dispute the accuracy of the statements which I have just quoted regarding the habits of the two Asiatic man-like Apes; and, if true, they must be admitted as evidence that such an Ape—

istly, May readily move along the ground in the erect, or semi-erect,



position, and without direct support from its arms.

2ndly, That it may possess an extremely loud voice, so loud as to be readily heard one or two miles.

3rdly, That it may be capable of great viciousness and violence when irritated: and this is especially true of adult males.

4thly, That it may build a nest to sleep in.

Such being well-established facts respecting the Asiatic Anthropoids, analogy alone might justify us in expecting the African species to offer similar peculiarities, separately or combined; or, at any rate, would destroy the force of any attempted *a priori* argument against such direct testimony as might be adduced in favour of their existence. And, if the organisation of any of the African Apes could be demonstrated to fit it better than either of its Asiatic allies for the erect position and for efficient attack, there would be still less reason for doubting its occasional adoption of the upright attitude or of aggressive proceedings.

From the time of Tyson and Tulpius downwards, the habits of the young CHIMPANZEE in a state of captivity have been abundantly reported and commented upon. But trustworthy evidence as to the manners and customs of adult Anthropoids of this species, in their native woods, was almost wanting up to the time of the publication of the paper by Dr. Savage, to which I have already referred; containing notes of the observations which he made, and of the information which he collected from sources which he considered trustworthy, while resident at Cape

Palmas, at the north-western limit of the Bight of Benin.

The adult Chimpanzees measured by Dr. Savage never exceeded, though the males may almost attain, five feet in height.

When at rest, the sitting posture is that generally assumed. They are sometimes seen standing and walking, but when thus detected they immediately take to all fours, and flee from the presence of the observer. Such is their organisation that they cannot stand erect, but lean forward. Hence they are seen, when standing, with the hands clasped over the occiput, or the lumbar region, which would seem necessary to balance or ease of posture.

The toes of the adult are strongly flexed and turned inwards, and cannot be perfectly straightened. In the attempt the skin gathers into thick folds on the back, showing that the full expansion of the foot, as is necessary in walking, is unnatural. The natural position is on all fours, the body anteriorly resting upon the knuckles. These are greatly enlarged, with the skin protuberant and thickened like the sole of the foot.

They are expert climbers, as one would suppose from their organization. In their gambols they swing from limb to limb to a great distance, and leap with astonishing agility. It is not unusual to see the "old folks" (in the language of an observer) sitting under a tree regaling themselves with fruit and friendly chat, while their "children" are leaping around them, and swinging from tree to tree with boisterous merriment.

As seen here, they cannot be called *gregarious*, seldom more than five, or ten at most, being found together. It has been said, on good authority, that they occasionally assemble in large numbers, in gambols. My informant asserts that he saw once not less than fifty so engaged; hooting, screaming, and drumming with sticks upon old logs, which is done in the latter case with equal facility by the four extremities. They do not appear ever to act on the offensive, and seldom, if ever really, on the defensive. When about to be captured, they resist by throwing their arms about their opponent, and attempting to draw him into contact with their teeth. (Savage, l. c., p. 384.)



With respect to this last point Dr. Savage is very explicit in another place :—

*Biting* is their principal art of defence. I have seen one man who had been thus severely wounded in the feet.

The strong development of the canine teeth in the adult would seem to indicate a carnivorous propensity ; but in no state save that of domestication do they manifest it. At first they reject flesh, but easily acquire a fondness for it. The canines are early developed, and evidently designed to act the important part of weapons of defence. When in contact with man, almost the first effort of the animal is—to bite.

They avoid the abodes of men, and build their habitations in trees. Their construction is more that of *nests* than *huts*, as they have been erroneously termed by some naturalists. They generally build not far above the ground. Branches or twigs are bent, or partly broken, and crossed, and the whole supported by the body of a limb or a crotch. Sometimes a nest will be found near the end of a *strong leafy branch* twenty or thirty feet from the ground. One I have lately seen that could not be less than forty feet, and more probably it was fifty. But this is an unusual height.

Their dwelling-place is not permanent, but changed in pursuit of food and solitude, according to the force of circumstances. We more often see them in elevated places ; but this arises from the fact that the low grounds, being more favourable for the natives' rice-farms, are the oftener cleared, and hence are almost always wanting in suitable trees for their nests.....It is seldom that more than one or two nests are seen upon the same tree, or in the same neighbourhood : five have been found, but it was an unusual circumstance.....

They are very filthy in their habits. ....It is a tradition with the natives generally here that they were once members of their own tribe ; that for their depraved habits they were expelled from all human society, and that, through an obstinate indulgence of their vile propensities, they have degenerated into their present state and organisation. They are, however, eaten by them, and when cooked with the oil and pulp of the palm-nut considered a highly palatable morsel.

They exhibit a remarkable degree of intelligence in their habits, and, on the part of the mother, much affection for their young. The second female described was upon a tree when first discovered, with her mate and two young ones (a male and a female). Her first impulse was to descend with great rapidity, and make off into the thicket, with her mate and female offspring. The young male remaining behind, she soon returned to the rescue. She ascended and took him in her arms, at which moment she was shot, the ball passing through the forearm of the young one, on its way to the heart of the mother.....

In a recent case the mother, when discovered, remained upon the tree with her offspring, watching intently the movements of the hunter. As he took aim, she motioned with her hand, precisely in the manner of a human being, to have him desist and go away. When the wound has not proved instantly fatal, they have been known to stop the flow of blood by pressing with the hand upon the part, and, when this did not succeed, to apply leaves and grass.....When shot, they give a sudden screech, not unlike that of a human being in sudden and acute distress.

The ordinary voice of the Chimpanzee, however, is affirmed to be hoarse, guttural, and not very loud, somewhat like "whoo-whoo" (l. c., p. 365).

The analogy of the Chimpanzee to the Orang, in its nest-building habit and in the mode of forming its nest, is exceedingly interesting ; while, on the other hand, the activity of this ape, and its tendency to bite, are particulars in which it rather resembles the Gibbons. In extent of geographical range, again, the Chimpanzees—which are found from Sierra Leone to Congo—remind one of the Gibbons, rather than of either of the other man-like apes ; and it seems not unlikely that, as is the case with the Gibbons, there may be several species spread over the geographical area of the genus

The same excellent observer from whom I have borrowed the preceding account of the habits of the adult Chimpanzee, published, fifteen years ago,<sup>1</sup> an account of the GORILLA,

has really been added that in justice to Dr. Savage I give it almost in full:—

It should be borne in mind that my account is based upon the statements of



FIG. 10.—The Gorilla (after Wolff).

which has, in its most essential points, been confirmed by subsequent observers, and to which so very little

<sup>1</sup> "Notice of the external character and habits of *Troglodytes Gorilla*." *Boston Journal of Natural History*, 1847.

the aborigines of that region (the Gaboon). In this connection, it may also be proper for me to remark, that having been a missionary resident for several years, studying, from habitual intercourse, the African mind and character, I felt myself prepared to discriminate and decide upon the probability of their



statements. Besides, being familiar with the history and habits of its interesting congener (*Trog. niger*, Geoff.), I was able to separate their accounts of the two animals, which, having the same locality and a similarity of habit, are confounded in the minds of the mass, especially as but few—such as traders to the interior and huntsmen—have ever seen the animal in question.

The tribe from which our knowledge of the animal is derived, and whose territory forms its habitat, is the *Mpongwe*, occupying both banks of the River Gaboon, from its mouth to some fifty or sixty miles upward.....

If the word "Pongo" be of African origin, it is probably a corruption of the word *Mpongwe*, the name of the tribe on the banks of the Gaboon, and hence applied to the region they inhabit. Their local name for the Chimpanzee is *Enché-*



FIG. 11.—Gorilla walking (after Wolff).

*eko*, as near as it can be Anglicised, from which the common term 'Jocko' probably comes. The *Mpongwe* appellation for its new congener is *Engé-ena*, prolonging the sound of the first vowel, and slightly sounding the second.

The habitat of the *Engé-ena* is the interior of Lower Guinea, whilst that of the *Enché-eko* is nearer the sea-board.

Its height is about five feet; it is disproportionately broad across the shoulders, thickly covered with coarse black hair, which is said to be similar in its arrangement to that of the *Enché-eko*; with age it becomes grey, which fact has given rise to the report that both animals are seen of different colours.

*Head.*—The prominent features of the head are, the great width and elongation of the face, the depth of the molar region, the branches of the lower jaw being very

deep and extending far backward, and the comparative smallness of the cranial portion; the eyes are very large, and said to be like those of the *Enché-eko*, a bright hazel; nose broad and flat, slightly elevated towards the root; the muzzle broad, and prominent lips and chin, with scattered grey hairs; the under lip highly mobile, and capable of great elongation when the animal is enraged, then hanging over the chin; skin of the face and ears naked, and of a dark brown, approaching to black.

The most remarkable feature of the head is a high ridge, or crest of hair, in the course of the sagittal suture, which meets posteriorly with a transverse ridge of the same, but less prominent, running round from the back of one ear to the other. The animal has the power of moving the scalp freely forward and back, and when enraged is said to contract it strongly over the brow, thus bringing down the hairy ridge and pointing the hair forward, so as to present an indescribably ferocious aspect.

Neck short, thick, and hairy; chest and shoulders very broad, said to be fully double the size of the *Enché-ekos*; arms very long, reaching some way below the knee—the fore-arm much the shortest; hands very large, the thumbs much larger than the fingers.....

The gait is shuffling; the motion of the body, which is never upright as in man, but bent forward, is somewhat rolling, or from side to side. The arms being longer than the Chimpanzee, it does not stoop as much in walking; like that animal, it makes progression by thrusting its arms forward, resting the hands on the ground, and then giving the body a half jumping, half swinging motion between them. In this act it is said not to flex the fingers, as does the Chimpanzee, resting on its knuckles, but to extend them, making a fulcrum of the hand. When it assumes the walking posture, to which it is said to be much inclined, it balances its huge body by flexing its arms upward.

They live in bands, but are not so numerous as the Chimpanzees: the females generally exceed the other sex in number. My informants all agree in the assertion that but one adult male is seen in a band; that when the young males grow up a contest takes place for mastery, and the strongest, by killing and driving out the others, establishes himself as the head of the community.



Dr. Savage repudiates the stories about the Gorillas carrying off women and vanquishing elephants, and then adds :—

Their dwellings, if they may be so called, are similar to those of the Chimpanzee, consisting simply of a few sticks and leafy branches, supported by the crotches and limbs of trees ; they afford no shelter, and are occupied only at night.

They are exceedingly ferocious, and always offensive in their habits, never running from man, as does the Chimpanzee. They are objects of terror to the natives, and are never encountered by them except on the defensive. The few that have been captured were killed by elephant-hunters and native traders, as they came suddenly upon them while passing through the forests.

It is said that when the male is first seen he gives a terrific yell that resounds far and wide through the forest, something like kh—ah ! kh—ah ! prolonged and shrill. His enormous jaws are widely opened at each expiration, his under lip hangs over the chin, and the hairy ridge and scalp are contracted upon the brow, presenting an aspect of indescribable ferocity.

The females and young, at the first cry, quickly disappear. He then approaches the enemy in great fury, pouring out his horrid cries in quick succession. The hunter awaits his approach with his gun extended : if his aim is not sure, he permits the animal to grasp the barrel, and as he carries it to his mouth (which is his habit) he fires. Should the gun fail to go off, the barrel (that of the ordinary musket, which is thin) is crushed between his teeth, and the encounter soon proves fatal to the hunter.

In the wild state their habits are in general like those of the *Troglodytes niger*, building their nests loosely in trees, living on similar fruits, and changing their place of resort from force of circumstances.

Dr. Savage's observations were confirmed and supplemented by those of Mr. Ford, who communicated an interesting paper on the Gorilla to the Philadelphian Academy of Sciences in 1852. With respect to the geo-

graphical distribution of this greatest of all the man-like Apes, Mr. Ford remarks :—

This animal inhabits the range of mountains that traverse the interior of Guineà, from the Cameroon in the north to Angola in the south, and about 100 miles inland, and called by the geographers Crystal Mountains. The limit to which this animal extends, either north or south, I am unable to define. But that limit is doubtless some distance north of this river [Gaboön]. I was able to certify myself of this fact in a late excursion to the headwaters of the Mooney (Danger) River, which comes into the sea some sixty miles from this place. I was informed (credibly, I think) that they were numerous among the mountains in which that river rises, and far north of that.

In the south this species extends to the Congo River, as I am told by native traders who have visited the coast between the Gaboon and that river. Beyond that I am not informed. This animal is only found at a distance from the coast in most cases, and, according to my best information, approaches it nowhere so nearly as on the south side of this river, where they have been found within ten miles of the sea. This, however, is only of late occurrence. I am informed by some of the oldest Mpongwe men that formerly he was only found on the sources of the river, but that at present he may be found within half-a-day's walk of its mouth. Formerly he inhabited the mountainous ridge where Bushmen alone inhabited, but now he boldly approaches the Mpongwe plantations. This is doubtless the reason of the scarcity of information in years past, as the opportunities for receiving a knowledge of the animal have not been wanting ; traders having for one hundred years frequented this river, and specimens, such as have been brought here within a year, could not have been exhibited without having attracted the attention of the most stupid.

One specimen Mr. Ford examined weighed 170 lbs., without the thoracic or pelvic viscera, and measured four feet four inches round the chest. This writer describes so minutely and graphically the onslaught of the Gorilla—though he does not for a

moment pretend to have witnessed the scene—that I am tempted to give this part of his paper in full for comparison with other narratives :—

He always rises to his feet, when making an attack, though he approaches his antagonist in a stooping posture.

Though he never lies in wait, yet when he hears, sees, or scents a man he immediately utters his characteristic cry, prepares for an attack, and always acts on the offensive. The cry he utters resembles a grunt more than a growl, and is similar to the cry of the Chimpanzee when irritated, but vastly louder. It is said to be audible at a great distance. His preparation consists in attending the females and young ones, by whom he is usually accompanied, to a little distance. He, however, soon returns, with his crest erect and projecting forward, his nostrils dilated, and his under-lip thrown down, at the same time uttering his characteristic yell, designed, it would seem, to terrify his antagonist. Instantly, unless he is disabled by a well-directed shot, he makes an onset, and striking his antagonist with the palm of his hands, or seizing him with a grasp from which there is no escape, he dashes him upon the ground, and lacerates him with his tusks.

He is said to seize a musket, and instantly crush the barrel between his teeth.....This animal's savage nature is very well shown by the implacable desperation of a young one that was brought here. It was taken very young and kept four months, and many means were used to tame it; but it was incorrigible, so that it bit me an hour before it died.

Mr. Ford discredits the house-building and elephant-driving stories, and says that no well-informed natives believe them. They are tales told to children.

I might quote other testimony to a similar effect, but, as it appears to me, less carefully weighed and sifted, from the letters of MM. Franquet and Gautier Laboullay, appended to the memoir of M. I. G. St. Hilaire, which I have already cited.

Bearing in mind what is known regarding the Orang and the Gibbon, the statements of Dr. Savage and Mr. Ford do not appear to me to be justly open to criticism on *a priori* grounds. The Gibbons, as we have seen, readily assume the erect posture, but the Gorilla is far better fitted by its organisation for that attitude than are the Gibbons: if the laryngeal pouches of the Gibbons; as is very likely, are important in giving volume to a voice which can be heard for half a league, the Gorilla, which has similar sacs, more largely developed, and whose bulk is five-fold that of a Gibbon, may well be audible for twice that distance. If the Orang fights with his hands, the Gibbons and Chimpanzees with their teeth, the Gorilla may, probably enough, do either or both; nor is there anything to be said against either Chimpanzee or Gorilla building a nest, when it is proved that the Orang-Utan habitually performs that feat.

With all this evidence, now ten to fifteen years old, before the world, it is not a little surprising that the assertions of a recent traveller, who, so far as the Gorilla is concerned, really does very little more than repeat, on his own authority, the statements of Savage and of Ford, should have met with so much and such bitter opposition. If subtraction be made of what was known before, the sum and substance of what M. Du Chaillu has affirmed as a matter of his own observation respecting the Gorilla is that, in advancing to the attack, the great brute beats his chest with his fists. I confess I see nothing very improbable, or very much worth disputing about, in this statement.



With respect to the other man-like Apes of Africa, M. Du Chaillu tells us absolutely nothing, of his own knowledge, regarding the common Chimpanzee; but he informs us of a bald-headed species or variety, the *nschiëgo mbouve*, which builds itself a shelter, and of another rare kind with a comparatively small face, large facial angle, and peculiar note, resembling "Kooloo."

As the Orang shelters itself with a rough coverlet of leaves, and the common Chimpanzee, according to that eminently trustworthy observer Dr. Savage, makes a sound like "Whoo-whoo," the grounds of the

summary repudiation with which M. Du Chaillu's statements on these matters have been met are not obvious.

If I have abstained from quoting M. Du Chaillu's work, then, it is not because I discern any inherent improbability in his assertions respecting the man-like Apes; nor from any wish to throw suspicion on his veracity; but because, in my opinion, so long as his narrative remains in its present state of unexplained and apparently inexplicable confusion it has no claim to original authority respecting any subject whatsoever.

It may be truth, but it is not evidence.

## II.

### ON THE RELATIONS OF MAN TO THE LOWER ANIMALS

Multis videri poterit, majorem esse differentiam Simiæ et Hominis, quam diei et noctis; verum tamen hi, comparatione instituta inter summos Europæ Heroës et Hottentottos ad Caput Bonæ Spei degentes, difficillime sibi persuadebunt, has eosdem habere natales; vel si virginem nobilem aulicam, maxime contam et humanissimam, conferre vellent cum homine sylvestri et sibi relicto, vix augurari possent, hunc et illam ejusdem esse speciei.—*Linnæi Amœnitates Acad.*, "*Anthropomorpha*."

THE question of questions for mankind—the problem which underlies all others, and is more deeply interesting than any other—is the ascertainment of the place which Man occupies in nature, and of his relations to the

universe of things. Whence our race has come; what are the limits of our power over nature, and of nature's power over us; to what goal we are tending; are the problems which present themselves anew and with undiminished interest to every man born into the world. Most of us, shrinking from the difficulties and dangers which beset the seeker after original answers to these riddles, are contented to ignore them altogether, or to smother the investigating spirit under the feather-bed of respected and respectable tradition. But, in every age, one or two restless spirits, blessed with that constructive genius which can only build on a secure



foundation, or cursed with the mere spirit of scepticism, are unable to follow in the well-worn and comfortable track of their forefathers and contemporaries, and, unmindful of thorns and stumbling-blocks, strike out into paths of their own. The sceptics end in the infidelity which asserts the problem to be insoluble, or in the atheism which denies the existence of any orderly progress and governance of things; the men of genius propound solutions which grow into systems of Theology or of Philosophy, or, veiled in musical language which suggests more than it asserts, take the shape of the Poetry of an epoch.

Each such answer to the great question, invariably asserted by the followers of its propounder, if not by himself, to be complete and final, remains in high authority and esteem, it may be for one century, or it may be for twenty; but, as invariably, time proves each reply to have been a mere approximation to the truth—tolerable chiefly on account of the ignorance of those by whom it was accepted, and wholly intolerable when tested by the larger knowledge of their successors.

In a well-worn metaphor a parallel is drawn between the life of man and the metamorphosis of the caterpillar into the butterfly; but the comparison may be more just as well as more novel if for its former term we take the mental progress of the race. History shows that the human mind, fed by constant accessions of knowledge, periodically grows too large for its theoretical coverings, and bursts them asunder to appear in new habiliments, as the feeding and grow-

ing grub at intervals casts its too narrow skin and assumes another, itself but temporary. Truly the imago state of Man seems to be terribly distant, but every moult is a step gained, and of such there have been many.

Since the revival of learning, whereby the Western races of Europe were enabled to enter upon that progress towards true knowledge which was commenced by the philosophers of Greece, but was almost arrested in subsequent long ages of intellectual stagnation, or, at most, gyration, the human larva has been feeding vigorously, and moulting in proportion. A skin of some dimension was cast in the sixteenth century, and another towards the end of the eighteenth, while, within the last fifty years, the extraordinary growth of every department of physical science has spread among us mental food of so nutritious and stimulating a character that a new ecdysis seems imminent. But this is a process not unusually accompanied by many throes and some sickness and debility, or, it may be, by graver disturbances; so that every good citizen must feel bound to facilitate the process, and, even if he have nothing but a scalpel to work withal, to ease the cracking integument to the best of his ability.

In this duty lies my excuse for the publication of these essays. For it will be admitted that some knowledge of man's position in the animate world is an indispensable preliminary to the proper understanding of his relations to the universe; and this again resolves itself, in the long run, into an inquiry into the nature and the closeness of the ties which connect him

with those singular creatures whose history<sup>1</sup> has been sketched in the preceding pages.

The importance of such an inquiry is indeed intuitively manifest. Brought face to face with these blurred copies of himself, the least thoughtful of men is conscious of a certain shock, due perhaps not so much to disgust at the aspect of what looks like an insulting caricature, as to the awakening of a sudden and profound mistrust of time-honoured theories and strongly-rooted prejudices regarding his own position in nature, and his relations to the under-world of life; while that which remains a dim suspicion for the unthinking becomes a vast argument, fraught with the deepest consequences, for all who are acquainted with the recent progress of the anatomical and physiological sciences.

I now propose briefly to unfold that argument, and to set forth, in a form intelligible to those who possess no special acquaintance with anatomical science, the chief facts upon which all conclusions respecting the nature and the extent of the bonds which connect man with the brute world must be based; I shall then indicate the one immediate conclusion which, in my judgment, is justified by those facts, and I shall finally discuss the bearing of that conclusion upon the hypotheses which have been entertained respecting the Origin of Man.

The facts to which I would first direct the reader's attention, though

<sup>1</sup> It will be understood that in the preceding Essay I have selected for notice, from the vast mass of papers which have been written upon the man-like Apes, only those which seem to me to be of special moment.

ignored by many of the professed instructors of the public mind, are easy of demonstration, and are universally agreed to by men of science; while their significance is so great that whoso has duly pondered over them will, I think, find little to startle him in the other revelations of Biology. I refer to those facts which have been made known by the study of Development.

It is a truth of very wide, if not of universal, application, that every living creature commences its existence under a form different from, and simpler than, that which it eventually attains.

The oak is a more complex thing than the little rudimentary plant contained in the acorn; the caterpillar is more complex than the egg; the butterfly than the caterpillar; and each of these beings, in passing from its rudimentary to its perfect condition, runs through a series of changes, the sum of which is called its Development. In the higher animals these changes are extremely complicated; but within the last half-century the labours of such men as Von Baer, Rathke, Reichert, Bischoff, and Remak have almost completely unravelled them, so that the successive stages of development which are exhibited by a Dog, for example, are now as well known to the embryologist as are the steps of the metamorphosis of the silkworm moth to the school-boy. It will be useful to consider with attention the nature and the order of the stages of canine development, as an example of the process in the higher animals generally.

The Dog, like all animals, save the



very lowest (and further inquiries may not improbably remove the apparent exception), commences its existence as an egg : as a body which is, in every sense, as much an egg as that of a hen, but is devoid of that accumulation of nutritive matter which confers upon the bird's egg its exceptional size and domestic utility ; and wants the shell, which would not only be useless to an animal incubated within the body of its parent, but would cut it off from access to the source of that nutriment which the young creature

more solid rounded body, termed the *germinal spot* (*b*).

The egg, or "Ovum," is originally formed within a gland, from which, in due season, it becomes detached, and passes into the living chamber fitted for its protection and maintenance during the protracted process of gestation. Here, when subjected to the required conditions, this minute and apparently insignificant particle of living matter becomes animated by a new and mysterious activity. The germinal vesicle and spot cease to be




FIG. 12.—A. Egg of the Dog, with the vitelline membrane burst, so as to give exit to the yolk, the germinal vesicle (*a*), and its included spot (*b*). B. C. D. E. F. Successive changes of the yolk indicated in the text. (After Bischoff.)

requires, but which the minute egg of the mammal does not contain within itself.

The Dog's egg is, in fact, a little spheroidal bag (Fig. 12), formed of a delicate transparent membrane called the *vitelline membrane*, and about  $\frac{1}{160}$  to  $\frac{1}{120}$ th of an inch in diameter. It contains a mass of viscid nutritive matter—the *yolk*—within which is enclosed a second much more delicate spheroidal bag, called the *germinal vesicle* (*a*). In this, lastly, lies a

discernible (their precise fate being one of the yet unsolved problems of embryology), but the yolk becomes circumferentially indented, as if an invisible knife had been drawn round it, and thus appears divided into two hemispheres (Fig. 12, C).

By the repetition of this process in various planes, these hemispheres become subdivided, so that four segments are produced (D) ; and these, in like manner, divide and subdivide again, until the whole yolk is



converted into a mass of granules, each of which consists of a minute spheroid of yolk-substance, enclosing a central particle, the so-called *nucleus* (F). Nature, by this process, has attained much the same result as that at which a human artificer arrives by his operations in a brickfield. She takes the rough plastic material of the yolk and breaks it up into well-shaped, tolerably even-sized masses, handy for building up into any part of the living edifice.

Next, the mass of organic bricks, or

the groove on each side next rises up into a fold, the rudiment of the side wall of that long cavity which will eventually lodge the spinal marrow and the brain; and in the floor of this chamber appears a solid cellular cord, the so-called *notochord*. One end of the enclosed cavity dilates to form the head (Fig. 13, B), the other remains narrow, and eventually becomes the tail; the side walls of the body are fashioned out of the downward continuation of the walls of the groove; and from them, by and by,




FIG. 13.—A. Earliest rudiment of the Dog. B. Rudiment further advanced, showing the foundations of the head, tail, and vertebral column. C. The very young puppy, with attached ends of the yolk-sac and allantois, and invested in the amnion.

*cells* as they are technically called, thus formed, acquires an orderly arrangement, becoming converted into a hollow spheroid with double walls. Then, upon one side of this spheroid, appears a thickening, and, by and by, in the centre of the area of thickening, a straight shallow groove (Fig. 13, A) marks the central line of the edifice which is to be raised, or, in other words, indicates the position of the middle line of the body of the future dog. The substance bounding

grow out little buds which, by degrees, assume the shape of limbs. Watching the fashioning process stage by stage, one is forcibly reminded of the modeller in clay. Every part, every organ, is at first, as it were, pinched up rudely, and sketched out in the rough; then shaped more accurately; and only at last receives the touches which stamp its final character.

Thus, at length, the young puppy assumes such a form as is shown in Fig. 13, C. In this condition it has

a disproportionately large head, as dissimilar to that of a dog as the bud-like limbs are unlike his legs.

The remains of the yelk, which have not yet been applied to the nutrition and growth of the young animal, are contained in a sac attached to the rudimentary intestines, and termed the yelk-sac, or *umbilical vesicle*. Two membranous bags, intended to subserve respectively the protection and nutrition of the young creature, have been developed from the skin and from the under and hinder surface of the body; the former, the so-called *amnion*, is a sac filled with fluid, which invests the whole body of the embryo, and plays the part of a sort of water-bed for it; the other, termed the *allantois*, grows out, loaded with blood-vessels, from the ventral region, and eventually applying itself to the walls of the cavity, in which the developing organism is contained, enables these vessels to become the channel by which the stream of nutriment required to supply the wants of the offspring is furnished to it by the parent.

The structure which is developed by the interlacement of the vessels of the offspring with those of the parent, and by means of which the former is enabled to receive nourishment, and to get rid of effete matters, is termed the *placenta*.

It would be tedious, and it is unnecessary for my present purpose, to trace the process of development further; suffice it to say that, by a long and gradual series of changes, the rudiment here depicted and described becomes a puppy, is born, and then, by still slower and less perceptible steps, passes into the adult Dog.

There is not much apparent resemblance between a barndoor Fowl and the Dog who protects the farm-yard. Nevertheless, the student of development finds, not only that the chick commences its existence as an egg, primarily identical, in all essential respects, with that of the Dog, but that the yelk of this egg undergoes division, that the primitive groove arises, and that the contiguous parts of the germ are fashioned, by precisely similar methods, into a young chick, which, at one stage of its existence, is so like the nascent Dog that ordinary inspection would hardly distinguish the two.

The history of the development of any other vertebrate animal, Lizard, Snake, Frog, or Fish, tells the same story. There is always, to begin with, an egg having the same essential structure as that of the Dog: the yelk of that egg always undergoes division, or *segmentation*, as it is often called: the ultimate products of that segmentation constitute the building materials for the body of the young animal; and this is built up round a primitive groove, in the floor of which a notochord is developed. Furthermore, there is a period in which the young of all these animals resemble one another, not merely in outward form, but in all essentials of structure, so closely that the differences between them are inconsiderable, while, in their subsequent course, they diverge more and more widely from one another. And it is a general law that, the more closely any animals resemble one another in adult structure, the longer and the more intimately do their embryos



resemble one another; so that, for example, the embryos of a Snake and of a Lizard remain like one another longer than do those of a Snake and of a Bird; and the embryo of a Dog and of a Cat remain like one another for a far longer period than do those of a Dog and a Bird; or of a Dog and an Opossum; or even than those of a Dog and a Monkey.

Thus the study of development affords a clear test of closeness of structural affinity, and one turns with impatience to inquire what results are yielded by the study of the develop-

ment of Man. It has been doubtful any time these thirty years. Without question, the mode of origin and the early stages of the development of Man are identical with those of the animals immediately below him in the scale:—without a doubt, in these respects he is far nearer the Apes than the Apes are to the Dog.

The human ovum is about  $\frac{1}{128}$  of an inch in diameter, and might be described in the same terms as that of the Dog, so that I need only refer to the figure illustrative (14 A) of its structure. It leaves the organ in




FIG. 14.—A. Human ovum (after Kölliker). *a*. germinal vesicle. *b*. germinal spot. B. A very early condition of Man, with yolk-sac, allantois and amnion (original). C. A more advanced stage (after Kölliker). (Compare Fig. 13, C.)

ment of Man. Is he something apart? Does he originate in a totally different way from Dog, Bird, Frog, and Fish, thus justifying those who assert him to have no place in nature and no real affinity with the lower world of animal life? Or does he originate in a single germ, pass through the same slow and gradually progressive modifications—depend on the same contrivances for protection and nutrition, and finally enter the world by the help of the same mechanism? The reply is not doubtful for a moment, and has not

which it is formed in a similar fashion and enters the organic chamber prepared for its reception in the same way, the conditions of its development being in all respects the same. It has not yet been possible (and only by some rare chance can it ever be possible) to study the human ovum in so early a developmental stage as that of yolk division; but there is every reason to conclude that the changes it undergoes are identical with those exhibited by the ova of other vertebrated animals; for the formative



materials of which the rudimentary human body is composed, in the earliest conditions in which it has been observed, are the same as those of other animals. Some of these earliest stages are figured on p. 45, and, as will be seen, they are strictly comparable to the very early states of the Dog; the marvellous correspondence between the two which is kept up, even for some time, as development advances, becoming apparent by the simple comparison of the figures with those on page 43.

Indeed, it is very long before the body of the young human being can be readily discriminated from that of the young puppy; but, at a tolerably early period, the two become distinguishable by the different form of their adjuncts, the yelk-sac and the allantois. The former, in the Dog, becomes long and spindle-shaped, while in Man it remains spherical; the latter, in the Dog, attains an extremely large size, and the vascular processes which are developed from it, and eventually give rise to the formation of the placenta (taking root, as it were, in the parental organism, so as to draw nourishment therefrom, as the root of a tree extracts it from the soil), are arranged in an encircling zone, while in Man the allantois remains comparatively small, and its vascular rootlets are eventually restricted to one disk-like spot. Hence, while the placenta of the Dog is like a girdle, that of Man has the cake-like form indicated by the name of the organ.

But, exactly in those respects in which the developing Man differs from the Dog, he resembles the Ape, which, like man, has a spheroidal

yelk-sac and a discoidal—sometimes partially lobed—placenta.

So that it is only quite in the later stages of development that the young human being presents marked differences from the young ape, while the latter departs as much from the dog in its development as the man does.

Startling as the last assertion may appear to be, it is demonstrably true, and it alone appears to me sufficient to place beyond all doubt the structural unity of man with the rest of the animal world, and more particularly and closely with the apes.

Thus identical in the physical processes by which he originates—identical in the early stages of his formation—identical in the mode of his nutrition before and after birth, with the animals which lie immediately below him in the scale—Man, if his adult and perfect structure be compared with theirs, exhibits, as might be expected, a marvellous likeness of organisation. He resembles them as they resemble one another—he differs from them as they differ from one another. And, though these differences and resemblances cannot be weighed and measured, their value may be readily estimated; the scale or standard of judgment touching that value being afforded and expressed by the system of classification of animals now current among zoologists.

A careful study of the resemblances and differences presented by animals has, in fact, led naturalists to arrange them into groups, or assemblages, all the members of each group presenting a certain amount of definable resemblance, and the number of points of similarity being smaller as the group

is larger and *vice versâ*. Thus, all creatures which agree only in presenting the few distinctive marks of animality form the "Kingdom" ANIMALIA. The numerous animals which agree only in possessing the special characters of Vertebrates form one "Sub-kingdom" of this Kingdom. Then the Sub-kingdom VERTEBRATA is subdivided into the five "Classes"—Fishes, Amphibians, Reptiles, Birds, and Mammals, and these into smaller groups called "Orders"; these into "Families" and "Genera"; while the last are finally broken up into the smallest assemblages, which are distinguished by the possession of constant, not-sexual, characters. These ultimate groups are Species.

Every year tends to bring about a greater uniformity of opinion throughout the zoological world as to the limits and characters of these groups, great and small. At present, for example, no one has the least doubt regarding the characters of the classes Mammalia, Aves, or Reptilia; nor does the question arise whether any thoroughly well-known animal should be placed in one class or the other. Again, there is a very general agreement respecting the characters and limits of the orders of Mammals, and as to the animals which are structurally necessitated to take a place in one or another order.

No one doubts, for example, that the Sloth and the Ant-eater, the Kangaroo and the Opossum, the Tiger and the Badger, the Tapir and the Rhinoceros, are respectively members of the same orders. These successive pairs of animals may, and some do, differ from one another immensely, in such matters as the

proportions and structure of their limbs; the number of their dorsal and lumbar vertebræ; the adaptation of their frames to climbing, leaping, or running; the number and form of their teeth; and the characters of their skulls and of the contained brain. But, with all these differences, they are so closely connected in all the more important and fundamental characters of their organisation, and so distinctly separated by these same characters from other animals, that zoologists find it necessary to group them together as members of one order. And if any new animal were discovered, and were found to present no greater difference from the Kangaroo and the Opossum, for example, than these animals do from one another, the zoologist would not only be logically compelled to rank it in the same order with these, but he would not think of doing otherwise.

Bearing this obvious course of zoological reasoning in mind, let us endeavour for a moment to disconnect our thinking selves from the mask of humanity; let us imagine ourselves scientific Saturnians, if you will, fairly acquainted with such animals as now inhabit the Earth, and employed in discussing the relations they bear to a new and singular "erect and featherless biped," which some enterprising traveller, overcoming the difficulties of space and gravitation, has brought from that distant planet for our inspection, well preserved, maybe, in a cask of rum. We should all at once agree upon placing him among the mammalian vertebrates; and his lower jaw, his molars, and his brain would leave no room for doubting the systematic



position of the new genus among those mammals, whose young are nourished during gestation by means of a placenta, or what are called the "placental mammals."

Further, the most superficial study would at once convince us that, among the orders of placental mammals, neither the Whales nor the hoofed creatures, nor the Sloths and Ant-eaters, nor the carnivorous Cats, Dogs, and Bears, still less the Rodent Rats and Rabbits, or the Insectivorous Moles and Hedgehogs, or the Bats, could claim our *Homo* as one of themselves.

There would remain then but one order for comparison—that of the Apes (using that word in its broadest sense), and the question for discussion would narrow itself to this: Is Man so different from any of these Apes that he must form an order by himself? Or does he differ less from them than they differ from one another, and hence must take his place in the same order with them?

Being happily free from all real or imaginary personal interest in the results of the inquiry thus set afoot, we should proceed to weigh the arguments on one side and on the other with as much judicial calmness as if the question related to a new Opossum. We should endeavour to ascertain, without seeking either to magnify or diminish them, all the characters by which our new Mammal differed from the Apes; and if we found that these were of less structural value than those which distinguish certain members of the Ape order from others universally admitted to be of the same order, we should undoubtedly place

the newly-discovered tellurian genus with them.

I now proceed to detail the facts which seem to me to leave us no choice but to adopt the last-mentioned course.

It is quite certain that the Ape which most nearly approaches man in the totality of its organisation is either the Chimpanzee or the Gorilla; and as it makes no practical difference, for the purposes of my present argument, which is selected for comparison, on the one hand with Man, and on the other hand with the rest of the Primates,\* I shall select the latter (so far as its organisation is known), as a brute now so celebrated in prose and verse that all must have heard of him, and have formed some conception of his appearance. I shall take up as many of the most important points of difference between man and this remarkable creature as the space at my disposal will allow me to discuss, and the necessities of the argument demand; and I shall inquire into the value and magnitude of these differences when placed side by side with those which separate the Gorilla from other animals of the same order.

In the general proportions of the body and limbs there is a remarkable difference between the Gorilla and Man which at once strikes the eye. The Gorilla's brain-case is smaller, its trunk larger, its lower limbs shorter, its upper limbs longer in proportion than those of Man.

I find that the vertebral column of

\* We are not at present thoroughly acquainted with the brain of the Gorilla, and, therefore, in discussing cerebral characters, I shall take that of the Chimpanzee as my highest term among the Apes.



a full-grown Gorilla, in the Museum of the Royal College of Surgeons, measures 27 inches along its anterior curvature, from the upper edge of the atlas, or first vertebra of the neck, to the lower extremity of the sacrum; that the arm, without the hand, is  $31\frac{1}{2}$  inches long; that the leg, without the foot, is  $26\frac{1}{2}$  inches long; that the hand is  $9\frac{3}{4}$  inches long; the foot  $11\frac{1}{4}$  inches long.

In other words, taking the length of the spinal column as 100, the arm equals 115, the leg 96, the hand 36, and the foot 41.

In the skeleton of a male Bosjesman, in the same collection, the proportions, by the same measurement, to the spinal column, taken as 100, are—the arm 78, the leg 110, the hand 26, and the foot 32. In a woman of the same race the arm is 83, and the leg 120, the hand and foot remaining the same. In a European skeleton I find the arm to be 80, the leg 117, the hand 26, the foot 35.

Thus the leg is not so different as it looks at first sight in its proportions to the spine in the Gorilla and in the Man, being very slightly shorter than the spine in the former, and between  $\frac{1}{16}$  and  $\frac{1}{8}$  longer than the spine in the latter. The foot is longer and the hand much longer in the Gorilla; but the great difference is caused by the arms, which are very much longer than the spine in the Gorilla, very much shorter than the spine in the Man.

The question now arises, how are the other Apes related to the Gorilla in these respects—taking the length of the spine, measured in the same way, at 100? In an adult Chimpanzee the arm is only 96, the leg 90, the

hand 43, the foot 39; so that the hand and the leg depart more from the human proportion and the arm less, while the foot is about the same as in the Gorilla.

In the Orang the arms are very much longer than in the Gorilla (122), while the legs are shorter (88); the foot is longer than the hand (52 and 48), and both are much longer in proportion to the spine.


In the other man-like Apes, again, the Gibbons, these proportions are still further altered; the length of the arms being to that of the spinal column as 19 to 11; while the legs are also a third longer than the spinal column, so as to be longer than in Man, instead of shorter. The hand is half as long as the spinal column, and the foot, shorter than the hand, is about  $\frac{1}{11}$ ths of the length of the spinal column.

Thus *Hylobates* is as much longer in the arms than the Gorilla as the Gorilla is longer in the arms than Man; while, on the other hand, it is as much longer in the legs than the Man as the Man is longer in the legs than the Gorilla, so that it contains within itself the extremest deviations from the average length of both pairs of limbs.<sup>1</sup>

The Mandrill presents a middle condition, the arms and legs being nearly equal in length, and both being shorter than the spinal column; while hand and foot have nearly the same proportions to one another and to the spine as in Man.

In the Spider monkey (*Ateles*) the leg is longer than the spine, and the

<sup>1</sup> See the figures of the skeleton of four anthropoid Apes and of Man, drawn to scale, on the following page.



Skeletons of the

CHIMPANZEE.

ORANG.

GORILLA.

MAN.

GIBBON.

Photographically reduced from Diagrams of the natural size (except that of the Gibbon, which was twice as large as nature), drawn by Mr. Waterhouse Hawkins from specimens in the Museum of the Royal College of Surgeons.

arm than the leg ; and, finally, in that remarkable Lemurine form, the Indri (*Lichanotus*), the leg is about as long as the spinal column, while the arm is not more than  $\frac{1}{3}$  of its length, the hand having rather less and the foot rather more than one-third the length of the spinal column.

These examples might be greatly multiplied, but they suffice to show that, in whatever proportion of its limbs the Gorilla differs from Man, the other Apes depart still more widely from the Gorilla, and that, consequently, such differences of proportion can have no ordinal value.

We may next consider the differences presented by the trunk, consisting of the vertebral column, or backbone, and the ribs and pelvis, or bony hip-basin, which are connected with it, in Man and in the Gorilla respectively.

In Man, in consequence partly of the disposition of the articular surfaces of the vertebræ, and largely of the elastic tension of some of the fibrous bands, or ligaments, which connect these vertebræ together, the spinal column, as a whole, has an elegant S-like curvature, being convex forwards in the neck, concave in the back, convex in the loins, or lumbar region, and concave again in the sacral region—an arrangement which gives much elasticity to the whole backbone, and diminishes the jar communicated to the spine, and through it to the head, by locomotion in the erect position.

Furthermore, under ordinary circumstances, Man has seven vertebræ in his neck, which are called *cervical* ; twelve succeed these, bearing ribs and

forming the upper part of the back, whence they are termed *dorsal* ; five lie in the loins, bearing no distinct, or free, ribs, and are called *lumbar* ; five, united together into a great bone, excavated in front, solidly wedged in between the hip bones to form the back of the pelvis, and known by the name of the *sacrum*, succeed these ; and, finally, three or four little more or less movable bones, so small as to be insignificant, constitute the *coccyx* or rudimentary tail.

In the Gorilla the vertebral column is similarly divided into cervical, dorsal, lumbar, sacral, and coccygeal vertebræ, and the total number of cervical and dorsal vertebræ, taken together, is the same as in Man ; but the development of a pair of ribs to the first lumbar vertebra, which is an exceptional occurrence in Man, is the rule in the Gorilla ; and hence, as lumbar are distinguished from dorsal vertebræ only by the presence or absence of free ribs, the seventeen "dorso-lumbar" vertebræ of the Gorilla are divided into thirteen dorsal and four lumbar, while in Man they are twelve dorsal and five lumbar.

Not only, however, does Man occasionally possess thirteen pair of ribs,<sup>1</sup> but the Gorilla sometimes has fourteen pair, while an Orang-Utan skeleton in the Museum of the Royal

<sup>1</sup> "More than once," says Peter Camper, "have I met with more than six lumbar vertebræ in man.... Once I found thirteen ribs and four lumbar vertebræ." Fallopius noted thirteen pair of ribs and only four lumbar vertebræ ; and Eustachius once found eleven dorsal vertebræ and six lumbar vertebræ. — "*Œuvres de Pierre Camper*," T. 1, p. 42. As Tyson states, his "Pygmic" had thirteen pair of ribs and five lumbar vertebræ. The question of the curves of the spinal column in the Apes requires further investigation.



College of Surgeons has twelve dorsal and five lumbar vertebræ, as in Man. Cuvier notes the same number in a dorsal and eight lumbar, and a Lemur (*Stenops tardigradus*) has fifteen dorsal and nine lumbar vertebræ.




FIG. 15.—Front and side views of the bony pelvis of Man, the Gorilla and Gibbon: reduced from drawings made from nature, of the same absolute length, by Mr. Waterhouse Hawkins.

*Hylobates*. On the other hand, among the lower Apes many possess twelve dorsal and six or seven lumbar vertebræ; the Douroucouli has fourteen The vertebral column of the Gorilla, as a whole, differs from that of Man in the less marked character of its curves, especially in the slighter

convexity of the lumbar region. Nevertheless, the curves are present, and are quite obvious in young skeletons of the Gorilla and Chimpanzee which have been prepared without removal of the ligaments. In young Orangs similarly preserved, on the other hand, the spinal column is either straight, or even concave forwards, throughout the lumbar region.

Whether we take these characters, then, or such minor ones as those which are derivable from the proportional length of the spines, of the cervical vertebræ, and the like, there is no doubt whatsoever as to the marked difference between Man and the Gorilla; but there is as little, that equally marked differences, of the very same order, obtain between the Gorilla and the lower Apes.

The Pelvis, or bony girdle of the hips, of Man is a strikingly human part of his organisation, the expanded haunch-bones affording support for his viscera during his habitually erect posture, and giving space for the attachment of the great muscles which enable him to assume and to preserve that attitude. In these respects the pelvis of the Gorilla differs very considerably from his (Fig. 15). But go no lower than the Gibbon, and see how vastly more he differs from the Gorilla than the latter does from Man, even in this structure. Look at the flat, narrow haunch-bones—the long and narrow passage—the coarse, outwardly curved, ischiatic prominences on which the Gibbon habitually rests, and which are coated by the so-called “callosities,” dense patches of skin, wholly absent in the Gorilla, in the Chimpanzee, and in the Orang, as in Man!

In the lower Monkeys and in the Lemurs the difference becomes more striking still, the pelvis acquiring an altogether quadrupedal character.

But now let us turn to a nobler and more characteristic organ—that by which the human frame seems to be, and indeed is, so strongly distinguished from all others—I mean the skull. The differences between a Gorilla's skull and a Man's are truly immense (Fig. 16). In the former the face, formed largely by the massive jaw-bones, predominates over the brain-case, or cranium proper; in the latter the proportions of the two are reversed. In the Man the occipital foramen, through which passes the great nervous cord connecting the brain with the nerves of the body, is placed just behind the centre of the base of the skull, which thus becomes evenly balanced in the erect posture; in the Gorilla, it lies in the posterior third of that base. In the Man the surface of the skull is comparatively smooth, and the supraciliary ridges or brow prominences usually project but little, while in the Gorilla vast crests are developed upon the skull, and the brow ridges overhang the cavernous orbits, like great penthouses.

Sections of the skulls, however, show that some of the apparent defects of the Gorilla's cranium arise, in fact, not so much from deficiency of brain-case as from excessive development of the parts of the face. The cranial cavity is not ill-shaped, and the forehead is not truly flattened or very retreating, its really well-formed curve being simply disguised by the mass of bone which is built up against it (Fig. 16).

But the roofs of the orbits rise more



obliquely into the cranial cavity, thus diminishing the space for the lower part of the anterior lobes of the brain, and the absolute capacity of the cranium is far less than that of Man. So far as I am aware, no human cranium belonging to an adult man has yet been observed with a less cubical capacity than 62 cubic inches, the smallest cranium observed in any race of men, by Morton, measuring 63 cubic inches; while, on the other hand, the most capacious Gorilla skull yet measured has a content of not more than  $34\frac{1}{2}$  cubic inches. Let us assume, for simplicity's sake, that the lowest Man's skull has twice the capacity of that of the highest Gorilla.<sup>1</sup>

<sup>1</sup> It has been affirmed that Hindoo crania sometimes contain as little as 27 ounces of water, which would give a capacity of about 46 cubic inches. The minimum capacity which I have assumed above, however, is based upon the valuable tables published by Professor R. Wagner, in his *Vorstudien zu einer wissenschaftlichen Morphologie und Physiologie des menschlichen Gehirns*. As the result of the careful weighing of more than 900 human brains, Professor Wagner states that one-half weighed between 1,200 and 1,400 grammes, and that about two-ninths, consisting for the most part of male brains, exceed 1,400 grammes. The lightest brain of an adult male, with sound mental faculties, recorded by Wagner, weighed 1,020 grammes. As a gramme equals 15.4 grains, and a cubic inch of water contains 252.4 grains, this is equivalent to 62 cubic inches of water; so that, as brain is heavier than water, we are perfectly safe against erring on the side of diminution in taking this as the smallest capacity of any adult male human brain. The only adult male brain weighing as little as 970 grammes is that of an idiot; but the brain of an adult woman, against the soundness of whose faculties nothing appears, weighed as little as 907 grammes (55.3 cubic inches of water); and Reid gives an adult female brain of still smaller capacity. The heaviest brain (1,872 grammes, or about 115 cubic inches) was, however, that of a woman; next to it comes the brain of Cuvier (1,861 grammes), then Byron (1,807 grammes), and then an insane

No doubt this is a very striking difference, but it loses much of its apparent systematic value when viewed by the light of certain other equally indubitable facts respecting cranial capacities.

The first of these is, that the difference in the volume of the cranial cavity of different races of mankind is far greater, absolutely, than that between the lowest Man and the highest Ape, while, relatively, it is about the same. For the largest human skull measured by Morton contained 114 cubic inches—that is to say, had very nearly double the capacity of the smallest; while its absolute preponderance, of 52 cubic inches, is far greater than that by which the lowest adult male human cranium surpasses the largest of the Gorillas ( $62 - 34\frac{1}{2} = 27\frac{1}{2}$ ). Secondly, the adult crania of Gorillas which have as yet been measured differ among themselves by nearly one-third, the maximum capacity being 34.5 cubic inches, the minimum 24 cubic inches; and, thirdly, after making all due allowance for difference of size, the cranial capacities of some of the lower Apes fall nearly as much, relatively, below those of the higher Apes as the latter fall below Man.

Thus, even in the important matter of cranial capacity, Men differ more widely from one another than they do from the Apes; while the lowest Apes

person (1,783 grammes). The lightest adult brain recorded (720 grammes) was that of an idiotic female. The brains of five children, four years old, weighed between 1,275 and 992 grammes. So that it may be safely said that an average European child of four years old has a brain twice as large as that of an adult Gorilla.



differ as much, in proportion, from the highest as the latter does from Man. The last proposition is still better illustrated by the study of the modifications which other parts of the cranium undergo in the Simian series.

It is the large proportional size of the facial bones and the great projection of the jaws which confer upon the Gorilla's skull its small facial angle and brutal character.

But if we consider the proportional size of the facial bones to the skull proper only, the little *Chrysothrix* (Fig. 16) differs very widely from the Gorilla, and in the same way as Man does; while the Baboons (*Cynocephalus*, Fig. 16) exaggerate the gross proportions of the muzzle of the great Anthropoid, so that its visage looks mild and human by comparison with theirs. The difference between the Gorilla and the Baboon is even greater than it appears at first sight; for the great facial mass of the former is largely due to a downward development of the jaws;




FIG. 16.—Sections of the skulls of Man and various Apes, drawn so as to give the cerebral cavity the same length in each case, thereby displaying the varying proportions of the facial bones. The line *b* indicates the plane of the tentorium, which separates the cerebrum from the cerebellum; *a*, the axis of the occipital outlet of the skull. The extent of cerebral cavity behind *c*, which is a perpendicular erected on *b* at the point where the tentorium is attached posteriorly, indicates the degree to which the cerebrum overlaps the cerebellum—the space occupied by which is roughly indicated by the dark shading. In comparing these diagrams, it must be recollected that figures on so small a scale as these simply exemplify the statements in the text, the proof of which is to be found in the objects themselves.

an essentially human character, super-added upon that almost purely forward, essentially brutal, development of the same parts which characterises the Baboon, and yet more remarkably distinguishes the Lemur.

Similarly, the occipital foramen of *Myctes* (Fig. 16), and still more of the Lemurs, is situated completely in the posterior face of the skull, or as much further back than that of the Gorilla, as that of the Gorilla is further back than that of Man; while, as if to render patent the futility of the attempt to base any broad classificatory distinction on such a character, the same group of Platyrrhine, or American monkeys, to which the *Myctes* belongs, contains the *Chrysotrix*, whose occipital foramen is situated far more forward than in any other Ape, and nearly approaches the position it holds in Man.

Again, the Orang's skull is as devoid of excessively developed supraciliary prominences as a Man's, though some varieties exhibit great crests elsewhere (see p. 32); and in some of the Cebine apes, and in the *Chrysotrix*, the cranium is as smooth and rounded as that of Man himself.

What is true of these leading characteristics of the skull holds good, as may be imagined, of all minor features; so that for every constant difference between the Gorilla's skull and the Man's a similar constant difference of the same order (that is to say, consisting in excess or defect of the same quality) may be found between the Gorilla's skull and that of some other Ape. So that, for the skull no less than for the skeleton in general, the proposition holds good, that the differences between Man and

the Gorilla are of smaller value than those between the Gorilla and some other Apes.

In connection with the skull, I may speak of the teeth—organs which have a peculiar classificatory value, and whose resemblances and differences of number, form, and succession, taken as a whole, are usually regarded as more trustworthy indicators of affinity than any others.

Man is provided with two sets of teeth—milk teeth and permanent teeth. The former consist of four incisors, or cutting teeth; two canines, or eye-teeth; and four molars, or grinders, in each jaw—making twenty in all. The latter (Fig. 17) comprise four incisors, two canines, four small grinders, called premolars or false molars, and six large grinders, or true molars, in each jaw—making thirty-two in all. The internal incisors are larger than the external pair in the upper jaw, smaller than the external pair in the lower jaw. The crowns of the upper molars exhibit four cusps, or blunt-pointed elevations, and a ridge crosses the crown obliquely from the inner anterior cusp to the outer posterior cusp (Fig. 17 *m*<sup>2</sup>). The anterior lower molars have five cusps, three external and two internal. The premolars have two cusps, one internal and one external, of which the outer is the higher.

In all these respects the dentition of the Gorilla may be described in the same terms as that of Man; but in other matters it exhibits many and important differences (Fig. 17).

Thus the teeth of Man constitute a regular and even series—without any break and without any marked



projection of one tooth above the level of the rest; a peculiarity which, as Cuvier long ago showed, is shared by no other mammal save one—as different a creature from Man as can well

*tema*, in both jaws: in front of the eye-tooth, or between it and the outer incisor, in the upper jaw; behind the eye-tooth, or between it and the front false molar, in the lower jaw. Into




FIG. 17.—Lateral views, of the same length, of the upper jaws of various Primates. *i*, incisors; *c*, canines; *pm*, premolars; *m*, molars. A line is drawn through the first molar of Man, Gorilla, Cynocephalus, and Cebus, and the grinding surface of the second molar is shown in each, its anterior and internal angle being just above the *m* of *m*<sup>2</sup>.

be imagined—namely, the long extinct *Anoplotherium*. The teeth of the Gorilla, on the contrary, exhibit a break, or interval, termed the *dias-*

this break in the series in each jaw fits the canine of the opposite jaw; the size of the eye-tooth in the Gorilla being so great that it projects, like a



tusk, far beyond the general level of the other teeth. The roots of the false molar teeth of the Gorilla, again, are more complex than in Man, and the proportional size of the molars is different. The Gorilla has the crown of the hindmost grinder of the lower jaw more complex, and the order of eruption of the permanent teeth is different, the permanent canines making their appearance before the second and third molars in Man, and after them in the Gorilla.

Thus, while the teeth of the Gorilla closely resemble those of Man in number, kind, and in the general pattern of their crowns, they exhibit marked differences from those of Man in secondary respects, such as relative size, number of fangs, and order of appearance.

But if the teeth of the Gorilla be compared with those of an Ape no further removed from it than a *Cynocephalus*, or Baboon, it will be found that differences and resemblances of the same order are easily observable; but that many of the points in which the Gorilla resembles Man are those in which it differs from the Baboon; while various respects in which it differs from Man are exaggerated in the *Cynocephalus*. The number and the nature of the teeth remain the same in the Baboon as in the Gorilla and in Man. But the pattern of the Baboon's upper molars is quite different from that described above (Fig. 17); the canines are proportionally longer and more knife-like; the anterior premolar in the lower jaw is specially modified; the posterior molar of the lower jaw is still larger and more complex than in the Gorilla.

Passing from the Old World Apes to those of the New World, we meet with a change of much greater importance than any of these. In such a genus as *Cebus*, for example (Fig. 17), it will be found that while in some secondary points, such as the projection of the canines and the diastema, the resemblance to the great Ape is preserved; in other and most important respects the dentition is extremely different. Instead of twenty teeth in the milk set, there are twenty-four; instead of thirty-two in the permanent set, there are thirty-six, the false molars being increased from eight to twelve. And in form the crowns of the molars are very unlike those of the Gorilla, and differ far more widely from the human pattern.

The Marmosets, on the other hand, exhibit the same number of teeth as Man and the Gorilla; but, notwithstanding this, their dentition is very different, for they have four more false molars, like the other American monkeys; but, as they have four fewer true molars, the total remains the same. And passing from the American Apes to the Lemurs, the dentition becomes still more completely and essentially different from that of the Gorilla. The incisors begin to vary both in number and in form. The molars acquire, more and more, a many-pointed, insectivorous character, and in one genus, the Aye-Aye (*Cheiromys*), the canines disappear, and the teeth completely simulate those of a Rodent (Fig. 17).

Hence it is obvious that, greatly as the dentition of the highest Ape differs from that of Man, it differs far more widely from that of the lower and lowest Apes.

Whatever part of the animal fabric—whatever series of muscles, whatever viscera—might be selected for comparison, the result would be the same: the lower Apes and the Gorilla would differ more than the Gorilla and the Man. I cannot attempt in this place to follow out all these comparisons in detail, and indeed it is unnecessary I should do so. But certain real or supposed structural distinctions between Man and the Apes remain, upon which so much stress has been laid that they require careful consideration, in order that the true value may be assigned to those which are real, and the emptiness of those which are fictitious may be exposed. I refer to the characters of the hand, the foot, and the brain.

Man has been defined as the only animal possessed of two hands terminating his fore limbs, and of two feet ending his hind limbs, while it has been said that all the Apes possess four hands; and he has been affirmed to differ fundamentally from all the Apes in the characters of his brain, which alone, it has been strangely asserted and re-asserted, exhibits the structures known to anatomists as the posterior lobe, the posterior cornu of the lateral ventricle, and the hippocampus minor.

That the former proposition should have gained general acceptance is not surprising—indeed, at first sight appearances are much in its favour; but, as for the second, one can only admire the surpassing courage of its enunciator, seeing that it is an innovation which is not only opposed to generally and justly accepted doctrines, but which is directly negatived by the testimony of all original

inquirers who have specially investigated the matter; and that it neither has been, nor can be, supported by a single anatomical preparation. It would, in fact, be unworthy of serious refutation, except for the general and natural belief that deliberate and reiterated assertions must have some foundation.

Before we can discuss the first point with advantage we must consider with some attention, and compare together, the structure of the human hand and that of the human foot, so that we may have distinct and clear ideas of what constitutes a hand and what a foot.

The external form of the human hand is familiar enough to everyone. It consists of a stout wrist followed by a broad palm, formed of flesh and tendons and skin, binding together four bones, and dividing into four long and flexible digits, or fingers, each of which bears on the back of its last joint a broad and flattened nail. The longest cleft between any two digits is rather less than half as long as the hand. From the outer side of the base of the palm a stout digit goes off, having only two joints instead of three; so short that it only reaches to a little beyond the middle of the first joint of the finger next it; and further remarkable by its great mobility, in consequence of which it can be directed outwards, almost at a right angle to the rest. This digit is called the *pollex*, or thumb; and, like the others, it bears a flat nail upon the back of its terminal joint. In consequence of the proportions and mobility of the thumb, it is what is termed “oppos-



able"; in other words, its extremity can, with the greatest ease, be brought into contact with the extremities of any of the fingers—a property upon which the possibility of our carrying into effect the conceptions of the mind so largely depends.

fingers; the great toe with the thumb. But the toes, or digits of the foot, are far shorter in proportion than the digits of the hand, and are less movable, the want of mobility being most striking in the great toe—which, again, is very much larger




FIG. 18.—The skeleton of the Hand and Foot of Man, reduced from Dr. Carter's drawings in Gray's *Anatomy*. The hand is drawn to a larger scale than the foot. The line *a a* in the hand indicates the boundary between the carpus and the metacarpus; *b b* that between the latter and the proximal phalanges; *c c* marks the ends of the distal phalanges. The line *a' a'* in the foot indicates the boundary between the tarsus and metatarsus; *b' b'* marks that between the metatarsus and the proximal phalanges; and *c' c'* bounds the ends of the distal phalanges; *ca*, the calcaneum; *as*, the astragalus; *sc*, the scaphoid bone in the tarsus.

The external form of the foot differs widely from that of the hand; and yet, when closely compared, the two present some singular resemblances. Thus the ankle corresponds in a manner with the wrist; the sole with the palm; the toes with the

in proportion to the other toes than the thumb to the fingers. In considering this point, however, it must not be forgotten that the civilised great toe, confined and cramped from childhood upwards, is seen to a great disadvantage, and that in uncivilised



and barefooted people it retains a great amount of mobility, and even some sort of opposability. The Chinese boatmen are said to be able to pull an oar, the artisans of Bengal to weave, and the Carajas to steal fishhooks, by its help; though, after all, it must be recollected that the structure of its joints and the arrangement of its bones necessarily render its prehensile action far less perfect than that of the thumb.

But to gain a precise conception of the resemblances and differences of the hand and foot, and of the distinctive characters of each, we must look below the skin, and compare the bony framework and its motor apparatus in each (Fig. 18).

The skeleton of the hand exhibits, in the region which we term the wrist, and which is technically called the *carpus*, two rows of closely fitted polygonal bones, four in each row, which are tolerably equal in size. The bones of the first row with the bones of the forearm form the wrist joint, and are arranged side by side, no one greatly exceeding or overlapping the rest.

The four bones of the second row of the carpus bear the four long bones which support the palm of the hand. The fifth bone of the same character is articulated in a much more free and movable manner than the others, with its carpal bone, and forms the base of the thumb. These are called *metacarpal* bones, and they carry the *phalanges*, or bones of the digits, of which there are two in the thumb and three in each of the fingers.

The skeleton of the foot is very like that of the hand in some respects. Thus there are three phalanges in

each of the lesser toes, and only two in the great toe, which answers to the thumb. There is a long bone, termed *metatarsal*, answering to the metacarpal, for each digit; and the *tarsus*, which corresponds with the carpus, presents four short polygonal bones in a row, which correspond very closely with the four carpal bones of the second row of the hand. In other respects the foot differs very widely from the hand. Thus the great toe is the longest digit but one; and its metatarsal is far less movably articulated with the tarsus than the metacarpal of the thumb with the carpus. But a far more important distinction lies in the fact that, instead of four more tarsal bones, there are only three; and that these three are not arranged side by side, or in one row. One of them, the *os calcis* or heel bone (*ca*), lies externally, and sends back the large projecting heel; another, the *astragalus* (*as*), rests on this by one face, and, by another, forms, with the bones of the leg, the ankle-joint; while a third face, directed forwards, is separated from the three inner tarsal bones of the row next the metatarsus by a bone called the *scaphoid* (*sc*).

Thus there is a fundamental difference in the structure of the foot and the hand, observable when the carpus and the tarsus are contrasted; and there are differences of degree noticeable when the proportions and the mobility of the metacarpals and metatarsals, with their respective digits, are compared together.

The same two classes of differences become obvious when the muscles of the hand are compared with those of the foot.

Three principal sets of muscles, called "flexors," bend the fingers and thumb, as in clenching the fist, and three sets—the extensors—extend them, as in straightening the fingers. These muscles are all "long muscles"; that is to say, the fleshy part of each, lying in and being fixed to the bones of the arm, is, at the other end, continued into tendons, or rounded cords, which pass into the hand, and are ultimately fixed to the bones which are to be moved. Thus, when the fingers are bent, the fleshy parts of the flexors of the fingers, placed in the arm, contract, in virtue of their peculiar endowment as muscles; and, pulling the tendinous cords connected with their ends, cause them to pull down the bones of the fingers towards the palm.

Not only are the principal flexors of the fingers and of the thumb long muscles, but they remain quite distinct from one another throughout their whole length.

In the foot there are also three principal flexor muscles of the digits or toes, and three principal extensors; but one extensor and one flexor are short muscles; that is to say, their fleshy parts are not situated in the leg (which corresponds with the arm), but in the back and in the sole of the foot—regions which correspond with the back and the palm of the hand.

Again, the tendons of the long flexor of the toes, and of the long flexor of the great toe, when they reach the sole of the foot, do not remain distinct from one another, as the flexors in the palm of the hand do, but they become united and commingled in a very curious manner; while their united tendons receive an

accessory muscle connected with the heel-bone.

But perhaps the most absolutely distinctive character about the muscles of the foot is the existence of what is termed the *peronæus longus*, a long muscle fixed to the outer bone of the leg, and sending its tendon to the outer ankle, behind and below which it passes, and then crosses the foot obliquely to be attached to the base of the great toe. No muscle in the hand exactly corresponds with this, which is eminently a foot muscle.

To resume, the foot of man is distinguished from his hand by the following absolute anatomical differences:—

1. By the arrangement of the tarsal bones.
2. By having a short flexor and a short extensor muscle of the digits.
3. By possessing the muscle termed *peronæus longus*.

And if we desire to ascertain whether the terminal division of a limb in other Primates is to be called a foot or a hand, it is by the presence or absence of these characters that we must be guided, and not by the mere proportions and greater or lesser mobility of the great toe, which may vary indefinitely without any fundamental alteration in the structure of the foot.

Keeping these considerations in mind, let us now turn to the limbs of the Gorilla. The terminal division of the fore limb presents no difficulty—bone for bone and muscle for muscle are found to be arranged essentially as in man, or with such minor differences as are found as varieties in man. The Gorilla's hand is clumsier, heavier, and has a thumb somewhat shorter in proportion than that of man; but no



one has ever doubted its being a true hand.

At first sight the termination of the hind limb of the Gorilla looks very hand-like, and, as it is still more so in many of the lower Apes, it is not wonderful that the appellation "*Quadrumanus*," or four-handed creatures, adopted from the older anatomists<sup>1</sup> by Blumenbach, and unfortunately rendered current by Cuvier, should have gained such wide acceptance as a name for the Simian group. But the most cursory anatomical investigation at once proves that the resemblance of the so-called "hind hand" to a true hand is only skin deep, and that, in all essential respects, the hind limb of the Gorilla is as truly terminated by a foot as that of Man. The tarsal bones, in all important circumstances of number, disposition, and form, resemble those of Man (Fig. 19). The metatarsals and digits, on the other hand, are proportionally longer and more slender, while the great toe is not only proportionally shorter and weaker, but its metatarsal bone is united by a more movable

joint with the tarsus. At the same time the foot is set more obliquely upon the leg than in man.

As to the muscles, there is a short flexor, a short extensor, and a *peronæus longus*, while the tendons of the long flexors of the great toe and of the other toes are united together and with an accessory fleshy bundle.

The hind limb of the Gorilla, therefore, ends in a true foot, with a very movable great toe. It is a prehensile foot, indeed, but is in no sense a hand; it is a foot which differs from that of Man not in any fundamental character, but in mere proportions, in the degree of mobility, and in the secondary arrangement of its parts.

It must not be supposed, however, because I speak of these differences as not fundamental, that I wish to underrate their value. They are important enough in their way, the structure of the foot being in strict correlation with that of the rest of the organism in each case. Nor can it be doubted that the greater division of physiological labour in Man, so that the function of support is thrown wholly on the leg and foot, is an advance in organisation of very great moment to him; but, after all, regarded anatomically, the resemblances between the foot of Man and the foot of the Gorilla are far more striking and important than the differences.

I have dwelt upon this point at length, because it is one regarding which much delusion prevails; but I might have passed it over without detriment to my argument, which only requires me to show that, be the differences between the hand and foot

<sup>1</sup> In speaking of the foot of his "*Pygmie*," Tyson remarks, p. 13:—

"But this part in the formation and in its function, too, being liker a Hand than a Foot: for the distinguishing this sort of animals from others, I have thought whether it might not be reckoned and called rather *Quadrumanus* than *Quadrupes*—*i.e.*, a four-handed rather than a four-footed animal."

As this passage was published in 1699, M. I. G. St. Hilaire is clearly in error in ascribing the invention of the term "*quadrumanous*" to Buffon, though "*bimanous*" may belong to him. Tyson uses "*Quadrumanus*" in several places, as at p. 91..... "*Our Pygmie* is no Man, nor yet the *common Ape*, but a sort of *Animal* between both; and though a *Biped*, yet of the *Quadrumanus*-kind: though some *Men*, too, have been observed to use their *Feet* like *Hands*, as I have seen several."



of Man and those of the Gorilla what they may, the differences between those of the Gorilla and those of the lower Apes are much greater.

It is not necessary to descend lower in the scale than the Orang for conclusive evidence on this head.

The thumb of the Orang differs more from that of the Gorilla than the thumb of the Gorilla differs from that of Man, not only by its short-

of articulation in the leg, and absence of a long flexor tendon to the great toe, separating it far more widely from the foot of the Gorilla than the latter is separated from that of Man.

But, in some of the lower Apes, the hand and foot diverge still more from those of the Gorilla than they do in the Orang. The thumb ceases to be opposable in the American monkeys ; is reduced to a mere rudiment covered




FIG. 19.—Foot of Man, Gorilla, and Orang-Utan of the same absolute length, to show the differences in proportion of each. Letters as in Fig. 18. Reduced from original drawings by Mr. Waterhouse Hawkins.

ness, but by the absence of any special long flexor muscle. The carpus of the Orang, like that of most lower Apes, contains nine bones, while in the Gorilla, as in Man and the Chimpanzee, there are only eight.

The Orang's foot (Fig. 19) is still more aberrant ; its very long toes and short tarsus, short great toe, short and raised heel, great obliquity

by the skin in the Spider Monkey ; and is directed forwards and armed with a curved claw, like the other digits, in the Marmosets ; so that, in all these cases, there can be no doubt but that the hand is more different from that of the Gorilla than the Gorilla's hand is from Man's.

And as to the foot, the great toe of the Marmoset is still more insignificant

in proportion than that of the Orang; while in the Lemurs it is very large, and as completely thumb-like and opposable as in the Gorilla; but in these animals the second toe is often irregularly modified, and in some species the two principal bones of the tarsus, the *astragalus* and the *os calcis*, are so immensely elongated as to render the foot, so far, totally unlike that of any other mammal.

So with regard to the muscles. The short flexor of the toes of the Gorilla differs from that of Man by the circumstance that one slip of the muscle is attached, not to the heel bone, but to the tendons of the long flexors. The lower Apes depart from the Gorilla by an exaggeration of the same character, two, three, or more slips becoming fixed to the long flexor tendons—or by a multiplication of the slips. Again, the Gorilla differs slightly from Man in the mode of interlacing of the long flexor tendons; and the lower Apes differ from the Gorilla in exhibiting yet other, sometimes very complex, arrangements of the same parts, and occasionally in the absence of the accessory fleshy bundle.

Throughout all these modifications it must be recollected that the foot loses no one of its essential characters. Every Monkey and Lemur exhibits the characteristic arrangement of tarsal bones, possesses a short flexor and short extensor muscle, and a *peronæus longus*. Varied as the proportions and appearance of the organ may be, the terminal division of the hind limb remains, in plan and principle of construction, a foot, and never, in those respects, can be confounded with a hand.

Hardly any part of the bodily frame, then, could be found better calculated to illustrate the truth that the structural differences between Man and the highest Ape are of less value than those between the highest and the lower Apes, than the hand or the foot; and yet, perhaps, there is one organ the study of which enforces the same conclusion in a still more striking manner—and that is the Brain.

But before entering upon the precise question of the amount of difference between the Ape's brain and that of Man, it is necessary that we should clearly understand what constitutes a great and what a small difference in cerebral structure; and we shall be best enabled to do this by a brief study of the chief modifications which the brain exhibits in the series of vertebrate animals.

The brain of a fish is very small compared with the spinal cord into which it is continued, and with the nerves which come off from it. Of the segments of which it is composed—the olfactory lobes, the cerebral hemispheres, and the succeeding divisions—no one predominates so much over the rest as to obscure or cover them; and the so-called optic lobes are, frequently, the largest masses of all. In Reptiles the mass of the brain, relatively to the spinal cord, increases, and the cerebral hemispheres begin to predominate over the other parts; while in Birds this predominance is still more marked. The brain of the lowest Mammals, such as the duck-billed Platypus and the Opossums and Kangaroos, exhibits a still more definite advance in the same direction. The cerebral hemispheres have now so much increased in size as, more or



less, to hide the representatives of the optic lobes, which remain comparatively small, so that the brain of a Marsupial is extremely different from that of a Bird, Reptile, or Fish. A step higher in the scale, among the placental Mammals, the structure of the brain acquires a vast modification—not that it appears much altered externally, in a Rat or in a Rabbit, from what it is in a Marsupial, nor that the proportions of its parts are much changed; but an apparently new structure is found between the cerebral hemispheres connecting them together, as what is called the “great commissure” or “corpus callosum.” The subject requires careful re-investigation; but, if the currently received statements are correct, the appearance of the “corpus callosum” in the placental Mammals is the greatest and most sudden modification exhibited by the brain in the whole series of vertebrated animals—it is the greatest leap anywhere made by Nature in her brain work. For the two halves of the brain being once thus knit together, the progress of cerebral complexity is traceable through a complete series of steps from the lowest Rodent, or Insectivore, to Man; and that complexity consists chiefly in the disproportionate development of the cerebral hemispheres and of the cerebellum, but especially of the former, in respect to the other parts of the brain.

In the lower placental Mammals the cerebral hemispheres leave the proper upper and posterior face of the cerebellum completely visible when the brain is viewed from above, but in the higher forms the hinder part of each hemisphere, separated only by the tentorium

(p. 68) from the anterior face of the cerebellum, inclines backwards and downwards, and grows out, as the so-called “posterior lobe,” so as at length to overlap and hide the cerebellum. In all Mammals each cerebral hemisphere contains a cavity which is termed the “ventricle,” and as this ventricle is prolonged, on the one hand forwards, and on the other downwards, into the substance of the hemisphere, it is said to have two horns or “cornua,” an “anterior cornu” and a “descending cornu.” When the posterior lobe is well developed a third prolongation of the ventricular cavity extends into it, and is called the “posterior cornu.”

In the lower and smaller forms of placental Mammals the surface of the cerebral hemispheres is either smooth or evenly rounded, or exhibits a very few grooves, which are technically termed “sulci,” separating ridges or “convolutions” of the substance of the brain; and the smaller species of all orders tend to a similar smoothness of brain. But in the higher orders, and especially the larger members of these orders, the grooves, or sulci, become extremely numerous, and the intermediate convolutions proportionately more complicated in their meanderings, until, in the Elephant, the Porpoise, the higher Apes, and Man, the cerebral surface appears a perfect labyrinth of tortuous foldings.

Where a posterior lobe exists and presents its customary cavity—the posterior cornu—it commonly happens that a particular sulcus appears upon the inner and under surface of the lobe, parallel with and beneath the floor of the cornu, which is, as it were, arched over the roof of the sulcus. It is as if

the groove had been formed by indenting the floor of the posterior horn from without with a blunt instrument, so that the floor should rise as a convex eminence. Now, this eminence is what has been termed the "hippocampus minor," the "hippocampus major" being a larger eminence in the floor of the descending cornu. What may be the functional importance of either of these structures we know not.

As if to demonstrate, by a striking example, the impossibility of erecting any cerebral barrier between Man and the Apes, Nature has provided us, in the latter animals, with an almost complete series of gradations from brains little higher than that of a Rodent to brains little lower than that of Man. And it is a remarkable circumstance that though, so far as our present knowledge extends, there *is* one true structural break in the series of forms of Simian brains, this hiatus does not lie between Man and the man-like Apes, but between the lower and the lowest Simians; or, in other words, between the Old and New World apes and monkeys and the Lemur. Every Lemur which has yet been examined, in fact, has its cerebellum partially visible from above, and its posterior lobe, with the contained posterior cornu and hippocampus minor, more or less rudimentary. Every Marmoset, American monkey, Old World monkey, Baboon, or man-like Ape, on the contrary, has its cerebellum entirely hidden, posteriorly, by the cerebral lobes, and possesses a large posterior cornu, with a well-developed hippocampus minor.

In many of these creatures, such as the Saimiri (*Chrysothrix*), the cerebral lobes overlap and extend much further behind the cerebellum, in proportion, than they do in Man (Fig. 16); and it is quite certain that in all the cerebellum is completely covered behind by well-developed posterior lobes. The fact can be verified by everyone who possesses the skull of any Old or New World monkey. For, inasmuch as the brain in all mammals completely fills the cranial cavity, it is obvious that a cast of the interior of the skull will reproduce the general form of the brain, at any rate with such minute and, for the present purpose, utterly unimportant differences as may result from the absence of the enveloping membranes of the brain in the dry skull. But if such a cast be made in plaster, and compared with a similar cast of the interior of a human skull, it will be obvious that the cast of the cerebral chamber, representing the cerebrum of the ape, as completely covers over and overlaps the cast of the cerebellar chamber, representing the cerebellum, as it does in the man (Fig. 20). A careless observer, forgetting that a soft structure like the brain loses its proper shape the moment it is taken out of the skull, may indeed mistake the uncovered condition of the cerebellum of an extracted and distorted brain for the natural relations of the parts; but his error must become patent even to himself if he try to replace the brain within the cranial chamber. To suppose that the cerebellum of an ape is naturally uncovered behind is a miscomprehension comparable only to that of one who should imagine that a man's lungs always



occupy but a small portion of the thoracic cavity, because they do so when the chest is opened, and their elasticity is no longer neutralised by the pressure of the air.

skull of any ape above a Lemur, without taking the trouble to make a cast of it. For there is a very marked groove in every such skull, as in the human skull, which indi-




FIG. 20.—Drawings of the internal casts of a Man's and of a Chimpanzee's skull, of the same absolute length, and placed in corresponding positions. *A.* Cerebrum; *B.* Cerebellum. The former drawing is taken from a cast in the Museum of the Royal College of Surgeons, the latter from the photograph of the cast of a Chimpanzee's skull, which illustrates the paper by Mr. Marshall "On the Brain of the Chimpanzee" in the *Natural History Review* for July, 1861. The sharper definition of the lower edge of the cast of the cerebral chamber in the Chimpanzee arises from the circumstance that the tentorium remained in that skull and not in the Man's. The cast more accurately represents the brain in the Chimpanzee than in the Man; and the great backward projection of the posterior lobes of the cerebrum of the former, beyond the cerebellum, is conspicuous.

And the error is the less excusable, as it must become apparent to every one who examines a section of the

skull of any ape above a Lemur, without taking the trouble to make a cast of it. For there is a very marked groove in every such skull, as in the human skull, which indicates the line of attachment of what is termed the *tentorium*—a sort of parchment-like shelf, or partition, which, in the recent state, is interposed between the cerebrum and cerebellum, and prevents the former from pressing upon the latter (see Fig. 16).

This groove, therefore, indicates the line of separation between that part of the cranial cavity which contains the cerebrum and that which contains the cerebellum; and as the brain exactly fills the cavity of the skull, it is obvious that the relations of these two parts of the cranial cavity at once inform us of the relations of their contents. Now, in man, in all the Old World, and in all the New World *Simiæ*, with one exception, when the face is directed forwards, this line of attachment of the tentorium, or impression for the lateral sinus, as it is technically called, is nearly horizontal, and the cerebral chamber in-

variably overlaps or projects behind the cerebellar chamber. In the Howler Monkey, or *Myctes* (see Fig. 16), the

line passes obliquely upwards and backwards, and the cerebral overlap is almost nil; while in the Lemurs, as in the lower Mammals, the line is much more inclined in the same direction, and the cerebellar chamber projects considerably beyond the cerebral.

When the gravest errors respecting points so easily settled as this question respecting the posterior lobes can be authoritatively propounded, it is no wonder that matters of observation of no very complex character, but still requiring a certain amount of care, should have fared worse. Anyone who cannot see the posterior lobe in an ape's brain is not likely to give a very valuable opinion respecting the posterior cornu or the hippocampus minor. If a man cannot see a church, it is preposterous to take his opinion about its altar-piece or painted window; so that I do not feel bound to enter upon any discussion of these points, but content myself with assuring the reader that the posterior cornu and the hippocampus minor have now been seen—usually at least as well developed as in man, and often better—not only in the Chimpanzee, the Orang, and the Gibbon, but in all the genera of the Old World baboons and monkeys, and in most of the New World forms, including the Marmosets.<sup>1</sup>

In fact, all the abundant and trustworthy evidence (consisting of the results of careful investigations directed to the determination of these very questions, by skilled anatomists)

which we now possess leads to the conviction that, so far from the posterior lobe, the posterior cornu, and the hippocampus minor being structures peculiar to and characteristic of Man, as they have been over and over again asserted to be, even after the publication of the clearest demonstration of the reverse, it is precisely these structures which are the most marked cerebral characters common to Man with the Apes. They are among the most distinctly Simian peculiarities which the human organism exhibits.

As to the convolutions, the brains of the Apes exhibit every stage of progress, from the almost smooth brain of the Marmoset to the Orang and the Chimpanzee, which fall but little below Man. And it is most remarkable that, as soon as all the principal sulci appear, the pattern according to which they are arranged is identical with that of the corresponding sulci of Man. The surface of the brain of a monkey exhibits a sort of skeleton map of man's, and in the man-like Apes the details become more and more filled in, until it is only in minor characters, such as the greater excavation of the anterior lobes, the constant presence of fissures usually absent in man, and the different disposition and proportions of some convolutions, that the Chimpanzee's or the Orang's brain can be structurally distinguished from Man's.

So far as cerebral structure goes, therefore, it is clear that Man differs less from the Chimpanzee or the Orang than these do even from the Monkeys, and that the difference between the brains of the Chimpanzee and of Man is almost insignificant

<sup>1</sup> See the note at the end of this essay for a succinct history of the controversy to which allusion is here made.



when compared with that between the Chimpanzee brain and that of a Lemur.

It must not be overlooked, however, that there is a very striking difference in the absolute mass and weight between the lowest human brain and that of the highest ape—a difference which is all the more remarkable when we recollect that a full-grown Gorilla is probably pretty nearly twice as heavy as a Bosjes man, or as many an European woman. It may be doubted whether a healthy human adult brain ever weighed less than thirty-one or two ounces, or that the heaviest Gorilla brain has exceeded twenty ounces.

This is a very noteworthy circumstance, and doubtless will one day help to furnish an explanation of the great gulf which intervenes between the lowest man and the highest ape in intellectual power;<sup>1</sup> but it has

<sup>1</sup> I say *help* to furnish, for I by no means believe that it was any original difference of cerebral quality, or quantity, which caused that divergence between the human and the pithecoïd stirpes which has ended in the present enormous gulf between them. It is no doubt perfectly true, in a certain sense, that all difference of function is a result of difference of structure; or, in other words, of difference in the combination of the primary molecular forces of living substance; and, starting from this undeniable axiom, objectors occasionally, and with much seeming plausibility, argue that the vast intellectual chasm between the Ape and Man implies a corresponding structural chasm in the organs of the intellectual functions; so that, it is said, the non-discovery of such vast differences proves, not that they are absent, but that Science is incompetent to detect them. A very little consideration, however, will, I think, show the fallacy of this reasoning. Its validity hangs upon the assumption that intellectual power depends altogether on the brain—whereas the brain is only one condition




FIG. 21.—Drawings of the cerebral hemispheres of a Man and of a Chimpanzee, of the same length, in order to show the relative proportions of the parts: the former taken from a specimen which Mr. Flower, Conservator of the Museum of the Royal College of Surgeons, was good enough to dissect for me; the latter, from the photograph of a similarly dissected Chimpanzee's brain, given in Mr. Marshall's paper above referred to. *a*, posterior lobe; *b*, lateral ventricle; *c*, posterior cornu; *x*, the hippocampus minor.

little systematic value, for the simple reason that, as may be concluded from what has been already said respecting cranial capacity, the difference in weight of brain between the highest and the lowest men is far greater, both relatively and absolutely, than that between the lowest man and the highest ape. The latter, as has been seen, is represented by, say, twelve ounces of cerebral sub-

stance absolutely, or by 32 : 20 relatively; but, as the largest recorded human brain weighed between 65 and 66 ounces, the former difference is represented by more than 33 ounces absolutely, or by 65 : 32 relatively. Regarded systematically, the cerebral differences of Man and Apes are not of more than generic value—his Family distinction resting chiefly on his dentition, his pelvis, and his lower limbs.

out of many on which intellectual manifestations depend; the others being, chiefly, the organs of the senses and the motor apparatuses, especially those which are concerned in prehension and in the production of articulate speech.

A man born dumb, notwithstanding his great cerebral mass and his inheritance of strong intellectual instincts, would be capable of few higher intellectual manifestations than an Orang or a Chimpanzee if he were confined to the society of dumb associates. And yet there might not be the slightest discernible difference between his brain and that of a highly intelligent and cultivated person. The dumbness might be the result of a defective structure of the mouth, or of the tongue, or a mere defective innervation of these parts; or it might result from congenital deafness, caused by some minute defect of the internal ear, which only a careful anatomist could discover.

The argument that, because there is an immense difference between a Man's intelligence and an Ape's, therefore there must be an equally immense difference between their brains, appears to me to be about as well based as the reasoning by which one should endeavour to prove that, because there is a "great gulf" between a watch that keeps accurate time and another that will not go at all, there is therefore a great structural hiatus between the two watches. A hair in the balance-wheel, a little rust on a pinion, a bend in a tooth of the escapement, a something so slight that only the practised eye of the watchmaker can discover it, may be the source of all the difference.

And believing, as I do, with Cuvier, that the possession of articulate speech is the grand distinctive character of man (whether it be absolutely peculiar to him or not), I find it very easy to comprehend that some equally inconspicuous structural difference may have been the primary cause of the immeasurable and practically infinite divergence of the Human from the Simian stirps.

Thus, whatever system of organs be studied, the comparison of their modifications in the ape series leads to one and the same result—that the structural differences which separate Man from the Gorilla and the Chimpanzee are not so great as those which separate the Gorilla from the lower Apes.

But in enunciating this important truth I must guard myself against a form of misunderstanding which is very prevalent. I find, in fact, that those who endeavour to teach what Nature so clearly shows us in this matter are liable to have their opinions misrepresented and their phraseology garbled, until they seem to say that the structural differences between Man and even the highest Apes are small and insignificant. Let me take this opportunity, then, of distinctly asserting, on the contrary, that they are great and significant; that every bone of a Gorilla bears marks by which it might be distinguished from the corresponding bone of a Man; and that, in the present creation at any rate, no intermediate link bridges over the gap between *Homo* and *Troglo-*  
*lodytes*.

It would be no less wrong than



absurd to deny the existence of this chasm; but it is at least equally wrong and absurd to exaggerate its magnitude, and, resting on the admitted fact of its existence, to refuse to inquire whether it is wide or narrow. Remember, if you will, that there is no existing link between Man and the Gorilla; but do not forget that there is a no less sharp line of demarcation, a no less complete absence of any transitional form, between the Gorilla and the Orang or the Orang and the Gibbon. I say, not less sharp, though it is somewhat narrower. The structural differences between Man and the man-like Apes certainly justify our regarding him as constituting a family apart from them; though, inasmuch as he differs less from them than they do from other families of the same order, there can be no justification for placing him in a distinct order.

And thus the sagacious foresight of the great lawgiver of systematic zoology, Linnæus, becomes justified, and a century of anatomical research brings us back to his conclusion, that Man is a member of the same order (for which the Linnæan term PRIMATES ought to be retained) as the Apes and Lemurs. This order is now divisible into seven families, of about equal systematic value: the first, the ANTHROPINI, contains Man alone; the second, the CATARRHINI, embraces the Old World apes; the third, the PLATYRHINI, all New World apes, except the Marmosets; the fourth, the ARCTOPITHECINI, contains the Marmosets; the fifth, the LEMURINI, the Lemurs—from which *Cheiromys* should probably be excluded to form a sixth distinct family, the CHEIROMYINI; while the seventh, the GALEOPITHECINI,

contains only the flying Lemur *Galeopithecus*—a strange form which almost touches on the Bats, as the *Cheiromys* puts on a Rodent clothing, and the Lemurs simulate Insectivora.

Perhaps no order of mammals presents us with so extraordinary a series of gradations as this, leading us insensibly from the crown and summit of the animal creation down to creatures, from which there is but a step, as it seems, to the lowest, smallest, and least intelligent of the placental Mammalia. It is as if Nature herself had foreseen the arrogance of man, and with Roman severity had provided that his intellect, by its very triumphs, should call into prominence the slaves admonishing the conqueror that he is but dust.

These are the chief facts, this the immediate conclusion from them, to which I adverted in the commencement of this Essay. The facts, I believe, cannot be disputed; and, if so, the conclusion appears to me to be inevitable.

But if Man be separated by no greater structural barrier from the brutes than they are from one another, then it seems to follow that, if any process of physical causation can be discovered by which the genera and families of ordinary animals have been produced, that process of causation is amply sufficient to account for the origin of Man. In other words, if it could be shown that the Marmosets, for example, have arisen by gradual modification of the ordinary Platyrrhini, or that both Marmosets and Platyrrhini are modified ramifications of a primitive stock, then there would be no rational ground for doubting

that Man might have originated, in the one case, by the gradual modification of a man-like Ape ; or, in the other case, as a ramification of the same primitive stock as those apes.

At the present moment but one such process of physical causation has any evidence in its favour ; or, in other words, there is but one hypothesis regarding the origin of species of animals in general which has any scientific existence—that propounded by Mr. Darwin. For Lamarck, sagacious as many of his views were, mingled them with so much that was crude and even absurd as to neutralise the benefit which his originality might have effected had he been a more sober and cautious thinker ; and, though I have heard of the announcement of a formula touching “the ordained continuous becoming of organic forms,” it is obvious that it is the first duty of a hypothesis to be intelligible, and that a *qua-quâ-versal* proposition of this kind, which may be read backwards or forwards or sideways with exactly the same amount of signification, does not really exist, though it may seem to do so.

At the present moment, therefore, the question of the relation of Man to the lower animals resolves itself, in the end, into the larger question of the tenability or untenability of Mr. Darwin’s views. But here we enter upon difficult ground, and it behoves us to define our exact position with the greatest care.

It cannot be doubted, I think, that Mr. Darwin has satisfactorily proved that what he terms selection, or selective modification, must occur, and does occur, in nature ; and he has also proved to superfluity that such

selection is competent to produce forms as distinct, structurally, as some genera even are. If the animated world presented us with none but structural differences, I should have no hesitation in saying that Mr. Darwin had demonstrated the existence of a true physical cause, amply competent to account for the origin of living species, and of Man among the rest.

But, in addition to their structural distinctions, the species of animals and plants, or at least a great number of them, exhibit physiological characters—what are known as distinct species, structurally, being for the most part either altogether incompetent to breed one with another ; or, if they breed, the resulting mule, or hybrid, is unable to perpetuate its race with another hybrid of the same kind.

A true physical cause is, however, admitted to be such only on one condition—that it shall account for all the phenomena which come within the range of its operation. If it is inconsistent with any one phenomenon, it must be rejected ; if it fails to explain any one phenomenon, it is so far weak, so far to be suspected, though it may have a perfect right to claim provisional acceptance.

Now, Mr. Darwin’s hypothesis is not, so far as I am aware, inconsistent with any known biological fact ; on the contrary, if admitted, the facts of Development, of Comparative Anatomy, of Geographical Distribution, and of Palæontology, become connected together, and exhibit a meaning such as they never possessed before ; and I for one am fully convinced that, if not precisely true, that



hypothesis is as near an approximation to the truth as, for example, the Copernican hypothesis was to the true theory of the planetary motions.

But, for all this, our acceptance of the Darwinian hypothesis must be provisional so long as one link in the chain of evidence is wanting; and so long as all the animals and plants certainly produced by selective breeding from a common stock are fertile, and their progeny are fertile with one another, that link will be wanting. For so long selective breeding will not be proved to be competent to do all that is required of it to produce natural species.

I have put this conclusion as strongly as possible before the reader, because the last position in which I wish to find myself is that of an advocate for Mr. Darwin's or any other views—if by an advocate is meant one whose business it is to smooth over real difficulties and to persuade where he cannot convince.

In justice to Mr. Darwin, however, it must be admitted that the conditions of fertility and sterility are very ill understood, and that every day's advance in knowledge leads us to regard the hiatus in his evidence as of less and less importance when set against the multitude of facts which harmonise with, or receive an explanation from, his doctrines.

I adopt Mr. Darwin's hypothesis, therefore, subject to the production of proof that physiological species may be produced by selective breeding, just as a physical philosopher may accept the undulatory theory of light, subject to the proof of the existence of the hypothetical ether; or as the chemist adopts the atomic theory,

subject to the proof of the existence of atoms; and for exactly the same reasons—namely, that it has an immense amount of *prima facie* probability; that it is the only means at present within reach of reducing the chaos of observed facts to order; and, lastly, that it is the most powerful instrument of investigation which has been presented to naturalists since the invention of the natural system of classification and the commencement of the systematic study of embryology.

But, even leaving Mr. Darwin's views aside, the whole analogy of natural operations furnishes so complete and crushing an argument against the intervention of any but what are termed secondary causes in the production of all the phenomena of the universe that, in view of the intimate relations between Man and the rest of the living world, and between the forces exerted by the latter and all other forces, I can see no excuse for doubting that all are co-ordinated terms of Nature's great progression, from the formless to the formed—from the inorganic to the organic—from blind force to conscious intellect and will.

Science has fulfilled her function when she has ascertained and enunciated truth; and were these pages addressed to men of science only, I should now close this Essay, knowing that my colleagues have learned to respect nothing but evidence, and to believe that their highest duty lies in submitting to it, however it may jar against their inclinations.

But desiring, as I do, to reach the wider circle of the intelligent public, it would be unworthy cowardice were

I to ignore the repugnance with which the majority of my readers are likely to meet the conclusions to which the most careful and conscientious study I have been able to give to this matter has led me.

On all sides I shall hear the cry: "We are men and women, not a mere better sort of apes, a little longer in the leg, more compact in the foot, and bigger in brain than your brutal Chimpanzees and Gorillas. The power of knowledge—the conscience of good and evil—the pitiful tenderness of human affections, raise us out of all real fellowship with the brutes, however closely they may seem to approximate to us."

To this I can only reply that the exclamation would be more just, and would have my own entire sympathy, if it were only relevant. But it is not I who seek to base Man's dignity upon his great toe, or insinuate that we are lost if an Ape has a hippocampus minor. On the contrary, I have done my best to sweep away this vanity. I have endeavoured to show that no absolute structural line of demarcation wider than that between the animals which immediately succeed us in the scale can be drawn between the animal world and ourselves; and I may add the expression of my belief that the attempt to draw a psychical distinction is equally futile, and that even the highest faculties of feeling and of intellect begin to germinate in lower forms of life.<sup>1</sup> At the same time, no one is

more strongly convinced than I am of the vastness of the gulf between civilised man and the brutes; or is more certain that, whether *from* them or not, he is assuredly not *of* them. No one is less disposed to think lightly of the present dignity, or despairingly of the future hopes, of the only consciously intelligent denizen of this world.

We are indeed told by those who assume authority in these matters that the two sets of opinions are incompatible, and that the belief in the unity of origin of man and brutes involves the brutalisation and degradation of the former. But is this really so? Could not a sensible child confute, by obvious arguments, the shallow rhetoricians who would force this conclusion upon us? Is it, indeed, true that the Poet, or the Philosopher, or the Artist whose genius is the glory of his age, is degraded from his high estate by the undoubted historical probability, not to say certainty, that he is the direct

*of the Linnean Society of London for 1857, but is unaccountably omitted in the "Reade Lecture" delivered before the University of Cambridge two years later, which is otherwise nearly a reprint of the paper in question. Professor Owen writes:—*

"Not being able to appreciate or conceive of the distinction between the psychical phenomena of a Chimpanzee and of a Boschisman or of an Aztec, with arrested brain-growth, as being of a nature so essential as to preclude a comparison between them, or as being other than a difference of degree, I cannot shut my eyes to the significance of that all-pervading similitude of structure—every tooth, every bone, strictly homologous—which makes the determination of the difference between *Homo* and *Pithecius* the anatomist's difficulty."

Surely it is a little singular that the "anatomist" who finds it "difficult" to "determine the difference" between *Homo* and *Pithecius* should yet range them, on anatomical grounds, in distinct sub-classes!

<sup>1</sup> It is so rare a pleasure for me to find Professor Owen's opinions in entire accordance with my own that I cannot forbear from quoting a paragraph which appeared in his Essay "On the Characters, &c., of the Class Mammalia," in the *Journal of the Proceedings*



descendant of some naked and bestial savage, whose intelligence was just sufficient to make him a little more cunning than the Fox, and by so much more dangerous than the Tiger? Or is he bound to howl and grovel on all fours because of the wholly unquestionable fact that he was once an egg, which no ordinary power of discrimination could distinguish from that of a Dog? Or is the philanthropist or the saint to give up his endeavours to lead a noble life because the simplest study of man's nature reveals, at its foundations, all the selfish passions and fierce appetites of the merest quadruped? Is mother-love vile because a hen shows it, or fidelity base because dogs possess it?

The common sense of the mass of mankind will answer these questions without a moment's hesitation. Healthy humanity, finding itself hard pressed to escape from real sin and degradation, will leave the brooding over speculative pollution to the cynics and the "righteous overmuch" who, disagreeing in everything else, unite in blind insensibility to the nobleness of the visible world, and in inability to appreciate the grandeur of the place Man occupies therein.

Nay, more; thoughtful men, once escaped from the blinding influences of traditional prejudice, will find in the lowly stock whence Man has sprung the best evidence of the splendour of his capacities; and will discern in his long progress through the Past a reasonable ground of faith in his attainment of a nobler Future.

They will remember that, in comparing civilised man with the animal world, one is as the Alpine traveller

who sees the mountains soaring into the sky and can hardly discern where the deep shadowed crags and roseate peaks end, and where the clouds of heaven begin. Surely the awe-struck voyager may be excused if at first he refuses to believe the geologist, who tells him that these glorious masses are, after all, the hardened mud of primeval seas, or the cooled slag of subterranean furnaces—of one substance with the dullest clay, but raised by inward forces to that place of proud and seemingly inaccessible glory.

But the geologist is right; and due reflection on his teachings, instead of diminishing our reverence and our wonder, adds all the force of intellectual sublimity to the mere æsthetic intuition of the uninstructed beholder.

And after passion and prejudice have died away the same result will attend the teachings of the naturalist respecting that great Alps and Andes of the living world—Man. Our reverence for the nobility of manhood will not be lessened by the knowledge that Man is, in substance and in structure, one with the brutes; for he alone possesses the marvellous endowment of intelligible and rational speech, whereby, in the secular period of his existence, he has slowly accumulated and organised the experience which is almost wholly lost with the cessation of every individual life in other animals; so that now he stands raised upon it as on a mountain top, far above the level of his humble fellows, and transfigured from his grosser nature by reflecting here and there a ray from the infinite source of truth.

A SUCCINCT HISTORY OF THE CONTRO-  
VERSY RESPECTING THE CEREBRAL  
STRUCTURE OF MAN AND THE APES.

UP to the year 1857 all anatomists of authority, who had occupied themselves with the cerebral structure of the Apes—Cuvier, Tiedemann, Sandifort, Vrolik, Isidore G. St. Hilaire, Schroeder van der Kolk, Gratiolet—were agreed that the brain of the Apes possesses a POSTERIOR LOBE.

Tiedemann, in 1825, figured and acknowledged, in the text of his *Icones*, the existence of the POSTERIOR CORNU of the lateral ventricle in the Apes, not only under the title of "Scrobiculus parvus loco cornu posterioris"—a fact which has been paraded—but as "Cornu posterioris" (*Icones*, p. 54), a circumstance which has been as sedulously kept in the background.

Cuvier (*Leçons*, T. iii., p. 103) says: "The anterior or lateral ventricles possess a digital cavity [posterior cornu] only in Man and the Apes.....Its presence depends on that of the posterior lobes."

Schroeder van der Kolk and Vrolik and Gratiolet had also figured and described the posterior cornu in various Apes. As to the HIPPOCAMPUS MINOR, Tiedemann had erroneously asserted its absence in the Apes; but Schroeder van der Kolk and Vrolik had pointed out the existence of what they considered a rudimentary one in the Chimpanzee, and Gratiolet had expressly affirmed its existence in these animals. Such was the state of our information on these subjects in the year 1856.

In the year 1857, however, Professor Owen, either in ignorance of these well-known facts or else unjustifiably suppressing them, submitted to the Linnæan Society a paper "On the Characters, Principles of Division, and Primary Groups of the Class Mammalia," which was printed in the Society's Journal, and contains the following passage:—"In Man the brain presents an ascensive step in development, higher and more strongly marked than that by which the preceding sub-class was distinguished from the one below it. Not only do the cerebral hemispheres overlap the olfactory lobes and cerebellum, but they extend in advance of the one and further back than the other. The posterior development is so marked that anatomists have assigned to that

part the character of a third lobe; it is peculiar to the genus *Homo*, and equally peculiar is the posterior horn of the lateral ventricle and the 'hippocampus minor,' which characterise the hind lobe of each hemisphere."—*Journal of the Proceedings of the Linnæan Society*, vol. ii., p. 19.

As the essay in which this passage stands had no less ambitious an aim than the remodelling of the classification of the Mammalia, its author might be supposed to have written under a sense of peculiar responsibility, and to have tested with especial care the statements he ventured to promulgate. And even if this be expecting too much, hastiness, or want of opportunity for due deliberation, cannot now be pleaded in extenuation of any shortcomings; for the propositions cited were repeated two years afterwards in the Reade Lecture, delivered before so grave a body as the University of Cambridge, in 1859.

When the assertions, which I have italicised in the above extract, first came under my notice, I was not a little astonished at so flat a contradiction of the doctrines current among well-informed anatomists; but, not unnaturally imagining that the deliberate statements of a responsible person must have some foundation in fact, I deemed it my duty to investigate the subject anew before the time at which it would be my business to lecture thereupon came round. The result of my inquiries was to prove that Mr. Owen's three assertions, that "the third lobe, the posterior horn of the lateral ventricle, and the hippocampus minor" are "peculiar to the genus *Homo*," are contrary to the plainest facts. I communicated this conclusion to the students of my class; and then, having no desire to embark in a controversy which could not redound to the honour of British science, whatever its issue, I turned to more congenial occupations.

The time speedily arrived, however, when a persistence in this reticence would have involved me in an unworthy paltering with truth.

At the meeting of the British Association at Oxford, in 1860, Professor Owen repeated these assertions in my presence, and, of course, I immediately gave them a direct and unqualified contradiction, pledging myself to justify that unusual procedure elsewhere. I redeemed that pledge by publishing, in the January number of the *Natural History Review*



for 1861, an article wherein the truth of the three following propositions was fully demonstrated (*l. c.*, p. 71):—

"1. That the third lobe is neither peculiar to, nor characteristic of, Man, seeing that it exists in all the higher quadrumana."

"2. That the posterior cornu of the lateral ventricle is neither peculiar to, nor characteristic of, Man, inasmuch as it also exists in the higher quadrumana."

"3. That the *hippocampus minor* is neither peculiar to, nor characteristic of, Man, as it is found in certain of the higher quadrumana."

Furthermore, this paper contains the following paragraph (p. 76):—

"And lastly, Schroeder van der Kolk and Vrolik (*op. cit.*, p. 271), though they particularly note that 'the lateral ventricle is distinguished from that of Man by the very defective proportions of the posterior cornu, wherein only a stripe is visible as an indication of the hippocampus minor;' yet the Figure 4, in their second Plate, shows that this posterior cornu is a perfectly distinct and unmistakable structure, quite as large as it often is in Man. It is the more remarkable that Professor Owen should have overlooked the explicit statement and figure of these authors, as it is quite obvious, on comparison of the figures, that his woodcut of the brain of a Chimpanzee (*l. c.*, p. 19) is a reduced copy of the second figure of Messrs. Schroeder van der Kolk and Vrolik's first Plate.

"As M. Gratiolet (*l. c.*, p. 18), however, is careful to remark, 'unfortunately, the brain which they have taken as a model was greatly altered (profondément affaîssé), whence the general form of the brain is given in these plates in a manner which is altogether incorrect.' Indeed, it is perfectly obvious, from a comparison of a section of the skull of the Chimpanzee with these figures, that such is the case; and it is greatly to be regretted that so inadequate a figure should have been taken as a typical representation of the Chimpanzee's brain."

From this time forth, the untenability of his position might have been as apparent to Professor Owen as it was to everyone else; but, so far from retracting the grave errors into which he had fallen, Professor Owen has persisted in and reiterated them; first in a lecture delivered before the Royal Institution on the 19th of March, 1861, which is

admitted to have been accurately reproduced in the *Athenæum* for the 23rd of the same month, in a letter addressed by Professor Owen to that journal on the 30th of March. The *Athenæum* report was accompanied by a diagram purporting to represent a Gorilla's brain, but in reality so extraordinary a misrepresentation that Professor Owen substantially, though not explicitly, withdraws it in the letter in question. In amending this error, however, Professor Owen fell into another of much graver import, as his communication concludes with the following paragraph: "For the true proportion in which the cerebrum covers the cerebellum in the highest Apes, reference should be made to the figure of the undissected brain of the Chimpanzee in my *Reade's Lecture on the Classification, &c., of the Mammalia*, p. 25, fig. 7, 8vo, 1859."

It would not be credible, if it were not unfortunately true, that this figure, to which the trusting public is referred, without a word of qualification, "for the true proportion in which the cerebrum covers the cerebellum in the highest Apes," is exactly that unacknowledged copy of Schroeder van der Kolk and Vrolik's figure whose utter inaccuracy had been pointed out years before by Gratiolet, and had been brought to Professor Owen's knowledge by myself in the passage of my article in the *Natural History Review* above quoted.

I drew public attention to this circumstance again in my reply to Professor Owen, published in the *Athenæum* for April 13th, 1861; but the exploded figure was reproduced once more by Professor Owen, without the slightest allusion to its inaccuracy, in the *Annals of Natural History* for June, 1861!

This proved too much for the patience of the original authors of the figure, Messrs. Schroeder van der Kolk and Vrolik, who, in a note addressed to the Academy of Amsterdam, of which they were members, declared themselves to be, though decided opponents of all forms of the doctrine of progressive development, above all things lovers of truth; and that, therefore, at whatever risk of seeming to lend support to views which they disliked, they felt it their duty to take the first opportunity of publicly repudiating Professor Owen's misuse of their authority.

In this note they frankly admitted the justice of the criticisms of M. Gratiolet,

quoted above, and they illustrated, by new and careful figures, the posterior lobe, the posterior cornu, and the hippocampus minor of the Orang. Furthermore, having demonstrated the parts, at one of the sittings of the Academy, they add, "la présence des parties contestées y a été universellement reconnue par les anatomistes présents à la séance. Le seul doute qui soit resté se rapporte aux pes *Hippocampi minor*.....A l'état frais l'indice du petit pied d'Hippocampe était plus prononcé que maintenant."

Professor Owen repeated his erroneous assertions at the meeting of the British Association in 1861, and again—without any obvious necessity, and without adducing a single new fact or new argument, or being able in any way to meet the crushing evidence from original dissections of numerous Apes' brains, which had in the meanwhile been brought forward by Professor Rolleston,<sup>1</sup> F.R.S., Mr. Marshall,<sup>2</sup> F.R.S., Mr. Flower,<sup>3</sup> Mr. Turner,<sup>4</sup> and myself<sup>5</sup>—revived the subject at the Cambridge meeting of the same body in 1862. Not content with the tolerably vigorous repudiation which these unprecedented proceedings met with in Section D, Professor Owen sanctioned the publication of a version of his own statements, accompanied by a strange misrepresentation of mine (as may be seen by comparison of the *Times* report of the discussion), in the *Medical Times* for October 11th, 1862. I subjoin the conclusion of my reply in the same journal for October 25th:—

"If this were a question of opinion, or a question of interpretation of parts or of terms; were it even a question of observation, in which the testimony of my own senses alone was pitted against that of another person, I should adopt a very

different tone in discussing this matter. I should, in all humility, admit the likelihood of having myself erred in judgment, failed in knowledge, or been blinded by prejudice.

"But no one pretends now that the controversy is one of terms or of opinions. Novel and devoid of authority as some of Professor Owen's proposed definitions may have been, they might be accepted without changing the great features of the case. Hence, though special investigations into these matters have been undertaken during the last two years by Dr. Allen Thomson, by Dr. Rolleston, by Mr. Marshall, and by Mr. Flower, all, as you are aware, anatomists of repute in this country, and by Professors Schroeder van der Kolk and Vrolik (whom Professor Owen incautiously tried to press into his own service) on the Continent, all these able and conscientious observers have with one accord testified to the accuracy of my statements, and to the utter baselessness of the assertions of Professor Owen. Even the venerable Rudolph Wagner, whom no man will accuse of progressionist proclivities, has raised his voice on the same side; while not a single anatomist, great or small, has supported Professor Owen.

"Now, I do not mean to suggest that scientific differences should be settled by universal suffrage, but I do conceive that solid proofs must be met by something more than empty and unsupported assertions. Yet, during the two years through which this preposterous controversy has dragged its weary length, Professor Owen has not ventured to bring forward a single preparation in support of his often-repeated assertions.

"The case stands thus, therefore: Not only are the statements made by me in consonance with the doctrines of the best older authorities, and with those of all recent investigators, but I am quite ready to demonstrate them on the first monkey that comes to hand; while Professor Owen's assertions are not only in diametrical opposition to both old and new authorities, but he has not produced, and, I will add, cannot produce, a single preparation which justifies them."

I now leave this subject, for the present. For the credit of my calling I should be glad to be, hereafter, for ever silent upon it. But, unfortunately, this is a matter upon which, after all that has occurred, no mistake or confusion of terms is

<sup>1</sup> "On the Affinities of the Brain of the Orang." *Natural History Review*, April, 1861.

<sup>2</sup> "On the Brain of a Young Chimpanzee." *Ibid.*, July, 1861.

<sup>3</sup> "On the Posterior Lobes of the Cerebrum of the Quadrumana." *Philosophical Transactions*, 1862.

<sup>4</sup> "On the Anatomical Relations of the Surfaces of the Tentorium to the Cerebrum and Cerebellum in Man and the Lower Mammals." *Proceedings of the Royal Society of Edinburgh*, March, 1862.

<sup>5</sup> "On the Brain of Ateles." *Proceedings of Zoological Society*, 1861.



possible; and in affirming that the posterior lobe, the posterior cornu, and the hippocampus minor exist in certain Apes, I am stating either that which is true or that which I must know to be false. The

question has thus become one of personal veracity. For myself, I will accept no other issue than this, grave as it is, to the present controversy.

### III.

#### ON SOME FOSSIL REMAINS OF MAN

I HAVE endeavoured to show, in the preceding Essay, that the ANTHROPINI, or Man Family, form a very well-defined group of the Primates, between which and the immediately following Family, the CATARRHINI, there is, in the existing world, the same entire absence of any transitional form or connecting link as between the CATARRHINI and PLATYRHINI.

It is a commonly received doctrine, however, that the structural intervals between the various existing modifications of organic beings may be diminished, or even obliterated, if we take into account the long and varied succession of animals and plants which have preceded those now living, and which are known to us only by their fossilised remains. How far this doctrine is well based, how far, on the other hand, as our knowledge at present stands, it is an over-statement of the real facts of the case, and an exaggeration of the conclusions fairly deducible from them, are points of grave importance, but into the discussion of which I do not at present propose to enter. It is enough that such a view of the relations of extinct to living beings has been propounded

to lead us to inquire, with anxiety, how far the recent discoveries of human remains in a fossil state bear out, or oppose, that view.

I shall confine myself, in discussing this question, to those fragmentary Human skulls from the caves of Engis in the valley of the Meuse, in Belgium, and of the Neanderthal, near Düsseldorf, the geological relations of which have been examined with so much care by Sir Charles Lyell, upon whose high authority I shall take it for granted that the Engis skull belonged to a contemporary of the Mammoth (*Elephas primigenius*) and of the woolly Rhinoceros (*Rhinoceros tichorhinus*), with the bones of which it was found associated; and that the Neanderthal skull is of great, though uncertain, antiquity. Whatever be the geological age of the latter skull, I conceive it is quite safe (on the ordinary principles of paleontological reasoning) to assume that the former takes us to, at least, the further side of the vague biological limit which separates the present geological epoch from that which immediately preceded it. And there can be no doubt that the physical geography of Europe has

changed wonderfully since the bones of Men and Mammoths, Hyænas and Rhinoceroses, were washed pell-mell into the cave of Engis.

The skull from the cave of Engis was originally discovered by Professor Schmerling, and was described by him, together with other human remains disinterred at the same time, in his valuable work, *Recherches sur les ossements fossiles découverts dans les cavernes de la Province de Liège*, published in 1833 (p. 59, *et seq.*), from which the following paragraphs are extracted, the precise expressions of the author being, as far as possible, preserved:—

In the first place, I must remark that these human remains, which are in my possession, are characterised, like the thousands of bones which I have lately been disinterring, by the extent of the decomposition which they have undergone, which is precisely the same as that of the extinct species; all, with a few exceptions, are broken; some few are rounded, as is frequently found to be the case in fossil remains of other species. The fractures are vertical or oblique; none of them are eroded; their colour does not differ from that of other fossil bones, and varies from whitish yellow to blackish. All are lighter than recent bones, with the exception of those which have a calcareous incrustation, and the cavities of which are filled with such matter.

The cranium which I have caused to be figured, Plate I., figs. 1, 2, is that of an old person. The sutures are beginning to be effaced: all the facial bones are wanting, and of the temporal bones only a fragment of that of the right side is preserved.

The face and the base of the cranium had been detached before the skull was deposited in the cave, for we were unable to find those parts, though the whole cavern was regularly searched. The cranium was met with at a depth of a metre and a half [five feet nearly] hidden under an osseous breccia, composed of the remains of small animals, and containing one rhinoceros tusk, with several teeth of horses and of ruminants. This breccia,

which has been spoken of above (p. 30), was a metre [ $3\frac{1}{4}$  feet about] wide, and rose to the height of a metre and a half about the floor of the cavern, to the walls of which it adhered strongly.

The earth which contained this human skull exhibited no trace of disturbance: teeth of rhinoceros, horse, hyæna, and bear, surrounded it on all sides.

The famous Blumenbach<sup>1</sup> has directed attention to the differences presented by the form and the dimensions of human crania of different races. This important work would have assisted us greatly if the face, a part essential for the determination of race with more or less accuracy, had not been wanting in our fossil cranium.

We are convinced that, even if the skull had been complete, it would not have been possible to pronounce, with certainty, upon a single specimen, for individual variations are so numerous in the crania of one and the same race that one cannot, without laying oneself open to large chances of error, draw any inference from a single fragment of a cranium to the general form of the head to which it belonged.

Nevertheless, in order to neglect no point respecting the form of this fossil skull, we may observe that, from the first, the elongated and narrow form of the forehead attracted our attention.

In fact, the slight elevation of the frontal, its narrowness, and the form of the orbit, approximate it more nearly to the cranium of an Ethiopian than to that of an European; the elongated form and the produced occiput are also characters which we believe to be observable in our fossil cranium; but to remove all doubt upon that subject I have caused the contours of the cranium of an European and of an Ethiopian to be drawn and the foreheads represented. Plate II., Figs. 1 and 2, and in the same Plate, Figs. 3 and 4, will render the differences easily distinguishable; and a single glance at the figures will be more instructive than a long and wearisome description.

At whatever conclusion we may arrive as to the origin of the man from whence this fossil skull proceeded, we may express an opinion without exposing ourselves to a fruitless controversy. Each may adopt the hypothesis which seems to

<sup>1</sup> *Decas Collectionis suæ craniorum diversarum gentium illustrata*.—Gottingæ, 1790-1820.



him most probable; for my own part, I hold it to be demonstrated that this cranium has belonged to a person of limited intellectual faculties, and we conclude thence that it belonged to a man of a low degree of civilisation—a deduction which is borne out by contrasting the capacity of the frontal with that of the occipital region.

Another cranium of a young individual was discovered in the floor of the cavern beside the tooth of an elephant; the skull was entire when found, but the moment it was lifted it fell into pieces,

Figure 4 is a fragment of a superior maxillary bone, the molar teeth of which are worn down to the roots.

I possess two vertebræ, a first and last dorsal.

A clavicle of the left side (see Plate III., Fig. 1); although it belonged to a young individual, this bone shows that he must have been of great stature.<sup>1</sup>

Two fragments of the radius, badly preserved, do not indicate that the height of the man to whom they belonged exceeded five feet and a half.

As to the remains of the upper extre-



FIG. 22.—The skull from the cave of Engis—viewed from the right side. One half the size of nature. *a* glabella, *b* occipital protuberance (*a* to *b* glabella-occipital line), *c* auditory foramen.

which I have not, as yet, been able to put together again. But I have represented the bones of the upper jaw, Plate I., Fig. 5. The state of the alveoli and the teeth shows that the molars had not yet pierced the gum. Detached milk molars and some fragments of a human skull proceed from this same place. The Figure 3 represents a human superior incisor tooth, the size of which is truly remarkable.<sup>1</sup>

<sup>1</sup> In a subsequent passage Schmerling remarks upon the occurrence of an incisor

mities, those which are in my possession consist merely of a fragment of an ulna and of a radius (Plate III., Figs. 5 and 6).

Figure 2, Plate IV., represents a metacarpal bone contained in the breccia,

tooth "of enormous size" from the caverns of Engihoul. The tooth figured is somewhat long, but its dimensions do not appear to me to be otherwise remarkable.

<sup>1</sup> The figure of this clavicle measures five inches from end to end in a straight line; so that the bone is rather a small than a large one.

of which we have spoken ; it was found in the lower part above the cranium ; add to this some metacarpal bones, found at very different distances, half-a-dozen metatarsals, three phalanges of the hand, and one of the foot.

This is a brief enumeration of the remains of human bones collected in the cavern of Engis, which has preserved for us the remains of three individuals surrounded by those of the Elephant, the Rhinoceros, and of Carnivora of species unknown in the present creation.

From the cave of Engihoul, opposite that of Engis, on the right bank of the Meuse, Schmerling obtained the remains of three other individuals of Man, among which were only two fragments of parietal bones, but many bones of the extremities. In one case a broken fragment of an ulna was soldered to a like fragment of a radius by stalagmite—a condition frequently observed among the bones of the Cave Bear (*Ursus spelæus*) found in the Belgian caverns.

It was in the cavern of Engis that Professor Schmerling found, incrustated with stalagmite and joined to a stone, the pointed bone implement which he has figured in Fig. 7 of his Plate XXXVI., and worked flints were found by him in all those Belgian caves which contained an abundance of fossil bones.

A short letter from M. Geoffroy St. Hilaire, published in the *Comptes Rendus* of the Academy of Sciences of Paris for July 2nd, 1838, speaks of a visit (and apparently a very hasty one) paid to the collection of Professor "Schermidt" (which is presumably a misprint for Schmerling) at Liège. The writer briefly criticises the drawings which illustrate Schmerling's work, and affirms that the "human cranium is a little longer than it is represented" in Schmerling's figure.

The only other remark worth quoting is this:—

The aspect of the human bones differs little from that of the cave bones with which we are familiar, and of which there is a considerable collection in the same place. With respect to their special forms, compared with those of the varieties of recent human crania, few *certain* conclusions can be put forward ; for much greater differences exist between the different specimens of well-characterised varieties than between the fossil cranium of Liège and that of one of those varieties selected as a term of comparison.

Geoffroy St. Hilaire's remarks are, it will be observed, little but an echo of the philosophic doubts of the describer and discoverer of the remains. As to the critique upon Schmerling's figures, I find that the side view given by the latter is really about  $\frac{3}{10}$ ths of an inch shorter than the original, and that the front view is diminished to about the same extent. Otherwise the representation is not in any way inaccurate, but corresponds very well with the cast which is in my possession.

A piece of the occipital bone, which Schmerling seems to have missed, has since been fitted on to the rest of the cranium by an accomplished anatomist, Dr. Spring, of Liège, under whose direction an excellent plaster cast was made for Sir Charles Lyell. It is upon and from a duplicate of that cast that my own observations and the accompanying figures, the outlines of which are copied from very accurate Camera lucida drawings by my friend Mr. Busk, reduced to one-half of the natural size, are made.

As Professor Schmerling observes, the base of the skull is destroyed, and the facial bones are entirely absent ; but the roof of the cranium, consisting of the frontal, parietal, and the greater



part of the occipital bones, as far as the middle of the occipital foramen, is entire, or nearly so. The left temporal

considerable portion of the squamous element of the temporal, are well preserved (Fig. 22).




FIG. 23.—The Engis skull viewed from above (*A*) and in front (*B*).

The lines of fracture which remain between the coadjusted pieces of the skull, and are faithfully displayed in Schmerling's figure, are readily traceable in the cast. The sutures are also discernible, but the complex disposition of their serrations shown in the figure is not obvious in the cast. Though the ridges which give attachment to muscles are not excessively prominent, they are well marked, and, taken together with the apparently well-developed frontal sinuses and the condition of the sutures, leave no doubt on my mind that the skull is that of an adult, if not middle-aged man.

The extreme length of the skull is 7.7 inches. Its extreme breadth, which corresponds very nearly with the interval between the parietal protuberances, is not more than 5.4 inches. The proportion of the length to the breadth is therefore very nearly as 100 to 70. If a line be drawn from the point at which the

bone is wanting. Of the right temporal, the parts in the immediate neighbourhood of the auditory foramen, the mastoid process, and a

brow curves in towards the root of the nose, and which is called the "glabella" (*a*, Fig. 22), to the occipital protuberance (*b*), and the distance to the

highest point of the arch of the skull be measured perpendicularly from this line, it will be found to be 4.75 inches. Viewed from above (Fig. 23, A), the forehead presents an evenly rounded curve, and passes into the contour of the sides and back of the skull, which describes a tolerably regular elliptical curve.

The front view (Fig. 23, B) shows that the roof of the skull was very regularly and elegantly arched in the transverse direction, and that the transverse diameter was a little less below the parietal protuberances than above them. The forehead cannot be called narrow in relation to the rest of the skull, nor can it be called a retreating forehead; on the contrary, the antero-posterior contour of the skull is well arched, so that the distance along that contour, from the nasal depression to the occipital protuberance, measures about 13.75 inches. The transverse arc of the skull, measured from one auditory foramen to the other, across the middle of the sagittal suture, is about 13 inches. The sagittal suture itself is 5.5 inches long.

The supraciliary prominences or brow-ridges (on each side of *a*, Fig. 22) are well, but not excessively, developed, and are separated by a median depression. Their principal elevation is disposed so obliquely that I judge them to be due to large frontal sinuses.

If a line joining the glabella and the occipital protuberance (*a*, *b*, Fig. 22) be made horizontal, no part of the occipital region projects more than  $\frac{1}{16}$ th of an inch behind the posterior extremity of that line, and the upper edge of the auditory fora-

men (*c*) is almost in contact with a line drawn parallel with this upon the outer surface of the skull.

A transverse line drawn from one auditory foramen to the other traverses, as usual, the forepart of the occipital foramen. The capacity of the interior of this fragmentary skull has not been ascertained.

The history of the human remains from the cavern in the Neanderthal may best be given in the words of their original describer, Dr. Schaaffhausen,<sup>1</sup> as translated by Mr. Busk:—

In the early part of the year 1857 a human skeleton was discovered in a limestone cave in the Neanderthal, near Hochdal, between Düsseldorf and Elberfeld. Of this, however, I was unable to procure more than a plaster cast of the cranium, taken at Elberfeld, from which I drew up an account of its remarkable conformation, which was, in the first instance, read on the 4th of February, 1857, at the meeting of the Lower Rhine Medical and Natural History Society, at Bonn.<sup>2</sup> Subsequently Dr. Fuhlrott, to whom science is indebted for the preservation of these bones, which were not at first regarded as human, and into whose possession they afterwards came, brought the cranium from Elberfeld to Bonn, and entrusted it to me for more accurate anatomical examination. At the General Meeting of the Natural History Society of Prussian Rhineland and Westphalia, at Bonn, on the 2nd of June, 1857,<sup>3</sup> Dr. Fuhlrott himself gave a full account of the locality, and of the circumstances under which the discovery was made. He was of opinion

<sup>1</sup> *On the Crania of the Most Ancient Races of Man*. By Professor D. Schaaffhausen, of Bonn. (From Müller's *Archiv*, 1858, p. 453.) With Remarks and Original Figures taken from a Cast of the Neanderthal Cranium. By George Busk, F.R.S., &c. *Natural History Review*, April, 1861.

<sup>2</sup> *Verhandl. d. Naturhist. Vereins der preuss. Rheinlande und Westphalens*, xiv. Bonn, 1857.

<sup>3</sup> *Ib.*, *Correspondenzblatt*. No. 2.



that the bones might be regarded as fossil; and, in coming to this conclusion, he laid especial stress upon the existence of dendritic deposits, with which their surface was covered, and which were first noticed upon them by Professor Mayer. To this communication I appended a brief report on the results of my anatomical examination of the bones. The conclusions at which I arrived were:—1st, That the extraordinary form of the skull was due to a natural conformation hitherto not known to exist, even in the most barbarous races. 2nd, That these remarkable human remains belonged to a period antecedent to the time of the Celts and Germans, and were in all probability derived from one of the wild races of North-Western Europe, spoken of by Latin writers, and which were encountered as autochthones by the German immigrants. And 3rdly, That it was beyond doubt that these human relics were traceable to a period at which the latest animals of the diluvium still existed; but that no proof of this assumption, nor, consequently, of their so-called *fossil* condition, was afforded by the circumstances under which the bones were discovered.

As Dr. Fuhlrott has not yet published his description of these circumstances, I borrow the following account of them from one of his letters:—"A small cave or grotto, high enough to admit a man, and about 15 feet deep from the entrance, which is 7 or 8 feet wide, exists in the southern wall of the gorge of the Neanderthal, as it is termed, at a distance of about 100 feet from the Düssel, and about 60 feet above the bottom of the valley. In its earlier and uninjured condition, this cavern opened upon a narrow plateau lying in front of it, and from which the rocky wall descended almost perpendicularly into the river. It could be reached, though with difficulty, from above. The uneven floor was covered to a thickness of 4 or 5 feet with a deposit of mud, sparingly intermixed with rounded fragments of chert. In the removing of this deposit the bones were discovered. The skull was first noticed, placed nearest to the entrance of the cavern; and, further in, the other bones, lying in the same horizontal plane. Of this I was assured, in the most positive terms, by two labourers who were employed to clear out the grotto, and who were questioned by me on the spot. At first no idea was entertained of

the bones being human; and it was not till several weeks after their discovery that they were recognised as such by me, and placed in security.

"But, as the importance of the discovery was not at the time perceived, the labourers were very careless in the collecting, and secured chiefly only the larger bones; and to this circumstance it may be attributed that fragments merely of the probably perfect skeleton came into my possession."

My anatomical examination of these bones afforded the following results:—

The cranium is of unusual size, and of a long elliptical form. A most remarkable peculiarity is at once obvious in the extraordinary development of the frontal sinuses, owing to which the supraciliary ridges, which coalesce completely in the middle, are rendered so prominent that the frontal bone exhibits a considerable hollow or depression above, or rather behind, them; whilst a deep depression is also formed in the situation of the root of the nose. The forehead is narrow and low, though the middle and hinder portions of the cranial arch are well developed. Unfortunately, the fragment of the skull that has been preserved consists only of the portion situated above the roof of the orbits and the superior occipital ridges, which are greatly developed, and almost conjoined so as to form a horizontal eminence. It includes almost the whole of the frontal bone, both parietals, a small part of the squamous, and the upper third of the occipital. The recently fractured surfaces show that the skull was broken at the time of its disinterment. The cavity holds 16,876 grains of water, whence its cubical contents may be estimated at 57.64 inches, or 1,033.24 cubic centimetres. In making this estimation, the water is supposed to stand on a level with the orbital plate of the frontal, with the deepest notch in the squamous margin of the parietal, and with the superior semi-circular ridges of the occipital. Estimated in dried millet-seed, the contents equalled 31 ounces, Prussian Apothecaries' weight. The semi-circular line indicating the upper boundary of the attachment of the temporal muscle, though not very strongly marked, ascends nevertheless to more than half the height of the parietal bone. On the right supraciliary ridge is observable an oblique furrow or depression, indicative

of an injury received during life.<sup>1</sup> The coronal and sagittal sutures are on the exterior nearly closed, and on the inside so completely ossified as to have left no traces whatever, whilst the lambdoidal remains quite open. The depressions for the Pacchionian glands are deep and numerous; and there is an unusually deep vascular groove immediately behind the coronal suture, which, as it terminates in a foramen, no doubt transmitted a *vena emissaria*. The course of the frontal suture is indicated externally by a slight ridge; and, where it joins the coronal, this ridge rises into a small protuberance. The course of the sagittal suture is grooved, and above the angle of the occipital bone the parietals are depressed.

mm.<sup>2</sup>

The length of the skull from the nasal process of the frontal over the vertex to the superior semi-circular lines of the occipital meas- sures .. .. .	303 (300) = 12.0"
Circumference over the orbital ridges and the superior semi-circular lines of the occipital .. .. .	590 (590) = 23.37" or 23"
Width of the frontal from the middle of the temporal line on one side to the same point on the opposite .. .. .	104 (114) = 4.1"—4.5"
Length of the frontal from the nasal process to the coronal suture .. .. .	133 (125) = 5.25"—5"
Extreme width of the frontal sinuses .. .. .	25 (23) = 1.0"—0.9"
Vertical height above a line joining the deepest notches in the squamous border of the parietals .. .. .	70 = 2.75"
Width of hinder part of skull from one parietal protuberance to the other .. .. .	138 (150) = 5.4"—5.9"
Distance from the upper angle of the occipital to the superior semi-circular lines .. .. .	51 (60) = 1.9"—2.4"
Thickness of the bone at the parietal protuberance .. .. .	8.
— at the angle of the occipital .. .. .	9.
— at the superior semi-circular line of the occipital .. .. .	10 = 0.3"

Besides the cranium, the following bones have been secured:—

1. Both thigh-bones, perfect. These,

<sup>1</sup> This, Mr. Busk has pointed out, is probably the notch for the frontal nerve.

<sup>2</sup> The numbers in brackets are those which I should assign to the different measures, as taken from the plaster cast. —G. B.

like the skull, and all the other bones, are characterised by their unusual thickness, and the great development of all the elevations and depressions for the attachment of muscles. In the Anatomical Museum at Bonn, under the designation of "Giant's bones," are some recent thigh-bones, with which, in thickness, the foregoing pretty nearly correspond, although they are shorter.

	Giant's bones. mm.	Fossil bones. mm.
Length .. .. .	542 = 21.4"	438 = 17.4"
Diameter of head of femur .. .. .	54 = 2.14"	53 = 2.0"
Diameter of lower articular end, from one condyle to the other .. .. .	89 = 3.5"	87 = 3.4"
Diameter of femur in the middle .. .. .	33 = 1.2"	30 = 1.1"

2. A perfect right humerus, whose size shows that it belongs to the thigh-bones.

	mm.
Length .. .. .	312 = 12.3"
Thickness in the middle .. .. .	26 = 1.0"
Diameter of head .. .. .	49 = 1.9"

Also a perfect right radius of corresponding dimensions, and the upper third of a right ulna corresponding to the humerus and radius.

3. A left humerus, of which the upper third is wanting, and which is so much slenderer than the right as apparently to belong to a distinct individual; a left ulna, which, though complete, is pathologically deformed, the coronoid process being so much enlarged by bony growth that flexure of the elbow beyond a right angle must have been impossible; the anterior fossa of the humerus for the reception of the coronoid process being also filled up with a similar bony growth. At the same time the olecranon is curved strongly downwards. As the bone presents no sign of rachitic degeneration, it may be supposed that an injury sustained during life was the cause of the ankylosis. When the left ulna is compared with the right radius, it might at first sight be concluded that the bones respectively belonged to different individuals, the ulna being more than half an inch too short for articulation with a corresponding radius. But it is clear that this shortening, as well as the attenuation of the left humerus, are both consequent upon the pathological condition above described.

4. A left ilium, almost perfect, and



belonging to the femur ; a fragment of the right scapula ; the anterior extremity of a rib of the right side ; and the same part of a rib of the left side ; the hinder part of a rib of the right side ; and, lastly, two hinder portions and one middle portion of ribs, which, from their unusually rounded shape and abrupt curvature, more resemble the ribs of a carnivorous animal than those of a man. Dr. H. v. Meyer, however, to whose judgment I defer, will not venture to declare them to be ribs of any animal ; and it only remains to suppose that this abnormal condition has arisen from an unusually powerful development of the thoracic muscles.

The bones adhere strongly to the tongue, although, as proved by the use of hydrochloric acid, the greater part of the cartilage is still retained in them, which appears, however, to have undergone that transformation into gelatine which has been observed by v. Bibra in fossil bones. The surface of all the bones is in many spots covered with minute black specks, which, more especially under a lens, are seen to be formed of very delicate *dendrites*. These deposits, which were first observed on the bones by Dr. Meyer, are most distinct on the inner surface of the cranial bones. They consist of a ferruginous compound, and from their black colour may be supposed to contain manganese. Similar dendritic formations also occur, not unfrequently, on laminated rocks, and are usually found in minute fissures and cracks. At the meeting of the Lower Rhine Society at Bonn on the 1st April, 1857, Prof. Meyer stated that he had noticed in the museum of Poppelsdorf similar dendritic crystallisations on several fossil bones of animals, and particularly on those of *Ursus spelæus*, but still more abundantly and beautifully displayed on the fossil bones and teeth of *Equus adamiticus*, *Elephas primigenius*, &c., from the caves of Bolve and Sundwig. Faint indications of similar *dendrites* were visible in a Roman skull from Siegburg ; whilst other ancient skulls, which had lain for centuries in the earth, presented no trace of them.<sup>1</sup> I am indebted to H. v. Meyer for the following remarks on this subject :—

“The incipient formation of dendritic deposits, which were formerly regarded as a sign of a truly fossil condition, is

interesting. It has even been supposed that in diluvial deposits the presence of *dendrites* might be regarded as affording a certain mark of distinction between bones mixed with the diluvium at a somewhat later period and the true diluvial relics, to which alone it was supposed that these deposits were confined. But I have long been convinced that neither can the absence of *dendrites* be regarded as indicative of recent age, nor their presence as sufficient to establish the great antiquity of the objects upon which they occur. I have myself noticed upon paper, which could scarcely be more than a year old, dendritic deposits, which could not be distinguished from those on fossil bones. Thus I possess a dog's skull from the Roman colony of the neighbouring Heddersheim, *Castrum Hadrianum*, which is in no way distinguishable from the fossil bones from the Frankish caves ; it presents the same colour, and adheres to the tongue just as they do ; so that this character also, which, at a former meeting of German naturalists at Bonn, gave rise to amusing scenes between Buckland and Schmerling, is no longer of any value. In disputed cases, therefore, the condition of the bone can scarcely afford the means for determining with certainty whether it be fossil—that is to say, whether it belonged to geological antiquity or to the historical period.”

As we cannot now look upon the primitive world as representing a wholly different condition of things, from which no transition exists to the organic life of the present time, the designation of *fossil*, as applied to a *bone*, has no longer the sense it conveyed in the time of Cuvier. Sufficient grounds exist for the assumption that man co-existed with the animals found in the diluvium ; and many a barbarous race may, before all historical time, have disappeared, together with the animals of the ancient world, whilst the races whose organisation is improved have continued the genus. The bones which form the subject of this paper present characters which, although not decisive as regards a geological epoch, are, nevertheless, such as indicate a very high antiquity. It may also be remarked that, common as is the occurrence of diluvial animal bones in the muddy deposits of caverns, such remains have not hitherto been met with in the caves of the Neanderthal ; and that the bones, which were covered by a deposit of mud not more than

<sup>1</sup> *Verh. des Naturhist. Vereins in Bonn*, xiv. 1857.

four or five feet thick, and without any protective covering of stalagmite, have retained the greatest part of their organic substance.

These circumstances might be adduced against the probability of a geological antiquity. Nor should we be justified in regarding the cranial conformation as perhaps representing the most savage primitive type of the human race; since crania exist among living savages, which, though not exhibiting such a remarkable conformation of the forehead, which gives the skull somewhat the aspect of that of the large apes, still in other respects—as, for instance, in the greater depth of the temporal fossæ, the crest-like, prominent temporal ridges, and a generally less capacious cranial cavity—exhibit an equally low stage of development. There is no reason for supposing that the deep frontal hollow is due to any artificial flattening, such as is practised in various modes by barbarous nations in the Old and New World. The skull is quite symmetrical, and shows no indication of counter-pressure at the occiput; whilst, according to Morton, in the Flat-heads of the Columbia, the frontal and parietal bones are always unsymmetrical. Its conformation exhibits the sparing development of the anterior part of the head which has been so often observed in very ancient crania, and affords one of the most striking proofs of the influence of culture and civilisation on the form of the human skull.

In a subsequent passage Dr. Schaaffhausen remarks:—

There is no reason whatever for regarding the unusual development of the frontal sinuses in the remarkable skull from the Neanderthal as an individual or pathological deformity; it is unquestionably a typical race-character, and is physiologically connected with the uncommon thickness of the other bones of the skeleton, which exceeds by about one-half the usual proportions. This expansion of the frontal sinuses, which are appendages of the air-passages, also indicates an unusual force and power of endurance in the movements of the body, as may be concluded from the size of all the ridges and processes for the attachment of the muscles or bones. That this conclusion may be drawn from the existence of large frontal sinuses, and a prominence of the lower frontal region, is confirmed in many ways by other observa-

tions. By the same characters, according to Pallas, the wild horse is distinguished from the domesticated, and, according to Cuvier, the fossil cave-bear from every recent species of bear; whilst, according to Roulin, the pig, which has become wild in America, and regained a resemblance to the wild boar, is thus distinguished from the same animal in the domesticated state, as is the chamois from the goat; and, lastly, the bull-dog, which is characterised by its large bones and strongly-developed muscles from every other kind of dog. The estimation of the facial angle, the determination of which, according to Professor Owen, is also difficult in the great apes, owing to the very prominent supra-orbital ridges, in the present case is rendered still more difficult from the absence both of the auditory opening and of the nasal spine. But if the proper horizontal position of the skull be taken from the remaining portions of the orbital plates, and the ascending line made to touch the surface of the frontal bone behind the prominent supra-orbital ridges, the facial angle is not found to exceed  $56^{\circ}$ .<sup>1</sup> Unfortunately, no portions of the facial bones, whose conformation is so decisive as regards the form and expression of the head, have been preserved. The cranial capacity, compared with the uncommon strength of the corporeal frame, would seem to indicate a small cerebral development. The skull, as it is, holds about 31 ounces of millet-seed; and as, from the proportionate size of the wanting bones, the whole cranial cavity should have about 6 ounces more added, the contents, were it perfect, may be taken at 37 ounces. Tiedemann assigns, as the cranial contents in the Negro, 40, 38, and 35 ounces. The cranium holds rather more than 36 ounces of water, which corresponds to a capacity of 1,033.24 cubic centimetres. Huschke estimates the cranial contents of a Negress at 1,127 cubic centimetres, of an old Negro at 1,146 cubic centimetres. The capacity of the Malay skulls, estimated by water, equalled 36, 33 ounces, whilst in the diminutive Hindoos it falls to as little as 27 ounces.

After comparing the Neanderthal cranium with many others, ancient

<sup>1</sup> Estimating the facial angle in the way suggested, on the cast I should place it at  $64^{\circ}$  to  $67^{\circ}$ .—G. B.



and modern, Professor Schaaffhausen concludes thus :—

But the human bones and cranium from the Neanderthal exceed all the rest in those peculiarities of conformation which lead to the conclusion of their belonging to a barbarous and savage race. Whether the cavern in which they were found, unaccompanied with any trace of human art, were the place of their interment, or whether, like the bones of extinct animals elsewhere, they had been washed into it, they may still be regarded as the most ancient memorial of the early inhabitants of Europe.

Mr. Busk, the translator of Dr. Schaaffhausen's paper, has enabled us to form a very vivid conception of the degraded character of the Neanderthal skull, by placing side by side with its outline that of the skull of a Chimpanzee, drawn to the same absolute size.

Some time after the publication of the translation of Professor Schaaffhausen's Memoir, I was led to study the cast of the Neanderthal cranium with more attention than I had previously bestowed upon it, in consequence of wishing to supply Sir Charles Lyell with a diagram exhibiting the special peculiarities of this skull as compared with other human skulls. In order to do this it was necessary to identify with precision those points in the skulls compared which corresponded anatomically. Of these points, the glabella was obvious enough ; but when I had distinguished another, defined by the occipital protuberance and superior semi-circular line, and had placed the outline of the Neanderthal skull against that of the Engis skull, in such a position that the glabella and occipital protuberance of both were

intersected by the same straight line, the difference was so vast and the flattening of the Neanderthal skull so prodigious (compare Figs. 22 and 24, A) that I at first imagined I must have fallen into some error. And I was the more inclined to suspect this, as, in ordinary human skulls, the occipital protuberance and superior semi-circular curved line on the exterior of the occiput correspond pretty closely with the "lateral sinuses" and the line of attachment of the tentorium internally. But on the tentorium rests, as I have said in the preceding Essay, the posterior lobe of the brain ; and, hence, the occipital protuberance, and the curved line in question, indicate, approximately, the lower limits of that lobe. Was it possible for a human being to have the brain thus flattened and depressed ; or, on the other hand, had the muscular ridges shifted their position ? In order to solve these doubts, and to decide the question whether the great supraciliary projections did or did not arise from the development of the frontal sinuses, I requested Sir Charles Lyell to be so good as to obtain for me from Dr. Fuhlrott, the possessor of the skull, answers to certain queries, and, if possible, a cast, or at any rate drawings or photographs, of the interior of the skull.

Dr. Fuhlrott replied, with a courtesy and readiness for which I am infinitely indebted to him, to my inquiries, and, furthermore, sent three excellent photographs. One of these gives a side view of the skull, and from it Fig. 24, A has been shaded. The second (Fig. 25, A) exhibits the wide openings of




FIG. 24.—The skull from the Neanderthal cavern. A, side, B, front, and C, top view. One-half the natural size. The outlines from camera lucida drawings, one-half the natural size, by Mr. Busk; the details from the cast and from Dr. Fuhlrott's photographs. *a* glabella; *b* occipital protuberance; *d* lambdoidal suture.



the frontal sinuses upon the inferior surface of the frontal part of the skull, into which, Dr. Fuhlrott writes, "a probe may be introduced to the depth of an inch," and demonstrates the great extension of the thickened supraciliary ridges beyond the cerebral cavity. The third, lastly (Fig. 25, B), exhibits the edge and the interior of the posterior, or occipital, part of the skull, and shows very

In truth, the Neanderthal cranium has most extraordinary characters. It has an extreme length of 8 inches, while its breadth is only 5.75 inches, or, in other words, its length is to its breadth as 100 : 72. It is exceedingly depressed, measuring only about 3.4 inches from the glabello-occipital line to the vertex. The longitudinal arc, measured in the same way as in the Engis skull, is 12 inches; the trans-

verse arc cannot be exactly ascertained, in consequence of the absence of the temporal bones, but was probably about the same, and certainly exceeded  $10\frac{1}{4}$  inches. The horizontal circumference is 23 inches. But this great circumference arises largely from the vast development of the supraciliary ridges, though the perimeter of the brain case itself is not small. The large supraciliary ridges give the forehead a far more retreating appearance than its internal contour would bear out.

To an anatomical eye the posterior part of the

skull is even more striking than the anterior. The occipital protuberance occupies the extreme posterior end of the skull when the glabello-occipital line is made horizontal, and, so far from any part of the occipital region extending beyond it, this region of the skull slopes obliquely upward and forward, so that the lambdoidal suture is situated




FIG. 25.—Drawings from Dr. Fuhlrott's photographs of parts of the interior of the Neanderthal cranium. A, view of the under and inner surface of the frontal region, showing the inferior apertures of the frontal sinuses (*a*). B, corresponding view of the occipital region of the skull, showing the impressions of the lateral sinuses (*aa*).

clearly the two depressions for the lateral sinuses, sweeping inwards towards the middle line of the roof of the skull, to form the longitudinal sinus. It was clear, therefore, that I had not erred in my interpretation, and that the posterior lobe of the brain of the Neanderthal man must have been as much flattened as I suspected it to be.

well upon the upper surface of the cranium. At the same time, notwithstanding the great length of the skull, the sagittal suture is remarkably short ( $4\frac{1}{2}$  inches), and the squamosal suture is very straight.

In reply to my questions Dr. Fuhlrott writes that the occipital bone "is in a state of perfect preservation as far as the upper semi-circular line, which is a very strong ridge, linear at its extremities, but enlarging towards the middle, where it forms two ridges (*bourrelets*), united by a linear continuation, which is slightly depressed in the middle."

"Below the left ridge the bone exhibits an obliquely inclined surface, six lines (French) long, and twelve lines wide."

This last must be the surface, the contour of which is shown in Fig. 24, A, below *b*. It is particularly interesting, as it suggests that, notwithstanding the flattened condition of the occiput, the posterior cerebral lobes must have projected considerably beyond the cerebellum, and as it constitutes one among several points of similarity between the Neanderthal cranium and certain Australian skulls.

Such are the two best known forms of human cranium which have been found in what may be fairly termed a fossil state. Can either be shown to fill up or diminish, to any appreciable extent, the structural interval which exists between Man and the man-like Apes? Or, on the other hand, does neither depart more widely from the average structure of the human cranium than normally-formed skulls of men are known to do at the present day?

It is impossible to form any opinion on these questions without some preliminary acquaintance with the range of variation exhibited by human structure in general—a subject which has been but imperfectly studied, while, even of what is known, my limits will necessarily allow me to give only a very imperfect sketch.

The student of anatomy is perfectly well aware that there is not a single organ of the human body the structure of which does not vary, to a greater or less extent, in different individuals. The skeleton varies in the proportions, and even to a certain extent in the connections, of its constituent bones. The muscles which move the bones vary largely in their attachments. The varieties in the mode of distribution of the arteries are carefully classified, on account of the practical importance of a knowledge of their shiftings to the surgeon. The characters of the brain vary immensely, nothing being less constant than the form and size of the cerebral hemispheres and the richness of the convolutions upon their surface, while the most changeable structures of all in the human brain are exactly those on which the unwise attempt has been made to base the distinctive characters of humanity—viz., the posterior cornu of the lateral ventricle, the hippocampus minor, and the degree of projection of the posterior lobe beyond the cerebellum. Finally, as all the world knows, the hair and skin of human beings may present the most extraordinary diversities in colour and in texture.

So far as our present knowledge goes, the majority of the structural varieties to which allusion is here



made are individual. The ape-like arrangement of certain muscles which is occasionally met with<sup>\*</sup> in the white races of mankind is not known to be more common among Negroes or Australians; nor, because the brain of the Hottentot Venus was found to be smoother, to have its convolutions more symmetrically disposed, and to be, so far, more ape-like than that of ordinary Europeans, are we justified in concluding a like condition of the brain to prevail universally among the lower races of mankind, however probable that conclusion may be.

We are, in fact, sadly wanting in information respecting the disposition of the soft and destructible organs of every race of mankind but our own; and even of the skeleton, our Museums are lamentably deficient in every part but the cranium. Skulls enough there are, and since the time when Blumenbach and Camper first called attention to the marked and singular differences which they exhibit, skull-collecting and skull-measuring has been a zealously pursued branch of Natural History, and the results obtained have been arranged and classified by various writers, among whom the late active and able Retzius must always be the first named.

Human skulls have been found to differ from one another, not merely in their absolute size and in the absolute capacity of the brain case, but in the proportions which the diameters of the latter bear to one another; in the relative size of the bones of the face (and more particularly of the jaws and teeth) as compared with those of the

skull; in the degree to which the upper jaw (which is of course followed by the lower) is thrown backwards and downwards under the fore-part of the brain case, or forwards and upwards in front of and beyond it. They differ further in the relations of the transverse diameter of the face, taken through the cheek bones, to the transverse diameter of the skull; in the more rounded or more gable-like form of the roof of the skull, and in the degree to which the hinder part of the skull is flattened or projects beyond the ridge, into and below which the muscles of the neck are inserted.

In some skulls the brain case may be said to be *round*, the extreme length not exceeding the extreme breadth by a greater proportion than 100 to 80, while the difference may be much less.<sup>\*</sup> Men possessing such skulls were termed by Retzius *brachycephalic*, and the skull of a Calmuck, of which a front and side view (reduced outline copies of which are given in Figure 26) are depicted by Von Baer in his excellent *Crania Selecta*, affords a very admirable example of that kind of skull. Other skulls, such as that of a Negro copied in Fig. 27 from Mr. Busk's *Crania Typica*, have a very different, greatly elongated form, and may be termed *oblong*. In this skull the extreme length is to the extreme breadth as 100 to not more than 67, and the transverse diameter of the human skull may fall below even this proportion. People having such skulls were called by Retzius *dolichocephalic*.

The most cursory glance at the side

<sup>\*</sup> See an excellent Essay by Mr. Church on the Myology of the Orang, in the *Natural History Review* for 1861.

<sup>\*</sup> In no normal human skull does the breadth of the brain case exceed its length.

views of these two skulls will suffice to prove that they differ, in another respect, to a very striking extent. The profile of the face of the Calmuck is almost vertical, the facial bones being thrown downwards and under the fore part of the skull. The profile of the face of the Negro, on the other hand, is singularly inclined, the front part of the jaws projecting far forward beyond the level of the fore part of the skull. In the former case the skull is said to be *orthognathous*, or straight-jawed; in the latter, it is called *prognathous*, a term which has been rendered with more force than elegance by the Saxon equivalent—"snouty."

Various methods have been devised in order to express with some accuracy the degree of prognathism or orthognathism of any given skull, most of these methods being essentially modifications of that devised by Peter Camper in order to attain what he called the "facial angle."

But a little consideration will show that any "facial angle" that has been devised can be competent to express the structural modifications involved in prognathism and orthognathism only in a rough and general sort of way. For the lines, the intersection of which forms the facial angle, are drawn through points of the skull, the position of each of which is modified by a number of circumstances, so that the

angle obtained is a complex resultant of all these circumstances, and is not the expression of any one definite organic relation of the parts of the skull.




FIG. 26.—Side and front views of the round and orthognathous skull of a Calmuck, after Von Baer. One-third the natural size.

I have arrived at the conviction that no comparison of crania is worth very much that is not founded upon the establishment of a relatively fixed base



line, to which the measurements, in all cases, must be referred. Nor do I think it is a very difficult matter to decide what that base line should be.

roof; it is converted into cartilage earlier and more completely than the sides and roof; and the cartilaginous base ossifies and becomes soldered into one piece long before the roof. I conceive, then, that the base of the skull may be demonstrated developmentally to be its relatively fixed part, the roof and sides being relatively movable.

The same truth is exemplified by the study of the modifications which the skull undergoes in ascending from the lower animals up to man.

In such a mammal as a Beaver (Fig. 28), a line (*a b*) drawn through the bones termed basi-occipital, basisphenoid, and presphenoid, is very long in proportion to the extreme length of the cavity which contains the cerebral hemispheres (*g h*). The plane of the occipital foramen (*b c*) forms a slightly acute angle with this "basicranial axis," while the plane of the tentorium (*i T*) is inclined at rather more than  $90^\circ$  to the "basicranial axis"; and so is the plane of the perforated plate (*a d*), by which




FIG. 27.—Ob'long and prognathous skull of a Negro; side and front views. One-third of the natural size.

The parts of the skull, like those of the rest of the animal framework, are developed in succession; the base of the skull is formed before its sides and

the filaments of the olfactory nerve leave the skull. Again, a line drawn through the axis of the face, between the bones called ethmoid and vomer—

the "basifacial axis" ( $fe$ )—forms an exceedingly obtuse angle where, when produced, it cuts the "basicranial axis."

If the angle made by the line  $b\ c$  with  $a\ b$  be called the "occipital angle," and the angle made by the line  $a\ d$  with  $a\ b$  be termed the "olfactory angle," and that made by  $i\ T$  with  $a\ b$  the "tentorial angle," then all these, in the mammal in question, are nearly right angles, varying between  $80^\circ$  and  $110^\circ$ . The angle  $e\ f\ b$ , or that made by the cranial with the facial axis, and which may be termed the "crano-facial angle," is extremely obtuse, amounting, in the case of the Beaver, to at least  $150^\circ$ .

But if a series of sections of mammalian skulls, intermediate between a Rodent and a Man (Fig. 28), be examined, it will be found that in the higher crania the basi-cranial axis becomes shorter relatively to the cerebral length; that the "olfactory angle" and the "occipital angle" become more obtuse; and that the "crano-facial angle" becomes more acute by the bending down, as it were, of the facial axis upon the cranial axis. At the same time, the roof of the cranium becomes more and more arched, to allow of the increasing height of the cerebral hemispheres, which is eminently characteristic of man, as well as of that backward extension beyond the cerebellum, which reaches its maximum in the South American Monkeys. So that, at last, in the human skull (Fig. 29), the cerebral length is between twice and thrice as great as the length of the basicranial axis; the olfactory plane is  $20^\circ$  or  $30^\circ$  on the *under* side of that axis; the occipital angle, instead of being less

than  $90^\circ$ , is as much as  $150^\circ$  or  $160^\circ$ ; the crano-facial angle may be  $90^\circ$  or less, and the vertical height of the skull may have a large proportion to its length.

It will be obvious, from an inspection of the diagrams, that the basi-cranial axis is, in the ascending series of Mammalia, a relatively fixed line, on which the bones of the sides and roof of the cranial cavity and of the face may be said to revolve downwards and forwards or backwards, according to their position. The arc described by any one bone or plane, however, is not by any means always in proportion to the arc described by another.

Now comes the important question: Can we discern, between the lowest and the highest forms of the human cranium, anything answering, in however slight a degree, to this revolution of the side and roof bones of the skull upon the basicranial axis observed upon so great a scale in the mammalian series? Numerous observations lead me to believe that we must answer this question in the affirmative.

The diagrams in Figure 29 are reduced from very carefully-made diagrams of sections of four skulls, two round and orthognathous, two long and prognathous, taken longitudinally and vertically through the middle. The sectional diagrams have then been superimposed in such a manner that the basal axes of the skulls coincide by their anterior ends, and in their direction. The deviations of the rest of the contours (which represent the interior of the skulls only) show the differences of the skulls from one another when these axes are regarded as relatively fixed lines.

The dark contours are those of an



Australian and of a Negro skull; the light contours are those of a Tartar skull, in the Museum of the Royal College of Surgeons; and of a well-developed round skull from a cemetery

the same way as, though to a far less degree than, the skulls of the lower mammals differ from those of Man. Furthermore, the plane of the occipital foramen ( $b c$ ) forms a somewhat




FIG. 28.—Longitudinal and vertical sections of the skulls of a Beaver (*Castor Canadensis*), a Lemur (*L. Catta*), and a Baboon (*Cynocephalus Papio*).  $ab$ , the basicranial axis;  $bc$ , the occipital plane;  $i T$ , the tentorial plane;  $ad$ , the olfactory plane;  $fe$ , the basifacial axis;  $cba$ , occipital angle;  $Tia$ , tentorial angle;  $dab$ , olfactory angle;  $efb$ , cranio-facial angle;  $gh$ , extreme length of the cavity which lodges the cerebral hemispheres, or "cerebral length." The length of the basicranial axis as to this length, or, in other words, the proportional length of the line  $gh$  to that of  $ab$  taken as 100, in the three skulls, is as follows:—Beaver, 70 to 100; Lemur, 119 to 100; Baboon, 144 to 100. In an adult male Gorilla the cerebral length is as 170 to the basicranial axis taken as 100, in the Negro (Fig. 29) as 236 to 100. In the Constantinople skull (Fig. 29) it is as 266 to 100. The difference between the highest Ape's skull and the lowest Man's is therefore very strikingly brought out by these measurements.

In the diagram of the Baboon's skull the dotted lines  $d^1 d^2$ , &c., give the angles of the Lemur's and Beaver's skull, as laid down upon the basicranial axis of the Baboon. The line  $ab$  has the same length in each diagram.

in Constantinople, of uncertain race, in my own possession.

It appears at once from these views that the prognathous skulls, so far as their jaws are concerned, do really differ from the orthognathous in much

smaller angle with the axis in these particular prognathous skulls than in the orthognathous; and the like may be slightly true of the perforated plate of the ethmoid, though this point is not so clear. But it is singular to

remark that, in another respect, the prognathous skulls are less ape-like than the orthognathous, the cerebral cavity projecting decidedly more beyond the anterior end of the axis in the prognathous than in the orthognathous skulls.

It will be observed that these diagrams reveal an immense range of variation in the capacity and relative proportion to the cranial axis of the different regions of the cavity which contains the brain in the different skulls. Nor is the difference in the extent to which the cerebral overlaps the cerebellar cavity less singular. A round skull (Fig. 29, *Const.*) may have a greater posterior cerebral projection than a long one (Fig. 29, *Negro*).

Until human crania have been largely worked out in a manner similar to that here suggested—until it shall be an opprobrium to an ethnological collection to possess a single skull which is not bisected longitudinally—until the angles and measurements here mentioned, together with a number of others of which I cannot speak in this place, are determined and tabulated, with reference to the basicranial axis as unity, for large numbers of skulls of the different races of Mankind, I do not think we shall have any very safe basis for that ethnological craniology which aspires to give the anatomical characters of the crania of the different races of mankind.

At present I believe that the general outlines of what may be safely said upon that subject may be summed up in a very few words. Draw a line on a globe from the Gold Coast in Western Africa to the steppes of Tartary. At the southern and western




FIG. 29.—Sections of orthognathous (light contour) and prognathous (dark contour) skulls, one-third of the natural size. *a b*, basicranial axis; *b c, b' c'*, plane of the occipital foramen; *d' d'*, hinder end of the palatine bone; *e' e'*, front end of the upper jaw; *T T'*, insertion of the tentorium.

end of that line there live the most dolichocephalic, prognathous, curly-haired, dark-skinned of men—the true Negroes. At the northern and eastern end of the same line there live the most brachycephalic, orthognathous, straight-haired, yellow-skinned of



men—the Tartars and Calmucks. The two ends of this imaginary line are indeed, so to speak, ethnological antipodes. A line drawn at right angles, or nearly so, to this polar line through Europe and Southern Asia to Hindostan, would give us a sort of equator, around which round-headed, oval-headed, and oblong-headed, prognathous and orthognathous, fair and dark races—but none possessing the excessively marked characters of Calmuck or Negro—group themselves.

It is worthy of notice that the regions of the antipodal races are antipodal in climate, the greatest contrast the world affords, perhaps, being that between the damp, hot, steaming, alluvial coast plains of the West Coast of Africa and the arid, elevated steppes and plateaux of Central Asia, bitterly cold in winter, and as far from the sea as any part of the world can be.

From Central Asia eastward to the Pacific Islands and subcontinents on the one hand, and to America on the other, brachycephaly and orthognathism gradually diminish, and are replaced by dolichocephaly and prognathism, less, however, on the American Continent (throughout the whole length of which a rounded type of skull prevails largely, but not exclusively)<sup>1</sup> than in the Pacific region, where, at length, on the Australian Continent and in the adjacent islands, the oblong skull, the projecting jaws, and the dark skin reappear, with so much departure, in other respects, from the

Negro type that ethnologists assign to these people the special title of “Negrito-toes.”

The Australian skull is remarkable for its narrowness and for the thickness of its walls, especially in the region of the supraciliary ridge, which is frequently, though not by any means invariably, solid throughout, the frontal sinuses remaining undeveloped. The nasal depression, again, is extremely sudden, so that the brows overhang, and give the countenance a particularly lowering, threatening expression. The occipital region of the skull, also, not unfrequently becomes less prominent; so that it not only fails to project beyond a line drawn perpendicular to the hinder extremity of the glabella-occipital line, but even, in some cases, begins to shelve away from it, forwards, almost immediately. In consequence of this circumstance, the parts of the occipital bone which lie above and below the tuberosity make a much more acute angle with one another than is usual, whereby the hinder part of the base of the skull appears obliquely truncated. Many Australian skulls have a considerable height, quite equal to that of the average of any other race; but there are others in which the cranial roof becomes remarkably depressed, the skull, at the same time, elongating so much that, probably, its capacity is not diminished. The majority of skulls possessing these characters, which I have seen, are from the neighbourhood of Port Adelaide, in South Australia, and have been used by the natives as water vessels; to which end the face has been knocked away, and a string passed through

<sup>1</sup> See Dr. D. Wilson's valuable paper “On the supposed prevalence of one Cranial Type throughout the American aborigines.”—*Canadian Journal*, vol. ii., 1857.

the vacuity and the occipital foramen, so that the skull was suspended by the greater part of its basis.

Figure 30 represents the contour of a skull of this kind from Western Port, with the jaw attached, and of the Neanderthal skull, both reduced to one-third of the size of nature. A small additional amount of flattening and lengthening, with a corresponding increase of the supraciliary ridge, would convert the Australian brain case into a form identical with that of the aberrant fossil.

And now to return to the fossil skulls, and to the rank which they occupy among, or beyond, these existing varieties of cranial conformation. In the first place, I must remark that, as Professor Schmerling well observed (*supra*, p. 81) in commenting upon the Engis skull, the formation of a safe judgment upon the question is greatly hindered by the absence of the jaws from both the crania, so that there is no means of deciding with certainty whether they were more or less prognathous than the lower existing races of mankind. And yet, as we have seen, it is more in this respect than any other that human skulls vary, towards and from, the brutal type—the brain case of an average dolichocephalic European differing far less from that of a Negro, for example, than his jaws do. In the absence of the jaws, then, any judgment on the relations of the fossil skulls to recent races must be accepted with a certain reservation.

But taking the evidence as it

stands, and turning first to the Engis skull, I confess I can find no character in the remains of that cranium which, if it were a recent skull, would give any trustworthy clue as to the race to which it might appertain. Its contours and measurements agree very well with those of some Australian skulls which I have examined, and especially has it a tendency towards that occipital flattening, to the great extent of which, in some




FIG. 30.—An Australian skull from Western Port, in the Museum of the Royal College of Surgeons, with the contour of the Neanderthal skull. Both reduced to one-third the natural size.

Australian skulls, I have alluded. But all Australian skulls do not present this flattening, and the supraciliary ridge of the Engis skull is quite unlike that of the typical Australians.

On the other hand, its measurements agree equally well with those of some European skulls. And assuredly there is no mark of degradation about any part of its structure. It is, in fact, a fair average human skull, which might have belonged to a philosopher, or might



have contained the thoughtless brains of a savage.

The case of the Neanderthal skull is very different. Under whatever aspect we view this cranium, whether we regard its vertical depression, the

Schaaffhausen states (*supra*, p. 89) that the cranium, in its present condition, holds 1,033.24 cubic centimetres of water, or about 63 cubic inches; and as the entire skull could hardly have held less than an addi-

tional 12 cubic inches, its capacity may be estimated at about 75 cubic inches, which is the average capacity given by Morton for Polynesian and Hottentot skulls.

So large a mass of brain as this would alone suggest that the pithecoïd tendencies indicated by this skull did not extend deep into the organisation; and this conclusion is borne out by the dimensions of the other bones of the skeleton given by Professor Schaaffhausen, which show that the absolute height and relative proportions of the limbs were quite those of an European of middle stature. The bones are, indeed, stouter, but this and the great development of the muscular ridges noted by Dr. Schaaffhausen are characters to be expected in savages. The Patagonians, exposed without shelter or protection to a climate possibly not very dissimilar from that of Europe at the time during

which the Neanderthal man lived, are remarkable for the stoutness of their limb bones.

In no sense, then, can the Neanderthal bones be regarded as the remains of a human being intermediate between




FIG. 31.—Ancient Danish skull from a tumulus at Borreby; one-third of the natural size. From a camera lucida drawing by Mr. Busk.

enormous thickness of its supraciliary ridges, its sloping occiput, or its long and straight squamosal suture, we meet with ape-like characters, stamping it as the most pithecoïd of human crania yet discovered. But Professor

Men and Apes. At most they demonstrate the existence of a man whose skull may be said to revert somewhat towards the pithecoïd type—just as a Carrier, or a Pouter, or a Tumbler may sometimes put on the plumage of its primitive stock, the *Columba livia*.

And, indeed, though truly the most pithecoïd of known human skulls, the Neanderthal cranium is by no means so isolated as it appears to be at first, but forms, in reality, the extreme term of a series leading gradually from it to the highest and best developed of human crania. On the one hand, it is closely approached by the flattened Australian skulls, of which I have spoken, from which other Australian forms lead us gradually up to skulls having very much the type of the Engis cranium. And, on the other hand, it is even more closely affined to the skulls of certain ancient people who inhabited Denmark during the "stone period," and were probably either contemporaneous with, or later than, the makers of the "refuse heaps," or "Kjokkenmøddings" of that country.

The correspondence between the longitudinal contour of the Neanderthal skull and that of some of those skulls from the tumuli at Borreby, very accurate drawings of which have been made by Mr. Busk, is very close. The occiput is quite as retreating, the supraciliary ridges are nearly as prominent, and the skull is as low. Furthermore, the Borreby skull resembles the Neanderthal form more closely than any of the Australian skulls do, by the much more rapid retrocession of the forehead. On the

other hand, the Borreby skulls are all somewhat broader, in proportion to their length, than the Neanderthal skull, while some attain that proportion of breadth to length (80 : 100) which constitutes brachycephaly.

In conclusion, I may say that the fossil remains of Man hitherto discovered do not seem to me to take us appreciably nearer to that lower pithecoïd form by the modification of which he has, probably, become what he is. And, considering what is now known of the most ancient races of men; seeing that they fashioned flint axes and flint knives and bone skewers, of much the same pattern as those fabricated by the lowest savages at the present day, and that we have every reason to believe the habits and modes of living of such people to have remained the same from the time of the Mammoth and the tichorhine Rhinoceros till now, I do not know that this result is other than might be expected.

Where, then, must we look for primæval Man? Was the oldest *Homo sapiens* pliocene or miocene, or yet more ancient? In still older strata, do the fossilised bones of an Ape more anthropoid, or a Man more pithecoïd, than any yet known await the researches of some unborn palæontologist?

Time will show. But, in the meanwhile, if any form of the doctrine of progressive development is correct, we must extend by long epochs the most liberal estimate that has yet been made of the antiquity of Man.



## IV.

## ON THE METHODS AND RESULTS OF ETHNOLOGY

[1865]

ETHNOLOGY is the science which determines the distinctive characters of the persistent modifications of mankind ; which ascertains the distribution of those modifications in present and past times, and seeks to discover the causes, or conditions of existence, both of the modifications and of their distribution. I say "persistent" modifications, because, unless incidentally, ethnology has nothing to do with chance and transitory peculiarities of human structure. And I speak of "persistent modifications" or "stocks" rather than of "varieties," or "races," or "species," because each of these last well-known terms implies, on the part of its employer, a preconceived opinion touching one of those problems, the solution of which is the ultimate object of the science ; and in regard to which, therefore, ethnologists are especially bound to keep their minds open and their judgments freely balanced.

Ethnology, as thus defined, is a branch of ANTHROPOLOGY, the great science which unravels the complexities of human structure ; traces out the relations of man to other animals ; studies all that is especially human in the mode in which man's complex functions are performed ; and searches after the conditions which have determined his presence in the world. And anthropology is a section of ZOOLOGY, which again is the animal half of

BIOLOGY—the science of life and living things.

Such is the position of ethnology, such are the objects of the ethnologist. The paths or methods, by following which he may hope to reach his goal, are diverse. He may work at man from the point of view of the pure zoologist, and investigate the anatomical and physiological peculiarities of Negroes, Australians, or Mongolians, just as he would inquire into those of pointers, terriers, and turnspits—"persistent modifications" of man's almost universal companion. Or he may seek aid from researches into the most human manifestation of humanity—Language ; and assuming that what is true of speech is true of the speaker—a hypothesis as questionable in science as it is in ordinary life—he may apply to mankind themselves the conclusions drawn from a searching analysis of their words and grammatical forms.

Or, the ethnologist may turn to the study of the practical life of men ; and, relying upon the inherent conservatism and small inventiveness of untutored mankind, he may hope to discover in manners and customs, or in weapons, dwellings, and other handiwork, a clue to the origin of the resemblances and differences of nations. Or, he may resort to that kind of evidence which is yielded by History proper, and consists of the beliefs of men concerning

past events, embodied in traditional, or in written, testimony. Or, when that thread breaks, Archæology, which is the interpretation of the unrecorded remains of man's works, belonging to the epoch since the world has reached its present condition, may still guide him. And, when even the dim light of archæology fades, there yet remains Palæontology, which, in these latter years, has brought to daylight once more the exuvia of ancient populations, whose world was not our world, who have been buried in river beds immemorially dry, or carried by the rush of waters into caves, inaccessible to inundation since the dawn of tradition.

Along each, or all, of these paths the ethnologist may press towards his goal; but they are not equally straight, or sure, or easy to tread. The way of palæontology has but just been laid open to us. Archæological and historical investigations are of great value for all those peoples whose ancient state has differed widely from their present condition, and who have the good or evil fortune to possess a history. But, on taking a broad survey of the world, it is astonishing how few nations present either condition. Respecting five-sixths of the persistent modifications of mankind, history and archæology are absolutely silent. For half the rest, they might as well be silent for anything that is to be made of their testimony. And, finally, when the question arises as to what was the condition of mankind more than a paltry two or three thousand years ago, history and archæology are, for the most part, mere dumb dogs. What light does either of these branches of knowledge throw

on the past of the men of the New World, if we except the Central Americans and the Peruvians; on that of the Africans, save those of the Valley of the Nile and a fringe of the Mediterranean; on that of all the Polynesian, Australian, and central Asiatic peoples, the former of whom probably, and the last certainly, were, at the dawn of history, substantially what they are now? While thankfully accepting what history has to give him, therefore, the ethnologist must not look for too much from her.

Is more to be expected from inquiries into the customs and handicrafts of man? It is to be feared not. In reasoning from identity of custom to identity of stock the difficulty always obtrudes itself that, the minds of men being everywhere similar, differing in quality and quantity but not in kind of faculty, like circumstances must tend to produce like contrivances; at any rate, so long as the need to be met and conquered is of a very simple kind. That two nations use calabashes or shells for drinking-vessels, or that they employ spears, or clubs, or swords and axes of stone and metal as weapons and implements, cannot be regarded as evidence that these two nations had a common origin, or even that intercommunication ever took place between them; seeing that the convenience of using calabashes or shells for such purposes, and the advantage of poking an enemy with a sharp stick, or hitting him with a heavy one, must be early forced by nature upon the mind of even the stupidest savage. And when he had found out the use of a stick, he would need no prompting to discover the value of



a chipped or whetted stone, or of an angular piece of native metal, for the same object. On the other hand, it may be doubted whether the chances are not greatly against independent peoples arriving at the manufacture of a boomerang or of a bow; which last, if one comes to think of it, is a rather complicated apparatus; and the tracing of the distribution of inventions as complex as these, and of such strange customs as betel-chewing and tobacco-smoking, may afford valuable ethnological hints.

Since the time of Leibnitz, and guided by such men as Humboldt, Abel Remusat, and Klaproth, Philology has taken far higher ground. Thus Prichard affirms that "the history of nations, termed Ethnology, must be mainly founded on the relations of their languages."

An eminent living philologist, August Schleicher, in a recent essay, puts forward the claims of his science still more forcibly:—

If, however, language is the human *κατ' ἐξοχήν*, the suggestion arises whether it should not form the basis of any scientific systematic arrangement of mankind; whether the foundation of the natural classification of the genus *Homo* has not been discovered in it.

How little constant are cranial peculiarities and other so-called race characters! Language, on the other hand, is always a perfectly constant diagnostic. A German may occasionally compete in hair and prognathism with a negro, but a negro language will never be his mother tongue. Of how little importance for mankind the so-called race characters are is shown by the fact that speakers of languages belonging to one and the same linguistic family may exhibit the peculiarities of various races. Thus the settled Osmanli Turk exhibits Caucasian characters, whilst other so-called Tartaric Turks exemplify the Mongol type. On the other hand, the Magyar and the Basque do not depart in

any essential physical peculiarity from the Indo-Germans; whilst the Magyar, Basque, and Indo-Germanic tongues are widely different. Apart from their inconstancy, again, the so-called race characters can hardly yield a scientifically natural system. Languages, on the other hand, readily fall into a natural arrangement, like that of which other vital products are susceptible, especially when viewed from their morphological side.....The externally visible structure of the cerebral and facial skeletons, and of the body generally, is less important than that no less material but infinitely more delicate corporeal structure, the function of which is speech. I conceive, therefore, that the natural classification of languages is also the natural classification of mankind. With language, moreover, all the higher manifestations of man's vital activity are closely interwoven, so that these receive due recognition in and by that of speech.<sup>1</sup>

Without the least desire to depreciate the value of philology as an adjuvant to ethnology, I must venture to doubt, with Rudolphi, Desmoulins, Crawfurd, and others, its title to the leading position claimed for it by the writers whom I have just quoted. On the contrary, it seems to me obvious that though, in the absence of any evidence to the contrary, unity of languages may afford a certain presumption in favour of the unity of stock of the peoples speaking those languages, it cannot be held to prove that unity of stock, unless philologists are prepared to demonstrate that no nation can lose its language and acquire that of a distinct nation without a change of blood corresponding with the change of language. Desmoulins long ago put this argument exceedingly well:—

Let us imagine the recurrence of one of

<sup>1</sup> August Schleicher, *Ueber die Bedeutung der Sprache für die Naturgeschichte des Menschen*, pp. 16-18. Weimar, 1858.

those slow, or sudden, political revolutions, or say of those secular changes which, among different people and at different epochs, have annihilated historical monuments and even extinguished tradition. In that case the evidence, now so clear, that the negroes of Hayti were slaves imported by a French colony, who, by the very effect of the subordination involved in slavery, lost their own diverse languages and adopted that of their masters, would vanish. And metaphysical philosophers, observing the identity of Haytian French with that spoken on the shores of the Seine and the Loire, would argue that the men of St. Domingo with woolly heads, black and oily skins, small calves, and slightly bent knees, are of the same race, descended from the same parental stock, as the Frenchmen with silky brown, chestnut, or fair hair, and white skins. For, they would say, their languages are more similar than French is to German or Spanish.<sup>1</sup>

It must not be imagined that the case put by Desmoulins is a merely hypothetical one. Events precisely similar to the transport of a body of Africans to the West India Islands, indeed, cannot have happened among uncivilised races, but similar results have followed the importation of bodies of conquerors among an enslaved people over and over again. There is hardly a country in Europe in which two or more nations speaking widely different tongues have not become intermixed; and there is hardly a language of Europe of which we have any right to think that its structure affords a just indication of the amount of that intermixture.

As Dr. Latham has well said:—

It is certain that the language of England is of Anglo-Saxon origin, and that the remains of the original Keltic are unimportant. It is by no means so certain that the blood of Englishmen is equally Germanic. A vast amount of

Kelticism, not found in our tongue, very probably exists in our pedigrees. The ethnology of France is still more complicated. Many writers make the Parisian a Roman on the strength of his language; whilst others make him a Kelt on the strength of certain moral characteristics, combined with the previous Kelticism of the original Gauls. Spanish and Portuguese, as languages, are derivations from the Latin; Spain and Portugal, as countries, are Iberic, Latin, Gothic, and Arab, in different proportions. Italian is modern Latin all the world over; yet surely there must be much Keltic blood in Lombardy, and much Etruscan intermixture in Tuscany.

In the ninth century every man between the Elbe and the Niemen spoke some Slavonic dialect; they now nearly all speak German. Surely the blood is less exclusively Gothic than the speech.<sup>2</sup>

In other words, what philologer, if he had nothing but the vocabulary and grammar of the French and English languages to guide him, would dream of the real causes of the unlikeness of a Norman to a Provençal, of an Orcadian to a Cornishman? How readily might he be led to suppose that the different climatal conditions to which these speakers of one tongue have so long been exposed have caused their physical differences; and how little would he suspect that these are due (as we happen to know they are) to wide differences of blood.

Few take duly into account the evidence which exists as to the ease with which unlettered savages gain or lose a language. Captain Erskine, in his interesting *Journal of a Cruise Among the Islands of the Western Pacific*, especially remarks upon the "avidity with which the inhabitants of the polyglot islands of Melanesia, from New Caledonia to the Solomon

<sup>1</sup> Desmoulins, *Histoire Naturelle des Races Humaines*, p. 345, 1826.

<sup>2</sup> Latham, *Man and His Migrations*, p. 171.



Islands, adopt the improvements of a more perfect language than their own, which different causes and accidental communication still continue to bring to them"; and he adds that "among the Melanesian islands scarcely one was found by us which did not possess, in some cases still imperfectly, the decimal system of numeration in addition to their own, in which they reckon only to five."

Yet how much philological reasoning in favour of the affinity or diversity of two distinct peoples has been based on the mere comparison of numerals!

But the most instructive example of the fallacy which may attach to merely philological reasonings is that afforded by the Feejeans, who are, physically, so intimately connected with the adjacent Negritos of New Caledonia, &c., that no one can doubt to what stock they belong, and who yet, in the form and substance of their language, are Polynesian. The case is as remarkable as if the Canary Islands should have been found to be inhabited by negroes speaking Arabic, or some other clearly Semitic dialect, as their mother tongue. As it happens, the physical peculiarities of the Feejeans are so striking, and the conditions under which they live are so similar to those of the Polynesians, that no one has ventured to suggest that they are merely modified Polynesians—a suggestion which could otherwise certainly have been made. But if languages may be thus transferred from one stock to another without any corresponding intermixture of blood, what ethnological value has philology?—what security does unity of language afford

us that the speakers of that language may not have sprung from two, or three, or a dozen distinct sources?

Thus we come, at last, to the purely zoological method, from which it is not unnatural to expect more than from any other, seeing that, after all, the problems of ethnology are simply those which are presented to the zoologist by every widely distributed animal he studies. The father of modern zoology seems to have had no doubt upon this point. At the twenty-eighth page of the standard twelfth edition of the *Systema Naturæ*, in fact, we find:—

#### I. PRIMATES.

*Dentes primores incisores: superiores IV. paralleli mammæ pectorales II.*

- |                      |   |
|----------------------|---|
| 1. HOMO.             | Nosce te ipsum.   |
| Sapiens.             | 1. H. diurnus: <i>varians cultura, loco.</i>  |
| <i>Ferus.</i>        | Tetrapus, mutus, hirsutus.  |
| <i>Americanus</i> α. | Rufus, cholericus, rectus— <i>Pilis</i> nigris, rectis, crassis— <i>Naribus</i> patulis— <i>Facie</i> ephelítica: <i>Mento</i> subimberbi. <i>Pertinax</i> , contentus, liber. <i>Pingit</i> se lineis dædaleis rubris. |
| <i>Europæus</i> β.   | <i>Regitur</i> Consuetudine. Albus sanguineus torosus. <i>Pilis</i> flavescentibus, prolixis. <i>Oculis</i> cœruleis. <i>Levis</i> , argutus, inventor. <i>Tegitur</i> Vestimentis arctis. <i>Regitur</i> Ritibus.      |
| <i>Asiaticus</i> γ.  | Luridus, melancholicus, rigidus. <i>Pilis</i> nigricantibus. <i>Oculis</i> fuscis. <i>Severus</i> , fastuosus, avarus. <i>Tegitur</i> Indumentis laxis.   |
| <i>Afer</i> δ.       | <i>Regitur</i> Opinionibus. Niger, phlegmaticus, laxus. <i>Pilis</i> atris, contortuplicatis. <i>Cute</i> holosericea. <i>Naso</i> simo. <i>Labiis</i> tumidis. <i>Feminis</i> sinus pudoris.                           |

- Mammæ lactantes prolixæ.*  
*Vafer, segnis, negligens.*  
*Ungit se pingui. Regitur Arbitrio.*
- Monstrosus* ε. Solo(a) et arte (b c) variat. :  
 a. *Alpini* parvi, agiles, timidi. *Patagonici* magni, segnes.  
 b. *Monorchides* ut minus fertiles : Hottentotti. *Junceæ* puellæ, abdomine attenuato: Europææ.  
 c. *Macrocephali* capiti conico : Chinenses. *Plagiocephali* capite antice compresso: Canadenses.

Turn a few pages further on in the same volume, and there appears, with a fine impartiality in the distribution of capitals and sub-divisional headings :—

### III. FERÆ.

*Dentes primores superiores sex, acutiusculi.*  
*Canini solitarii.*

12. CANIS. *Dentes primores superiores* VI. : laterales longiores distantes : intermedii lobati. Inferiores VI. : laterales lobati.  
*Laniarii* solitarii, incurvati.  
 Molares VI. s. VII. (pluresve quam in reliquis).
- familiaris* 1. C. cauda (sinistrorsum) recurvata.....
- domesticus* a. auriculis erectis, cauda subtus lanata.
- sagax* β. auriculis pendulis, digito spurio ad tibias posticas.
- grajus* γ. magnitudine lupi, trunco curvato, rostro attenuato, &c., &c.

Linnæus's definition of what he considers to be mere varieties of the species Man are, it will be observed, as completely free from any illusion to

linguistic peculiarities as those brief and pregnant sentences in which he sketches the characters of the varieties of the species Dog: "Pilis nigris, naribus patulis" may be set against "auriculis erectis, cauda subtus lanata"; while the remarks on the morals and manners of the human subjects seem as if they were thrown in merely by way of makeweight.

Buffon, Blumenbach (the founder of ethnology as a special science), Rudolphi, Bory de St. Vincent, Desmoulins, Cuvier, Retzius—indeed, I may say all the naturalists proper—have dealt with man from a no less completely zoological point of view; while, as might have been expected, those who have been least naturalists, and most linguists, have most neglected the zoological method, the neglect culminating in those who have been altogether devoid of acquaintance with anatomy.

Prichard's proposition, that language is more persistent than physical characters, is one which has never been proved, and indeed admits of no proof, seeing that the records of language do not extend so far as those of physical characters. But, until the superior tenacity of linguistic over physical peculiarities is shown, and until the abundant evidence which exists, that the language of a people may change without corresponding physical change in that people, is shown to be valueless, it is plain that the zoological court of appeal is the highest for the ethnologist, and that no evidence can be set against that derived from physical characters.

What, then, will a new survey of mankind from the Linnean point of view teach us?



The great antipodal block of land we call Australia has, speaking roughly, the form of a vast quadrangle, 2,000 miles on the side, and extends from the hottest tropical to the middle of the temperate zone. Setting aside the foreign colonists introduced within the last century, it is inhabited by people no less remarkable for the uniformity than for the singularity of their physical characters and social state. For the most part of fair stature, erect and well built, except for an unusual slenderness of the lower limbs, the AUSTRALIANS have dark, usually chocolate-coloured skins; fine dark wavy hair; dark eyes, overhung by beetle brows; coarse, projecting jaws; broad and dilated, but not especially flattened, noses and lips, which, though prominent, are eminently flexible.

The skulls of these people are always long and narrow, with a smaller development of the frontal sinuses than usually corresponds with such largely developed brow ridges. An Australian skull of a round form, or one the transverse diameter of which exceeds eight-tenths of its length, has never been seen. These people, in a word, are eminently "dolichocephalic," or long-headed; but, with this one limitation, their crania present considerable variations, some being comparatively high and arched, while others are more remarkably depressed than almost any other human skulls. The female pelvis differs comparatively little from the European; but in the pelves of male Australians which I have examined the antero-posterior and transverse diameters approach equality more nearly than is the case in Europeans.

No Australian tribe has ever been known to cultivate the ground, to use metals, pottery, or any kind of textile fabric. They rarely construct huts. Their means of navigation are limited to rafts or canoes, made of sheets of bark. Clothing, except skin cloaks for protection from cold, is a superfluity with which they dispense; and though they have some singular weapons, almost peculiar to themselves, they are wholly unacquainted with bows and arrows.

It is but a step, as it were, across Bass's Straits to Tasmania. Neither climate nor the characteristic forms of vegetable or animal life change largely on the south side of the Straits, but the early voyagers found Man singularly different from him on the north side. The skin of the Tasmanian was dark, though he lived between parallels of latitude corresponding with those of middle Europe in our own hemisphere; his jaws projected, his head was long and narrow; his civilisation was about on a footing with that of the Australian, if not lower, for I cannot discover that the Tasmanian understood the use of the throwing-stick. But he differed from the Australian in his woolly, negro-like hair; whence the name of NEGRITO, which has been applied to him and his congeners.

Such Negritos—differing more or less from the Tasmanian, but agreeing with him in dark skin and woolly hair—occupy New Caledonia, the New Hebrides, the Louisiade Archipelago; and, stretching to the Papuan Islands, and for a doubtful extent beyond them to the north and west, form a sort of belt, or zone, of Negrito population, interposed between the Australians on

the west and the inhabitants of the great majority of the Pacific islands on the east.

The cranial characters of the Negritos vary considerably more than those of their skin and hair, the most notable circumstance being the strong Australian aspect which distinguishes many Negrito skulls, while others tend rather towards forms common in the Polynesian islands.

In civilisation New Caledonia exhibits an advance upon Tasmania, and farther north there is a still greater improvement. But the bows and arrows, the perched houses, the outrigger canoes, the habits of betel-chewing and of kawa-drinking, which abound more or less among the northern Negritos, are probably to be regarded, not as the products of an indigenous civilisation, but merely as indications of the extent to which foreign influences have modified the primitive social state of these people.

From Tasmania or New Caledonia to New Zealand or Tongataboo is, again, but a brief voyage; but it brings about a still more notable change in the aspect of the indigenous population than that effected by the passage of Bass's Straits. Instead of being chocolate-coloured people, the Maories and Tongans are light brown; instead of woolly, they have straight, or wavy, black hair. And if from New Zealand we travel some 5,000 miles east to Easter Island; and from Easter Island for as great a distance north-west, to the Sandwich Islands; and thence 7,000 miles, westward and southward, to Sumatra; and even across the Indian Ocean, into the interior of Madagascar, we shall everywhere meet with

people whose hair is straight or wavy, and whose skins exhibit various shades of brown. These are the Polynesians, Micronesians, Indonesians, whom Latham has grouped together under the common title of AMPHINESIANS.

The cranial characters of these people, as of the Negritos, are less constant than those of their skin and hair. The Maori has a long skull; the Sandwich Islander a broad skull. Some, like these, have strong brow ridges; others, like the Dayaks and many Polynesians, have hardly any nasal indentation. It is only in the westernmost parts of their area that the Amphinesian nations know anything about bows and arrows as weapons, or are acquainted with the use of metals or with pottery. Everywhere they cultivate the ground, construct houses, and skilfully build and manage outrigger, or double, canoes; while almost everywhere they use some kind of fabric for clothing.

Between Easter Island, or the Sandwich Islands, and any part of the American coast is a much wider interval than that between Tasmania and New Zealand, but the ethnological interval between the American and the Polynesian is less than that between either of the previously named stocks.

The typical AMERICAN has straight, black hair and dark eyes, his skin exhibiting various shades of reddish or yellowish brown, sometimes inclining to olive. The face is broad and scantily bearded; the skull wide and high. Such people extend from Patagonia to Mexico, and much farther north along the west coast. In the main a race of hunters, they



had, nevertheless, at the time of the discovery of the Americas, attained a remarkable degree of civilisation in some localities. They had domesticated ruminants, and not only practised agriculture, but had learned the value of irrigation. They manufactured textile fabrics, were masters of the potter's art, and knew how to erect massive buildings of stone. They understood the working of the precious, though not of the useful, metals; and had even attained to a rude kind of hieroglyphic, or picture, writing. The Americans not only employ the bow and arrow, but, like some Amphinesians, the blow-pipe, as offensive weapons; but I am not aware that the outrigger canoe has ever been observed among them.

I have reason to suspect that some of the Fuegian tribes differ cranially from the typical Americans; and the Northern and Eastern American tribes have longer skulls than their Southern compatriots. But the *ESQUIMAUX*, who roam on the desolate and ice-bound coast of Arctic America, certainly present us with a new stock. The *Esquimaux* (among whom the Greenlanders are included), in fact, though they share the straight, black hair of the proper Americans, are generally a duller-complexioned, shorter, and a more squat people, and they have still more prominent cheek-bones. But the circumstance which most completely separates them from the typical Americans is the form of their skulls, which, instead of being broad, high, and truncated behind, are eminently long, usually low, and prolonged backwards. These Hyperborean people

clothe themselves in skins, know nothing of pottery, and hardly anything of metals. Dependent for existence upon the produce of the chase, the seal and the whale are to them what the cocoa-nut tree and the plantain are to the savages of more genial climates. Not only are those animals meat and raiment, but they are canoes, sledges, weapons, tools, windows, and fire; while they support the dog, who is the indispensable ally and beast of burden of the *Esquimaux*.

It is admitted that the *Tchuktchi*, on the eastern side of Behring's Straits, are, in all essential respects, *Esquimaux*; and I do not know that there is any satisfactory evidence to show that the *Tunguses* and *Samoiedes* do not essentially share the same physical characters. Southward, there are indications of *Esquimaux* characters among the Japanese, and it is possible that their influence may be traced yet further.

However this may be, Eastern Asia, from Mantchouria to Siam, Thibet, and Northern Hindostan, is continuously inhabited by men usually of short stature, with skins varying in colour from yellow to olive; with broad cheek-bones, and faces that, owing to the insignificance of the nose, are exceedingly flat; and with small, obliquely-set black eyes and straight black hair, which sometimes attains a very great length upon the scalp, but is always scanty upon the face and body. The skull, never much elongated, is generally remarkably broad and rounded, with hardly any nasal depression, and but slight, if any, projection of the jaws. Many of these people, for whom the

old name of MONGOLIANS may be retained, are nomades; others, as the Chinese, have attained a remarkable and apparently indigenous civilisation, only surpassed by that of Europe.

At the north-western extremity of Europe the Lapps repeat the characters of the Eastern Asiatics. Between these extreme points, the Mongolian stock is not continuous, but is represented by a chain of more or less isolated tribes who pass under the name of Calmucks and Tartars, and form Mongolian islands, as it were, in the midst of an ocean of other people.

The waves of this ocean are the nations for whom, in order to avoid the endless confusion produced by our present half-physical, half-philological classification, I shall use a newname—XANTHOCHROI—indicating that they are "yellow" haired and "pale" in complexion. The Chinese historians of the Han dynasty, writing in the third century before our era, describe, with much minuteness, certain numerous and powerful barbarians with "yellow hair, green eyes, and prominent noses," who, the black-haired, skew-eyed, and flat-nosed annalists remark in passing, are "just like the apes from whom they are descended." These people held, in force, the upper waters of the Yenisei, and thence under various names stretched southward to Thibet and Kashgar. Fair-haired and blue-eyed Northern enemies were no less known to the ancient Hindoos, to the Persians, and to the Egyptians, on the south and west of the great central Asiatic area; while the testimony of all European antiquity is to the effect

that, before and since the period in question, there lay beyond the Danube, the Rhine, and the Seine, a vast and dangerous yellow or red-haired, fair-skinned, blue-eyed population. Whether the disturbers of the marches of the Roman Empire were called Gauls or Germans, Goths, Alans, or Scythians, one thing seems certain, that until the invasion of the Huns they were largely tall, fair, blue-eyed men.

If anyone should think fit to assume that in the year 100 B.C. there was one continuous Xanthochroic population from the Rhine to the Yenisei, and from the Ural mountains to the Hindoo Koosh, I know not that any evidence exists by which that position could be upset, while the existing state of things is rather in its favour than otherwise. For the Scandinavians, the Germans, the Slavonian and the Finnish tribes, to a great extent; some of the inhabitants of Greece, many Turks, some Kirghis, and some Mantchous, the Ossetes in the Caucasus, the Siahposh, the Rohillas, are at the present day fair, yellow or red haired, and blue-eyed; and the interpolation of tribes of Mongolian hair and complexion, as far west as the Caspian Steppes and the Crimea, might justly be accounted for by those subsequent westward irruptions of the Mongolian stock of which history furnishes abundant testimony. The furthestmost limit of the Xanthochroi north-westward is Iceland and the British Isles; south-westward they are traceable at intervals through Syria and the Berber country, ending in the Canary Islands. The cranial characters of the Xanthochroi are not, at present, strictly definable. The



Scandinavians are certainly long-headed ; but many Germans, the Swiss so far as they are Germanised, the Slavonians, the Finns, and the Turks, are short-headed. What were the cranial characters of the ancient "U-suns" and "Tinglings" of the valley of the Yenisei is unknown.

West and south of the area occupied by the chief mass of the Xanthochroi, and north of the Sahara, is a broad belt of land, shaped like a  $\triangleright$ . Between the forks of the Y lies the Mediterranean; the stem of it is Arabia. The stem is bathed by the Indian Ocean, the western ends of the forks by the Atlantic. The majority of the people inhabiting the area thus roughly defined have, like the Xanthochroi, prominent noses, pale skins, and wavy hair, with abundant beards; but, unlike them, the hair is black or dark, and the eyes usually so. They may thence be called the MELANOCHROI. Such people are found in the British Islands, in Western and Southern Gaul, in Spain, in Italy south of the Po, in parts of Greece, in Syria and Arabia, stretching as far northward and eastward as the Caucasus and Persia. They are the chief inhabitants of Africa north of the Sahara, and, like the Xanthochroi, they end in the Canary Islands. They are known as Kelts, Iberians, Etruscans, Romans, Pelasgians, Berbers, Semites. The majority of them are long-headed, and of smaller stature than the Xanthochroi. It is needless to remark upon the civilisation of these two great stocks. With them has originated everything that is highest in science, in art, in law, in politics, and in mechanical inventions. In their hands, at the present moment, lies the order

of the social world, and to them its progress is committed.

South of the Atlas, and of the Great Desert, Middle Africa exhibits a new type of humanity in the NEGRO, with his dark skin, woolly hair, projecting jaws, and thick lips. As a rule, the skull of the Negro is remarkably long; it rarely approaches the broad type, and never exhibits the roundness of the Mongolian. A cultivator of the ground, and dwelling in villages; a maker of pottery, and a worker in the useful as well as the ornamental metals; employing the bow and arrow as well as the spear, the typical negro stands high in point of civilisation above the Australian.

Resembling the Negroes in cranial characters, the BUSHMEN of South Africa differ from them in their yellowish brown skins, their tufted hair, their remarkably small stature, and their tendency to fatty and other integumentary outgrowths; nor is the wonderful click with which their speech is interspersed to be overlooked in enumerating the physical characteristics of this strange people.

The so-called "Dravidian" populations of Southern Hindostan lead us back, physically as well as geographically, towards the Australians; while the diminutive MINCOPIES of the Andaman Islands lie midway between the Negro and Negrito races, and, as Mr. Busk has pointed out, occasionally present the rare combination of brachycephaly, or short-headedness, with woolly hair.

In the preceding progress along the outskirts of the habitable world, eleven readily distinguishable stocks, or persistent modifications, of mankind

have been recognised. I have purposely omitted such people as the Abyssinians and the Hindoos of the valleys of the Ganges and Indus, who, there is every reason to believe, result from the intermixture of distinct stocks. Perhaps I ought, for like reasons, to have ignored the Min-copies. But I do not pretend that my enumeration is complete or, in any sense, perfect. It is enough for my purpose if it be admitted (and I think it cannot be denied) that those which I have mentioned exist, are well marked, and occupy the greater part of the habitable globe.

to accept a faulty name which has the merit of existence than to burthen it with a faultless newly-invented one.

Under each of these divisions are two columns, one for the Brachycephali, or short heads, and one for the Dolichocephali,<sup>1</sup> or long heads. Again, each column is subdivided transversely into four compartments, one for the "leucous," people with fair complexions and yellow or red hair; one for the "leucomelanous," with dark hair and pale skins; one for the "xanthomelanous," with black hair and yellow, brown, or olive skins; and one for the "melanous," with

## LEIOTRICHI.

Dolichocephali. Brachycephali.  
Leucous.

Leucous.

..... Xanthochroi.....

Leucomelaneous.

..... Melanochroi.....

Xanthomelanous.

*Esquimaux.*      Mongolians.

*Amphinesians.*

Americans.

Melanous.

*Australians.*

ULOTRICHI.

Dolichocephali. Brachycephali.

*Bushmen.*

Negroes. *Mincopies* (?)  
Negritos.

\* \* The names of the stocks known only since the fifteenth century are put into italics. If the "Skrällings" of the Norse discoverers of America were Esquimaux, Europeans became acquainted with the latter six or seven centuries earlier.

In attempting to classify these persistent modifications after the manner of naturalists, the first circumstance that attracts one's attention is the broad contrast between the people with straight and wavy hair and those with crisp, woolly, or tufted hair. Bory de St. Vincent, noting this fundamental distinction, divided mankind accordingly into the two primary groups of *Leiotrichi* and *Ulotrichi*—terms which are open to criticism, but which I adopt in the accompanying table, because they have been used. It is better for science

black hair and dark brown or blackish  
skins.

It is curious to observe that almost all the woolly-haired people are also long-headed; while among the straight-haired nations broad heads preponderate, and only two stocks, the Esquimaux and the Australians, are exclusively long-headed.

One of the acutest and most original

<sup>1</sup> Skulls, the transverse diameter of which is more than eight-tenths the long diameter, are short; those which have the transverse diameter less than eight-tenths the longitudinal, are long.



of ethnologists, Desmoulins, originated the idea, which has subsequently been fully developed by Agassiz, that the distribution of the persistent modifications of man is governed by the same laws as that of other animals, and that both fall into the same great distributional provinces. Thus, Australia, America, south of Mexico, the Arctic regions, Europe, Syria, Arabia, and North Africa, taken together, are each regions eminently characterised by the nature of their animal and vegetable populations, and each, as we have seen, has its peculiar and characteristic form of man. But it may be doubted whether the parallel thus drawn will hold good strictly, and in all cases. The Tasmanian Fauna and Flora are essentially Australian, and the like is true, to a less extent, of many, if not of all, the Papuan islands; but the Negritos who inhabit these islands are strikingly different from the Australians. Again, the differences between the Mongolians and the Xanthochroi are out of all proportion greater than those between the Faunæ and Floræ of Central and Eastern Asia. But whatever the difficulties in the way of the detailed application of this comparison, of the distribution of men with that of animals, it is well worthy of being borne in mind, and carried as far as it will go.

Apart from all speculation, a very curious fact regarding the distribution of the persistent modifications of mankind becomes apparent on inspecting an ethnological chart, projected in such a manner that the Pacific Ocean occupies its centre. Such a chart exhibits an Australian area occupied by dark, smooth-haired

people, separated by an incomplete inner zone of dark, woolly-haired Negritos and Negroes, from an outer zone of comparatively pale and smooth-haired men, occupying the Americas, and nearly all Asia and North Africa.

Such is a brief sketch of the characters and distribution of the persistent modifications, or stocks, of mankind at the present day. If we seek for direct evidence of how long this state of things has lasted, we shall find little enough, and that little far from satisfactory. Of the eleven different stocks enumerated, seven have been known to us for less than 400 years; and of these seven not one possessed a fragment of written history at the time it came into contact with European civilisation. The other four—the Negroes, Mongolians, Xanthochroi, and Melanochroi—have always existed in some of the localities in which they are now found; nor do the Negroes ever seem to have voluntarily travelled beyond the limits of their present area. But ancient history is in a great measure the record of the mutual encroachments of the other three stocks.

On the whole, however, it is wonderful how little change has been effected by these mutual invasions and intermixtures. As at the present time, so at the dawn of history, the Melanochroi fringed the Atlantic and the Mediterranean; the Xanthochroi occupied most of Central and Eastern Europe, and much of Western and Central Asia; while Mongolians held the extreme east of the Old World. So far as history teaches us, the populations of Europe, Asia, and

Africa were, twenty centuries ago, just what they are now, in their broad features and general distribution.

The evidence yielded by Archæology is not very definite, but, so far as it goes, it is to much the same effect. The mound-builders of Central America seem to have had the characteristic short and broad head of the modern inhabitants of that continent. The tumuli and tombs of ancient Scandinavia, of pre-Roman Britain, of Gaul, of Switzerland, reveal two types of skull—a broad and a long—of which, in Scandinavia, the broad seems to have belonged to the older stock, while the reverse was probably the case in Britain, and certainly in Switzerland. It has been assumed that the broad-skulled people of ancient Scandinavia were Lapps; but there is no proof of the fact, and they may have been, like the broad-skulled Swiss and Germans, Xanthochroi. One of the greatest of ethnological difficulties is to know where the modern Swedes, Norsemen, and Saxons got their long heads, as all their neighbours, Finns, Lapps, Slavonians, and South Germans, are broad-headed. Again, who were the small-handed, long-headed people of the “bronze epoch,” and what has become of the infusion of their blood among the Xanthochroi?

At present Palæontology yields no safe data to the ethnologist. We know absolutely nothing of the ethnological characters of the men of Abbeville and Hoxne, but must be content with the demonstration, in itself of immense value, that man existed in Western Europe when its physical condition was widely differ-

ent from what it is now, and when animals existed which, though they belong to what is, properly speaking, the present order of things, have long been extinct. Beyond the limits of a fraction of Europe, Palæontology tells us nothing of man or of his works.

To sum up our knowledge of the ethnological past of man. So far as the light is bright, it shows him substantially as he is now; and, when it grows dim, it permits us to see no sign that he was other than he is now.

It is a general belief that men of different stocks differ as much physiologically as they do morphologically; but it is very hard to prove, in any particular case, how much of a supposed national characteristic is due to inherent physiological peculiarities, and how much to the influence of circumstances. There is much evidence to show, however, that some stocks enjoy a partial or complete immunity from diseases which destroy or decimate others. Thus there seems good ground for the belief that Negroes are remarkably exempt from yellow fever; and that, among Europeans, the Melanochroic people are less obnoxious to its ravages than the Xanthochroic. But many writers, not content with physiological differences of this kind, undertake to prove the existence of others of far greater moment; and, indeed, to show that certain stocks of mankind exhibit, more or less distinctly, the physiological characters of true species. Unions between these stocks, and still more between the half-breeds arising from their mixture, are affirmed to be either infertile or less fertile than those which take place



between males and females of either stock under the same circumstances. Some go so far as to assert that no mixed breeds of mankind can maintain themselves without the assistance of one or other of the parent stocks, and that, consequently, they must inevitably be obliterated in the long run.

Here, again, it is exceedingly difficult to obtain trustworthy evidence, and to free the effects of the pure physiological experiment from adventitious influences. The only trial which, by a strange chance, was kept clear of all such influences—the only instance in which two distinct stocks of mankind were crossed, and their progeny intermarried without any admixture from without—is the famous case of the Pitcairn Islanders, who were the progeny of Bligh's English sailors by Tahitian women. The results of this experiment, as everybody knows, are dead against those who maintain the doctrine of human hybridity, seeing that the Pitcairn Islanders, even though they necessarily contracted consanguineous marriages, thrived and multiplied exceedingly.

But those who are disposed to believe in this doctrine should study the evidence brought forward in its support by M. Broca, its latest and ablest advocate, and compare this evidence with that which the botanists, as represented by a Gaertner, or by a Darwin, think it indispensable to obtain before they will admit the infertility of crosses between two allied kinds of plants. They will then, I think, be satisfied that the doctrine in question rests upon a very unsafe foundation; that the

facts adduced in its support are capable of many other interpretations; and, indeed, that, from the very nature of the case, demonstrative evidence one way or the other is almost unattainable. *A priori*, I should be disposed to expect a certain amount of infertility between some of the extreme modifications of mankind; and still more between the offsprings of their intermixture. *A posteriori*, I cannot discover any satisfactory proof that such infertility exists.

From the facts of ethnology I now turn to the theories and speculations of ethnologists, which have been devised to explain these facts, and to furnish satisfactory answers to the inquiry—what conditions have determined the existence of the persistent modifications of mankind, and have caused their distribution to be what it is?

These speculations may be grouped under three heads—firstly, the Monogenist hypotheses; secondly, those of the Polygenists; and, thirdly, that which would result from a simple application of Darwinian principles to mankind.

According to the Monogenists, all mankind have sprung from a single pair, whose multitudinous progeny spread themselves over the world, such as it now is, and became modified into the forms we meet with in the various regions of the earth, by the effect of the climatal and other conditions to which they were subjected.

The advocates of this hypothesis are divisible into several schools. There are those who represent the most numerous, respectable, and

would-be orthodox of the public, and are what may be called "Adamites," pure and simple. They believe that Adam was made out of earth somewhere in Asia, about six thousand years ago; that Eve was modelled from one of his ribs; and that, the progeny of these two having been reduced to the eight persons who were landed on the summit of Mount Ararat after an universal deluge, all the nations of the earth have proceeded from these last, have migrated to their present localities, and have become converted into Negroes, Australians, Mongolians, &c., within that time. Five-sixths of the public are taught this Adamitic Monogenism as if it were an established truth, and believe it. I do not; and I am not acquainted with any man of science, or duly instructed person, who does.

A second school of Monogenists, not worthy of much attention, attempts to hold a place midway between the Adamites and a third division, who take up a purely scientific position, and require to be dealt with accordingly. This third division, in fact, numbers in its ranks Linnæus, Buffon, Blumenbach, Cuvier, Prichard, and many distinguished living ethnologists.

These "Rational Monogenists," or, at any rate, the more modern among them, hold, firstly, that the present condition of the earth has existed for untold ages; secondly, that at a remote period, beyond the ken of Archbishop Usher, man was created, somewhere between the Caucasus and the Hindoo Koosh; thirdly, that he might have migrated thence to all parts of the inhabited world, seeing that none of them are unattainable

from some other inhabited part, by men provided with only such means of transport as savages are known to possess and must have invented; fourthly, that the operation of the existing diversities of climate and other conditions upon peoples so migrating is sufficient to account for all the diversities of mankind.

Of the truth of the first of these propositions no competent judge now entertains any doubt. The second is more open to discussion, for in these latter days many question the special creation of man; and even if his special creation be granted, there is not a shadow of a reason why he should have been created in Asia rather than anywhere else. Of all the odd myths that have arisen in the scientific world, the "Caucasian mystery," invented quite innocently by Blumenbach, is the oddest. A Georgian woman's skull was the handsomest in his collection. Hence it became his model exemplar of human skulls, from which all others might be regarded as deviations; and out of this, by some strange intellectual hocus-pocus, grew up the notion that the Caucasian man is the prototypic "Adamic" man, and his country the primitive centre of our kind. Perhaps the most curious thing of all is that the said Georgian skull, after all, is not a skull of average form, but distinctly belongs to the brachycephalic group.

With the third proposition I am quite disposed to agree, though it must be recollected that it is one thing to allow that a given migration is possible, and another to admit there is good reason to believe it has really taken place.



But I can find no sufficient ground for accepting the fourth proposition; and I doubt if it would ever have obtained its general currency except for the circumstance that fair Europeans are very readily tanned and embrowned by the sun. Yet I am not aware that there is a particle of proof that the cutaneous change thus effected can become hereditary, any more than that the enlarged livers which plague our countrymen in India can be transmitted; while there is very strong evidence to the contrary. Not only, in fact, are there such cases as those of the English families in Barbadoes who have remained for six generations unaltered in complexion, but which are open to the objection that they may have received infusions of fresh European blood; but there is the broad fact that not a single indigenous Negro exists either in the great alluvial plains of tropical South America, or in the exposed islands of the Polynesian Archipelago, or among the populations of equatorial Borneo or Sumatra. No satisfactory explanation of these obvious difficulties has been offered by the advocates of the direct influence of conditions. And as for the more important modifications observed in the structure of the brain, and in the form of the skull, no one has ever pretended to show in what way they can be effected directly by climate.

It is here, in fact, that the strength of the Polygenists, or those who maintain that men primitively arose, not from one, but from many stocks, lies. Show us, they say to the Monogenists, a single case in which the characters of a human stock have been essentially modified without its being

demonstrable, or, at least, highly probable, that there has been intermixture of blood with some foreign stock. Bring forward any instance in which a part of the world formerly inhabited by one stock is now the dwelling-place of another, and we will prove the change to be the result of migration or of intermixture, and not of modification of character by climatic influences. Finally, prove to us that the evidence in favour of the specific distinctness of many animals, admitted to be distinct species by all zoologists, is a whit better than that upon which we maintain the specific distinctness of men.

If presenting unanswerable objections to your adversary were the same thing as proving your own case, the Polygenists would be in a fair way towards victory; but, unfortunately, as I have already observed, they have as yet completely failed to adduce satisfactory positive proof of the specific diversity of mankind. Like the Monogenists, the Polygenists are of several sects; some imagine that their assumed species of mankind were created where we find them—the African in Africa, and the Australian in Australia, along with the other animals of their distributional province; others conceive that each species of man has resulted from the modification of some antecedent species of ape—the American from the broad-nosed Simians of the New World, the African from the Troglydytic stock, the Mongolian from the Orangs.

The first hypothesis is hardly likely to win much favour. The whole tendency of modern science is to thrust

the origination of things further and further into the background ; and the chief philosophical objection to Adam being, not his oneness, but the hypothesis of his special creation, the multiplication of that objection tenfold is, whatever it may look, an increase, instead of a diminution, of the difficulties of the case. And, as to the second alternative, it may safely be affirmed that, even if the differences between men are specific, they are so small that the assumption of more than one primitive stock for all is altogether superfluous. Surely no one can now be found to assert that any two stocks of mankind differ as much as a chimpanzee and an orang do ; still less that they are as unlike as either of these is to any New World Simian !

Lastly, the granting of the Polygenist premises does not in the slightest degree necessitate the Polygenist conclusion. Admit that Negroes and Australians, Negritos and Mongols, are distinct species, or distinct genera if you will, and you may yet, with perfect consistency, be the strictest of Monogenists, and even believe in Adam and Eve as the primæval parents of all mankind.

It is to Mr. Darwin we owe this discovery ; it is he who, coming forward in the guise of an eclectic philosopher, presents his doctrine as the key to ethnology, and as reconciling and combining all that is good in the Monogenistic and Polygenistic schools. It is true that Mr. Darwin has not, in so many words, applied his views to ethnology ; but even he who "runs and reads" the *Origin of Species* can hardly fail to do so ; and, furthermore, Mr. Wallace and M.

Pouchet have recently treated of ethnological questions from this point of view. Let me, in conclusion, add my own contribution to the same store.

I assume Man to have arisen in the manner which I have discussed elsewhere, and probably, though by no means necessarily, in one locality. Whether he arose singly, or a number of examples appeared contemporaneously, is also an open question for the believer in the production of species by the gradual modification of pre-existing ones. At what epoch of the world's history this took place, again, we have no evidence whatever. It may have been in the older tertiary, or earlier ; but what is most important to remember is that the discoveries of late years have proved that man inhabited Western Europe, at any rate, before the occurrence of those great physical changes which have given Europe its present aspect. And as the same evidence shows that man was the contemporary of animals which are now extinct, it is not too much to assume that his existence dates back at least as far as that of our present Fauna and Flora, or before the epoch of the drift.

But, if this be true, it is somewhat startling to reflect upon the prodigious changes which have taken place in the physical geography of this planet since man has been an occupant of it.

During that period the greater part of the British Islands, of Central Europe, of Northern Asia, have been submerged beneath the sea and raised up again. So has the great Desert of Sahara, which occupies the major part of Northern Africa. The Caspian and the Aral Seas have been one,



and their united waters have probably communicated with both the Arctic and the Mediterranean oceans. The greater part of North America has been under water, and has emerged. It is highly probable that a large part of the Malayan Archipelago has sunk, and that its primitive continuity with Asia has been destroyed. Over the great Polynesian area subsidence has taken place to the extent of many thousands of feet—subsidence of so vast a character, in fact, that, if a continent like Asia had once occupied the area of the Pacific, the peaks of its mountains would now show not more numerous than the islands of the Polynesian Archipelago.

What lands may have been thickly populated for untold ages, and subsequently have disappeared and left no sign above the waters, it is, of course, impossible for us to say; but, unless we are to make the wholly unjustifiable assumption that no dry land rose elsewhere when our present dry land sank, there must be half-a-dozen Atlantises beneath the waves of the various oceans of the world. But, if the regions which have undergone these slow and gradual, but immense alterations, were wholly or in part inhabited before the changes I have indicated began—and it is more probable that they were than that they were not—what a wonderfully efficient “Emigration Board” must have been at work all over the world long before canoes, or even rafts, were invented, and before men were impelled to wander by any desire nobler or stronger than hunger. And as these rude and primitive families were thrust, in the course of long series of generations, from land to

land, impelled by encroachments of sea or of marsh, or by severity of summer heat or winter cold, to change their positions, what opportunities must have been offered for the play of natural selection in preserving one family variation and destroying another!

Suppose, for example, that some families of a horde which had reached a land charged with the seeds of yellow fever varied in the direction of woolliness of hair and darkness of skin. Then, if it be true that these physical characters are accompanied by comparative or absolute exemption from that scourge, the inevitable tendency would be to the preservation and multiplication of the darker and woollier families, and the elimination of the whiter and smoother haired. In fact, by the operation of causes precisely similar to those which, in the famous instance cited by Mr. Darwin, have given rise to a race of black pigs in the forests of Louisiana, a negro stock would eventually people the region. Again, how often, by such physical changes, must a stock have been isolated from all others for innumerable generations, and have found ample time for the hereditary hardening of its special peculiarities into the enduring characters of a persistent modification.

Nor, if it be true that the physiological differences of species may be produced by variation and natural selection, as Mr. Darwin supposes, would it be at all astonishing if in some of these separated stocks the process of differentiation should have gone so far as to give rise to the phenomena of hybridity. In the face of the overwhelming evidence in

favour of the unity of the origin of mankind afforded by anatomical considerations, satisfactory proof of the existence of any degree of sterility in the unions of members of two of the

"persistent modifications" of mankind might well be appealed to by Mr. Darwin as crucial evidence of the truth of his views regarding the origin of species in general.

---

*Later in the present year there will be included in this Series the other Essays of Huxley which are now out of copyright.*



## Works dealing with Evolution.

---

**THE EVOLUTION OF MAN.** By PROFESSOR ERNST HAECKEL. In two vols., royal 8vo, 948 pp., 28 plates, 512 illustrations; 42s. net, carriage extra. Abridged edition, cloth, 2s. net, by post 2s. 5d.; in paper cover, 1s. net, by post 1s. 5d.

**THE WONDERS OF LIFE.** A Popular Study of Biological Philosophy. By the same Author. xiv.-501 pp.; cloth, 6s. net, by post 6s. 4d. Popular edition (similar to 6s. edition, unabridged, but in cheaper binding), cloth, 2s. 6d. net, by post 2s. 10d.

**THE ORIGIN OF SPECIES.** By CHARLES DARWIN. Cloth, 1s., by post 1s. 3d.; in paper cover, 6d., by post 8d.

**AN EASY OUTLINE OF EVOLUTION.** By DENNIS HIRD. Cloth, 1s., by post 1s. 3d.; in paper cover, 6d., by post 8d.

**A PICTURE BOOK OF EVOLUTION.** By the same Author. In two Parts, 2s. 6d. net each, by post 2s. 10d.; the two Parts post free 5s. 6d. Part I. deals with Astronomy, Geology, and Zoology; Part II. with Comparative Anatomy and Embryology, and gives the Pedigree of Man.

**THE STORY OF CREATION.** By EDWARD CLODD. In paper cover, 6d., by post 8d.

---

*Agents for the Rationalist Press Association, Ltd. :*

WATTS & Co., 17, Johnson's Court, Fleet St., London, E.C.

# List of Publications

(Alphabetically Arranged) Issued for the

**Rationalist Press Association, Limited,**

BY

**WATTS & CO.,**

17, JOHNSON'S COURT, FLEET STREET, LONDON, E.C.

(For a complete list of Sixpenny Publications see page 4 of cover.)

---

ANONYMOUS.—**Supernatural Religion:** An Inquiry into the Reality of Divine Revelation. Popular Edition, thoroughly revised by the Author. xvi.—920 pp.; paper covers, 2s. 6d. net, by post 2s. 11d.; cloth, 6s. net, by post 6s. 6d.; half morocco, gilt edges, 10s. net, by post 10s. 6d. Also the original edition, in three volumes, 6s. net, by post 7s.

A really great work, pronounced by Matthew Arnold to be "learned and exact," and by John Morley to be "decisive, trenchant, and far-reaching," the author stating "his case with a force which no previous English writer on the negative side can have the smallest claim to rival."

BITHELL, RICHARD, Ph.D.—**A Handbook of Scientific Agnosticism.** 64 pp.; cloth, 2s.; by post 2s. 3d.

BÜCHNER, PROFESSOR LUDWIG.—**Last Words on Materialism** and Kindred Subjects. Translated by Joseph McCabe. With Portrait of the Author and Biographical Sketch by his brother, Professor Alex Büchner. xxxiv.—299 pp.; 1s. 6d. net, by post 1s. 10d.

CALLAWAY, CHARLES, M.A., D.Sc.—**Does Determinism Destroy Responsibility?** 16 pp.; 2d., by post 2½d.

—**King David of Israel:** A Study in the Evolution of Ethics. Cloth, 2s. 6d. net, by post 2s. 9d.

CLODD, EDWARD.—**Thomas Henry Huxley.** (Modern English Writers' Series.) Cloth, 2s. 6d., by post 2s. 9d.

COLUMBINE, W. B.—**Mr. Balfour's Apologetics Critically Examined.** 232 pp.; cloth, 1s. net, by post 1s. 4d.

"A piece of thorough good work : exhaustive, demolishing, and withal high-toned."—EDWARD CLODD.

CONWAY, MONCURE D.—**Lessons for the Day.** 232 pp., 2s. 6d. net, by post 2s. 10d.

CROSSFIELD, HENRY.—**England and Islam.** 55 pp.; 3d. net, by post 4d.

DIXIE, LADY FLORENCE.—**Towards Freedom.** An Appeal to Thoughtful Men and Women. 14 pp.; 2d., by post 2½d.



EDWARDS, CHILPERIC.—**The Hammurabi Code.** 2s. 6d. net, by post 2s. 10d.

FORESTER, GEORGE.—**The Faith of an Agnostic;** or, First Essays in Rationalism. Cloth, 5s.

GASKELL, G. A.—**Exeunt Mahatmas!** 16 pp.; 2d., by post 2½d.

GLANVILLE, W. (ex-Baptist Minister).—**The Web Unwoven;** or, The Dolus Theory of the Book of Acts, as presented in a Critique of Chapters X., XI., and XII. of same. 3d. net, by post 5d.

GODFREY, W. S.—**Theism Found Wanting.** 2d., by post 3d.

GORHAM, CHARLES T.—**The Ethics of the Great French Rationalists.** 101 pp.; cloth, 1s., by post 1s. 3d.

— **Faith: its Freaks and Follies.** 104 pp.; 6d. net, by post 8d.

— **The Transformation of Christianity.** A Summary of the Principal Conclusions embodied in the *Encyclopædia Biblica*. 16 pp.; 1d., by post 1½d.

GOULD, F. J.—**A Concise History of Religion.** 3 vols. Vol. I., 3s.; Vol. II., 3s. 6d.; Vol. III., 5s.

No work of the same size and dealing with this important theme contains such a mass of information. All the highest authorities have been carefully consulted, and the book gives the main results of Biblical criticism, together with other valuable matter, in what is, by comparison at least, a nutshell. The *First Volume* treats of the superstitions of savages and primitive man, and delineates the characteristics of the religions of America, Finland, China, Egypt, Arabia, Chaldaea, Syria, India, Japan, Persia, the Kelts, Greeks, and Romans. The *Second Volume* takes to pieces the whole of the Old Testament literature, and explains the origin of the various parts. The last chapter describes the Religious Environment of Early Christianity. The *Third Volume* traces the growth of the Christian movement, the lives of Paul and Jesus (with due separation of the mythical elements), and affords a Rationalistic analysis of the whole of the New Testament books.

— **The Children's Plutarch.** With six full-page Illustrations by Walter Crane. viii.—286 pp.; cloth, 2s. 6d. net, by post 2s. 10d.

— **Tales from the New Testament.** 176 pp.; cloth, 1s. net, by post 1s. 3d.

— **Will Women Help?** 100 pp.; 6d. net, by post 8d.

— **The Children's Book of Moral Lessons.** First Series. Cheap Edition; with design by Walter Crane; 128 pp.; paper covers, 6d.; cloth 1s., by post 1s. 3d. Second Series ("Kindness" and "Work and Duty"), 204 pp.; cloth, 2s., by post 2s. 3d. Third Series ("The Family"; "People of Other Lands"; "History of Industry, Art, Science, and Religion"), 203 pp.; cloth, 2s., by post 2s. 3d. Fourth Series ("Justice," "The Common Weal," "Our Country," "Social Responsibilities," "Political and Industrial Progress," etc.), 216 pp.; cloth, 2s., by post 2s. 3d.

— **The Religion of the First Christians.** Beautifully bound, gold lettered, 2s. 6d.

— **Funeral Services, Without Theology.** A Series of Addresses adapted to various occasions. With an Appendix containing (1) examples of method of treating personal recollections, and (2) poetical quotations. 64 pp.; limp cloth, 1s. net, by post 1s. 1½d.

— **The Ethical Riches.** 14 pp.; 2d., by post 2½d.

- GOULD, F. J.—**The Building of the Bible.** Showing the Chronological Order in which the Books of the Old and New Testaments appeared according to recent Biblical Criticism; with Notes on Contemporary Events. 24 pp.; 3d., by post 4d. (Third edition, revised.)
- **Plan of Moral Instruction.** 15 pp. No price is marked on this publication, as it is intended for free distribution. The publishers will send single copies gratis to any applicants, and quantities will be supplied on the following terms: One dozen, 8d.; a hundred, 5s.; a thousand, £1 10s.
- GREGORY, MRS. F. K.—**New Stories for Children.** Boards, 1s. 6d., by post 1s. 10d.
- HARDWICKE, W. W., M.D.—**Sunday Observance:** Its Origin and Meaning. 64 pp.; cloth, 1s., by post 1s. 1½d.; paper covers, 6d., by post 7d.
- HEDDERWICK, J. A.—**Do We Believe?** A Rationalist's Analysis of the *Daily Telegraph* Controversy. 64 pp.; cloth, 1s., by post 1s. 3d.; paper covers, 6d., by post 8d.
- HOLYOAKE, G. J.—**The Origin and Nature of Secularism:** Showing that where Freethought commonly ends Secularism begins. 136 pp.; cloth, 1s. net, by post 1s. 3d.
- **Two Great Preachers:** The Rev. Hugh Price Hughes, the Rev. Dr. Joseph Parker. Or, Appreciation Distinct from Concurrence. Paper wrapper, 3d., by post 3½d.
- **The Logic of Death.** 1d., by post 1½d.
- HOOVER, C. E.—**The Anatomy of Knowledge:** An Essay in Objective Logic. (Part I. The Meanings of Reality and Truth. Part II. The Distinctive Grounds of the Sciences.) 226 pp.; cloth, 3s. 6d. net, by post 3s. 10d.
- HUXLEY, THOMAS HENRY.—**Possibilities and Impossibilities.** With Addenda. Paper wrapper, 2d., by post 2½d.
- JEKYLL, WALTER, M.A.—**The Bible Untrustworthy.** A critical comparison of contradictory passages in the Scriptures, with a view of testing their historical accuracy. Cloth, xii.—284 pp.; 3s. 6d. net, by post 3s. 10d.; paper covers, 2s. net, by post 2s. 3d.
- KALTHOFF, ALBERT.—**The Rise of Christianity.** Translated by JOSEPH McCABE. Cloth, 201 pp.; 2s. 6d. net, by post 2s. 10d.
- LAING, SAMUEL.—**A Modern Zoroastrian.** Cloth, 2s. net, by post 2s. 5d.
- **Problems of the Future, and Essays.** Cloth, 2s. net, by post 2s. 5d.
- LEONARD, WILLIAM A.—**The New Story of the Bible.** 109 pp.; cloth, 2s., by post 2s. 3d.; paper, 1s., by post 1s. 3d.
- McCABE, J. (lately Very Rev. FATHER ANTONY, O.S.F.)—**From Rome to Rationalism;** or, Why I Left the Church. 32 pp.; 3d., by post 4d.
- **The Religion of Woman:** An Historical Study. 207 pp.; 2s. 6d. net, by post 2s. 10d.
- **The Truth about Secular Education:** Its History and Results. 96 pp.; 6d., by post 7d.
- **The Bible in Europe:** An Inquiry into the Contribution of the Christian Religion to Civilisation. Cloth, 224 pp.; 2s. 6d. net, by post 2s. 10d.
- **One Hundred Years of Education Controversy.** 16 pp. and paper covers, 3d., by post 4d.; cheaper edition, 1d., by post 1½d.



- MORIES, A. S.—**Haeckel's Contribution to Religion.** 88 pp.; cloth, 1s., by post 1s. 2d.
- MORTIMER, GEOFFREY.—**The New Morality.** 96 pp.; 6d. net, by post 8d.; cloth, 1s. net, by post 1s. 3d.
- PLUMPTRE, CONSTANCE E.—**On the Progress of Liberty of Thought during Queen Victoria's Reign.** 6d. net, by post 8d.; cloth, 1s. net, by post 1s. 3d.
- R. P. A. ANNUAL, THE.—**First issue (for 1908).** 80 pp.; 6d. net, by post 8d.
- ROBERTSON, JOHN M.—**A Short History of Freethought, Ancient and Modern.** In 2 vols., xiii.-480—xiii.-455 pp.; 21s. net, by post 21s. 7d.
- **Christianity and Mythology.** xviii.-484 pp.; 8s. 6d. net, by post 9s.
- **Pagan Christs:** Studies in Comparative Hierology. xviii.-442 pp.; 8s. 6d. net, by post 9s.
- **A Short History of Christianity.** 400 pp.; cloth, 6s. net, by post 6s. 4d.
- **Letters on Reasoning.** Second, revised and enlarged, edition. xxix.-260 pp.; cloth, 3s. 6d. net, by post 3s. 10d.
- **Pioneer Humanists.** 399 pp.; cloth, 6s. net, by post 6s. 4d.
- **What to Read:** Suggestions for the Better Utilisation of Public Libraries. New edition. 16 large pp.; 2d., by post 3d.
- SALT, HENRY S.—**The Faith of Richard Jefferies.** 16 pp.; 2d., by post 2½d.
- SHELLEY, PERCY BYSSHE.—**The Necessity of Atheism.** Being a reprint of the Original Edition produced in 1811. Edited by Thomas J. Wise and Percy Vaughan. 1s. net, by post 1s. 1d.
- SPILLER, GUSTAV (Compiled by).—**Songs of Love and Duty for the Young.** 80 pp.; 8d. net, by post 9d.  
Comprising 90 songs and two sets of responses—one on ethical ideas and duties, the other on the Sacred Books of the World.
- VARIOUS WRITERS.—**Christianity and Rationalism on Trial.** The Christian Defences (which appeared in the *Clarion*) Answered. 96 pp.; 6d., by post 8d.; cloth, 1s., by post 1s. 3d.
- WATTS, CHARLES.—**The Meaning of Rationalism,** and Other Essays. 210 pp.; cloth, 2s. 6d. net, by post 2s. 10d.
- WHITTAKER, THOMAS.—**The Origins of Christianity.** With an Outline of Van Manen's Analysis of the Pauline Literature. Cloth, xvi.-216 pp.; 3s. 6d. net, by post 3s. 10d.
- WRIGHT, FRANCES.—**On the Nature of Knowledge;** And Other Essays. 6d., by post 8d.
- ZILWA, LUCIAN DE, M.B., B.Sc. (Lond.).—**Reason and Emotion in Relation to Conduct.** A Paper read before the Critical Society of University College, London. 2d., by post 2½d.

